

ACE-M-ALL SANITIZER

A BACTERICIDE — DISINFECTANT

DOSAGE

Level Table-
spoon per
Gallon of
Water

1/2

Active Ingredients:

Sodium Hypochlorite Over 3.25%

1

Sodium Phosphate Expressed as $Na_3PO_4 \cdot 12H_2O$ Over 91.75%

2

5

Inert Ingredients:

Sodium Chloride Under 5.00%

10

Total 100.00%

**CLOSED
DRY PLACE**

ACCEPTED

JUL 15 1968

UNDER THE FEDERAL INSECTIONS
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC PESTS, REGISTERED
ED UNDER NO. 6484-1 SUBJECT
TO ATTACHED COMMENTS.

USDA REG. NO. 6484-1

LBS. NET

ACE CHEMICAL PRODUCTS, INC.

622 W. OREGON STREET MILWAUKEE, WIS. 53204
Phone 272-3087

WARNING - - Harmful if swallowed. May cause burns or damage to the eyes. Do not get on skin or in eyes. If contact with eyes occurs flush with plenty of water for 15 minutes and get medical attention. If contact with skin occurs flush with water. Avoid contamination of food and avoid inhalation of dust.

DANGER: KEEP OUT OF REACH OF CHILDREN

DIRECTIONS FOR USE

1. All equipment should be thoroughly cleaned and rinsed, after the daily processing. Disinfect equipment shortly before it is to be used or at the time specified by local regulations.
2. When used as a bactericide for dairy utensils, milking machines and equipment, use 1 1/2 ounces per 3 1/2 gallons of water. Test frequently to maintain a concentration above 50 ppm. If facilities for testing are not available, use solution containing 200 ppm available chlorine or 3 ounces to each 3 gallons of water. Immerse all utensils for at least two minutes.
3. When used as a disinfectant for equipment in food and beverage plants, use a solution of 1 pound per 40 gallons of water and apply in such a manner that all parts and surfaces are properly reached with solution.
4. For domestic and household, use a solution of 1 ounce per gallon of water for disinfecting glasses, dishes, etc. For cleaning and deodorizing of porcelain and enameled surfaces such as sinks, refrigerators, etc., use a solution of 1 ounce per quart of water.
5. Rinse thoroughly with potable water.



THE RELIANCE CHEMICAL COMPANY

MANUFACTURERS

3301 LAKESIDE AVENUE
CLEVELAND, OHIO 44114
PHONE 771-0424

(LEFT PANEL)

DIRECTIONS

IDOSOL Germicidal Cleaner is its own indicator of Germicidal Activity. As long as faint yellow color remains, solution is germicidally active. When yellow color is gone, solution must be replaced. Care must be taken, when using "complexed Iodine" for cleaning, sanitizing or disinfecting, that the surface, being treated is free from alkaline matter.

This material has penetrating and wetting properties getting into openings providing maximum sanitation. IDOSOL is a concentrate. It must be diluted before using. It must not come in contact with food. IDOSOL solution should be freshly prepared and should not be used in combination with any other cleaner or sanitizer.

RECOMMENDED IDOSOL DILUTIONS

1. 1/2 oz. IDOSOL to 10 gallons cold water for sanitizing: milking machines and equipment, dishes, glassware and cutlery, carbonated beverage equipment. Immerse for a few minutes, then rinse with clean water.

2. 1/2 oz. IDOSOL to 10 gallons cold water for sanitizing: other equipment. Immerse for a few minutes, then rinse with clean water.

3. 1/2 oz. IDOSOL to 2 gallons cold water for disinfecting: cold storage equipment, milk cans, etc. Rinse with clean water.

ACCEPTED