

# KITO L

ACCEPTED  
NOV 2 1972  
UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENT CIDE ACT  
FOR ECONOMIC POISON REG. STER-  
ED UNDER NO. 6520-U SUBJECT  
TO ATTACHED COMMENTS.

#### ACTIVE INGREDIENT

Sodium 2, 4, 5 - Trichlorophenate 38.7%

INERT INGREDIENTS - 61.3%

#### WARNING

#### KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed or inhaled. Causes skin irritation. Goggles and gloves should be worn when handling KITO L. Do not breath vapors. Do not get in eyes, on skin or clothing. Wash contact areas with soap & water. In case of eye contact flush with a large quantity of water and call physician. Rinse empty container thoroughly with water and discard it.

This product is toxic to fish. Treated effluent should not be discharged where it will drain into lakes, streams, ponds, or public water. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on our technical data sheet.


Directions for use of KITO L are given in accompanying Hamblet & Hayes Bulletin titled KITO L

NET CONTENTS 50 GALLONS, 510 LBS.

U. S. D. A. REG. NO. 6520-U

MADE IN U.S.A.

## HAMBLET & HAYES COMPANY SALEM, MASSACHUSETTS



# HAMBLET & HAYES Co.

CHEMICALS

SALEM, MASSACHUSETTS 01970

TECHNICAL  
DATA

## KITO L

KITO L is a fungistat produced especially for the control of fungi in leather manufacturing processes.

### CHARACTERISTICS

Type: Water soluble mixture of phenolic compounds in solvents.  
Appearance: Light brown colored liquid  
pH: 11  
Wt. per gal: 10.25 lbs.  
Stability: Volatile mixture. Keep container closed to avoid evaporation.

### DIRECTIONS FOR USING KITO L

#### Pickling:

----- Mold damage of pickled stock will be prevented by the use of 1 to 5 lbs. of KITO L per 1000 lbs. of stock, depending upon concentration, storage conditions, and nature of pickled stock. KITO L should be added at least 30 minutes after the addition of the Sulfuric Acid.

#### Tanning:

----- KITO L will protect chrome and vegetable tanned stock against mold damage. If blue splitting is done, the protection is extended to the splits which may be held in the moist condition for some time before processing. KITO L should be added near the end of the tanning run. The recommended amount of KITO L is 1 to 4 lbs. per 1000 lbs. of stock, depending upon the climate and length of holding time.

### CAUTION

#### KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed or inhaled. Causes skin irritation. Goggles and gloves should be worn when handling KITO L. Do not breath vapors. Do not get in eyes, on skin or clothing. Wash contact areas with soap and water. In case of eye contact flush with a large quantity of water and call physician. Rinse empty container thoroughly with water and discard it.

#### FISH AND/OR WILDLIFE CAUTIONS

This product is toxic to fish. Treated effluent should not be discharged where it will drain into lakes, streams, ponds, or public water. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this sheet.



**HAMBLET & HAYES Co.**

CHEMICALS

SALEM, MASSACHUSETTS 01970

TECHNICAL  
DATA

-2-

Our recommendations for use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice. The buyer must assume all responsibility, including injury or damage resulting from its misuse as such, or in combination with other materials.

TECHNICAL SERVICE IS AVAILABLE