



**ACCEPTED**

AUG 13 1976

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under  
EPA Reg. No. 652-4

# KAY DEE RABON<sup>®</sup> CATTLE SUPPLEMENT

(CONTAINS RABON<sup>®</sup> ORAL LARVICIDE)

To prevent the development of Horn Flies, Face Flies, House Flies and Stable Flies in the manure of treated cattle.

ACTIVE INGREDIENT:	BY WEIGHT:
2-chloro-1-(2,4,5-trichlorophenyl)vinyl dimethyl phosphate	0.18%*
INERT INGREDIENTS:	99.82%**
	Total 100.00%

\* Each pound contains 792 mg RABON  
\*\* Refers only to ingredients which are not larvicidal

### GUARANTEED ANALYSIS

Total Crude Protein, not less than . . . . .	40.00%
(This includes not more than 15.0% equivalent crude protein from non-protein nitrogen)	
Crude Fat, not less than . . . . .	2.5%
Crude Fiber, not more than . . . . .	5.0%
Calcium (Ca), not more than . . . . .	1.0%
Calcium (Ca), not less than . . . . .	0.5%
Phosphorus (P), not less than . . . . .	0.75%
Salt (NaCl), not more than . . . . .	1.00%
Salt (NaCl), not less than . . . . .	0.50%

### VITAMIN GUARANTEE

Vitamin A . . . . .	50,000 USP Units Per Lb.
Vitamin D-3 . . . . .	10,000 USP Units Per Lb.

### FEED INGREDIENTS

Soybean Meal, Urea, Salt, Vitamin A Acetate, Irradiated Animal Sterol (Source of Vitamin D-3), Hominy, Cane Molasses, Ground Wheat Screenings, Dicalcium Phosphate, DeFluorinated Rock Phosphate.

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS

### CAUTION

Keep out of reach of children. Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water for 15 minutes. If irritation persists, see a physician.

#### ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

### DIRECTIONS FOR USE GENERAL CLASSIFICATION

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Feed KAY DEE RABON CATTLE SUPPLEMENT to feeder cattle, weighing 400 to 1100 lbs., at the rate of 1 pound per head per day. KAY DEE RABON CATTLE SUPPLEMENT is to be blended into the total ration. Start feeding early in the spring before flies begin to appear and continue feeding throughout the summer and into the fall, until cold weather restricts fly activity.

KAY DEE RABON CATTLE SUPPLEMENT may be fed up to slaughter.

KAY DEE RABON CATTLE SUPPLEMENT prevents the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle but is not effective against existing adult flies.

In some cases, supplemental fly control measures may be needed in and around cattle lots and

treated cattle.

<b>ACTIVE INGREDIENT:</b>	<b>BY WEIGHT:</b>
2-chloro-1-(2,4,5-trichlorophenyl)vinyl dimethyl phosphate	0.18%*
<b>INERT INGREDIENTS:</b>	99.82%**
Total	100.00%

\* Each pound contains 792 mg RABON  
 \*\* Refers only to ingredients which are not larvicidal

**GUARANTEED ANALYSIS**

Total Crude Protein, not less than	40.00%
(This includes not more than 15.0% equivalent crude protein from non-protein nitrogen)	
Crude Fat, not less than	2.5%
Crude Fiber, not more than	5.0%
Calcium (Ca), not more than	1.0%
Calcium (Ca), not less than	0.5%
Phosphorus (P), not less than	0.75%
Salt (NaCl), not more than	1.00%
Salt (NaCl), not less than	0.50%

**VITAMIN GUARANTEE**

Vitamin A	50,000 USP Units Per Lb.
Vitamin D-3	10,000 USP Units Per Lb.

**FEED INGREDIENTS**

Soybean Meal, Urea, Salt, Vitamin A Acetate, Irradiated Animal Sterol (Source of Vitamin D-3), Hominy, Cane Molasses, Ground Wheat Screenings, Dicalcium Phosphate, DeFluorinated Rock Phosphate.

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

See reverse side for use directions and other precautions

**50 POUNDS  
NET WEIGHT**

EPA Reg. No. 6552 -9  
 Made in U. S. A.  
 U.S. Patent No. 3,102,842

Manufactured by

**KAY DEE FEED COMPANY**  
**Sioux City, Iowa 51107**

eyes, wash with plenty of water for 15 minutes. If irritation persists, see a physician.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

**DIRECTIONS FOR USE  
GENERAL CLASSIFICATION**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Feed KAY DEE RABON CATTLE SUPPLEMENT to feeder cattle, weighing 400 to 1100 lbs., at the rate of 1 pound per head per day. KAY DEE RABON CATTLE SUPPLEMENT is to be blended into the total ration. Start feeding early in the spring before flies begin to appear and continue feeding throughout the summer and into the fall, until cold weather restricts fly activity.

KAY DEE RABON CATTLE SUPPLEMENT may be fed up to slaughter.

KAY DEE RABON CATTLE SUPPLEMENT prevents the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle but is not effective against existing adult flies.

In some cases, supplemental fly control measures may be needed in and around cattle lots and buildings to control adult house flies and stable flies which can breed not only in cattle manure but in other decaying vegetable matter or silage on the premises.

In order to achieve optimum fly control, KAY DEE RABON CATTLE SUPPLEMENT should be used in conjunction with other good management and sanitation practices.

**STORAGE AND DISPOSAL**

Do not contaminate water or food by storage or disposal. Open dumping is prohibited. Dispose of in an incinerator or landfill approved for pesticide containers, or bury in a safe place.