

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

OCT 27 1989

17/1

Mrs. Herbert
1000
1000
1000

Mrs. Herbert

AGREEMENT - Trenching and Registration of
Ration Mineral Block 6
EPA Registration No. 6000-15
Ration Mineral Block 10
EPA Registration No. 6000-19
Ration Mineral Block 11
EPA Registration No. 6000-20
Ration Block 1
EPA Registration No. 6000-21

Your letters dated August 27, 1989 and January 21, 1989

- 1. The applicant shall...
- 2. The applicant shall...
- 3. The applicant shall...
- 4. The applicant shall...
- 5. The applicant shall...

BEST AVAILABLE COPY

55378:1:CR-44:Herbert:LR-5:KENC0:10/15/89:12/05/89:de:VO:CL

CONCURRENCES							
SYMBOL							
SURNAM							
DATE							

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely yours,

George T. LaRocca
Product Manager (15)
Insecticide-Rodenticide Branch
Registration Division (H7505C)

Enclosure

RECEIVED COPY

ACCEPTED
with COMMENTS
in EPA Letter Dated:

OCT 27

Under the Federal Insecticide,
Fungicide, and Rodenticide Act as
amended, for the pesticide
registered under EPA Reg. No.

6552-14



RABON BLOCK 10

With Rabon Oral Larvacide

To prevent the development of horn flies, face flies, house flies and stable flies in the manure of cattle.

Active Ingredient
Tetrachlorvinphos: 2-chloro-1-(2, 4, 5-trichlorophenyl)
vinyl dimethyl phosphate 1.25%*
Inert Ingredients 98.75%**
..... 100.00%

*RABON Insecticide - Contains 5.625 grams of RABON per lb.
** Refers only to ingredients which aren't larvical

Keep Out Of Reach Of Children
CAUTION

See Side Panel For Additional Precautionary Statements

GUARANTEED ANALYSIS

Calcium (Ca), max.	12.00%
Calcium (Ca), min.	10.00%
Phosphorus (P), min.	10.00%
Salt (NaCl), max.	10.80%
Salt (NaCl), min.	9.00%
Magnesium (Mg), min.	1.00%
Manganese (Mn), min.	10%
Cobalt (Co), min.	0.01%
Copper (Cu), min.	0.1%
Iron (Fe), min.	15%
Vitamin A, min.	300,000 IU/Lb.
Vitamin D3, min.	45,000 IU/Lb.
Vitamin E, min.	150 IU/Lb.

FEED INGREDIENTS

Dicalcium Phosphate, Cane Molasses, Salt, Processed Grain-By-Products, Dried Molasses, Iron Oxide, Bentonite, Sodium Selenite, Iron Carbonate, Zinc Oxide, Manganous Oxide, Copper Oxide, Cobalt Carbonate, Magnesium Oxide, Vitamin A Acetate, D-Activated Animal Sterol (source of Vitamin D3), Vitamin E Supplement & Lignin Sulfonate

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Provide 1 block per 10-15 head of cattle. Add an additional block when each block has been 1/2 consumed. Cattle should consume an average 6 oz. of the block per 100 pounds of body weight per month. This will supply the recommended average intake of 70 mg RABON Oral Larvacide per 100 pounds of body weight. Animals should not be starved for mineral before initiation of the Kay Dee Rabon Block 10 program, but omit all salt from other feed after the program is started. Do not feed free choice sources of salt or mineral containing salt in any form. Place near watering and loafing areas. Protect from rain. If consumption of Kay Dee Rabon Block 10 is above 6 oz. of the block per 100 lbs. of body weight per month, reduce the number of feeding locations or relocate. If consumption is below 6 oz. of the block per 100 lbs. per month increase the

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS

Caution: Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water. If irritation persists, see a physician. Wear long sleeved shirt and pants, chemical resistant gloves, shoes and socks for protection when handling.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not use manure/droppings on land where food crops are to be grown during the current year.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Storage: Store in a dry place in original container. Disposal: Completely empty container into application (mixer/feeder) equipment. Then dispose of empty bag in a sanitary landfill or by incineration or if allowed, by state and local authorities by burning. If burned, stay out of smoke. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

number of feeding locations. Start feeding Kay Dee Rabon Block 10 early in the spring before flies begin to appear, and continue feeding throughout the summer and into fall.

Kay Dee Rabon Block 10 may be fed up to slaughter and to lactating dairy cows.

Kay Dee Rabon Block 10 prevents the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle, but is not effective against existing adult flies. When starting a feeding program during the fly season, it may be necessary to use other control measures to reduce the population of existing adult flies. Supplemental fly control measures may be needed in and around cattle lots and buildings to control adult house flies and stable flies which can breed not only in cattle manure, but in other decaying vegetable matter or silage on the premises. In order to achieve optimum fly control the Kay Dee Rabon Block 10 should be used in conjunction with other good management and sanitation practices.

NET WEIGHT 50 POUNDS

Manufactured by
KAY DEE FEED COMPANY
Sioux City, Iowa 51167

8-89
EPA REG NO 6552-14
EPA EXT NO 6552-1A-1 LOT NO

BEST AVAILABLE COPY