

SEC - 2



6552-17

RABON® BLOCK 4
With Rabon® Oral Larvicide

NET WEIGHT 331/3 POUNDS

LOT NO. _____

To prevent the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle.

ACTIVE INGREDIENT	BY WEIGHT
* 2-chloro-1 (2,4,5-trichlorophenyl) vinyl dimethyl phosphate	0.473%
INERT INGREDIENTS**	99.527%
Total	100.000%

* Each pound contains 2.14 g. RABON®
** Refers only to ingredients which are not larvicidal.

GUARANTEED ANALYSIS

Calcium (Ca), max.	7.20%
Calcium (Ca), min.	6.00%
Phosphorus (P), min.	4.00%
Salt (NaCl), max.	19.20%
Salt (NaCl), min.	16.00%
Total Sugars (as invert), min.	17.00%
Vitamin A, min.	60,000 USP Units/Lb.
Vitamin D ₃ , min.	5,000 USP Units/Lb.

FEED INGREDIENTS

Cane Molasses, Soybean Mill Feed, Dicalcium Phosphate, Salt, Calcium Carbonate, Dried Molasses, Manganous Oxide, Iron Sulfate, Magnesium Oxide, Zinc Oxide, Copper Oxide, Cobalt Carbonate, Ethylenediamine Dihydrochloride, Sodium Selenite, D-Activated Animal Sterol (source of Vitamin D₃), Vitamin A Acetate.

DIRECTIONS FOR USE

It is a violation of Federal law to use the product in a manner inconsistent with its labeling.
Provide 1 block per 5 head of cattle. Add an additional block when each block has been 1/2 consumed. Fed at a daily rate of .07 grams of larvicide in 0.5 ounces of block per 100 pounds of body weight. Example: 5 ozs. (approximately 1/2 pound) of block for a 1000 pound animal daily. This will supply the recommended average intake of 70 mg. RABON® ORAL LARVICIDE per 100 lbs. of body weight. Do not feed salt or mineral supplements of any kind containing salt. Place blocks near water and loafing areas. Protect from rain. If consumption of ROYAL RABON® BLOCK 4 is above the recommended level, decrease the number of feeding stations and or raise the block per head ratio. Start feeding ROYAL RABON® BLOCK 4 early in the spring before flies begin to appear and continue feeding throughout the summer and into the fall. ROYAL RABON® BLOCK 4 may be fed up to slaughter and to lactating dairy cows.
ROYAL RABON® BLOCK 4 prevents the development of horn flies, face flies, house flies and stable flies in the manure of

KEEP OUT OF REACH OF CHILDREN
CAUTION

See below for additional precautionary, storage and disposal statements.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS

Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water. If irritation persists, see a physician.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL

Pesticides that cannot be used according to label instructions must be disposed of according to applicable Federal, State or Local procedures.

CONTAINER DISPOSAL

Completely empty bag by shaking & tapping sides & bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of bags in a sanitary landfill or by incineration if allowed by State and Local authorities.

treated cattle but is not effective against existing adult flies. Supplemental fly control measures may be required in and around cattle lots and buildings to control adult house flies and stable flies which can breed not only in cattle manure, but in other decaying vegetable matter or sludge on the premises. Because face flies migrate, work with your neighbors to more effectively control the adult face fly population.

In order to achieve optimum fly control, ROYAL RABON® BLOCK 4 should be used in conjunction with other good management and sanitation practices.

Manufactured by
KAY DEE FEED COMPANY
Sioux City, Iowa 51107

EPA REG. NO. 6552-17
EPA EST. NO. 6552-1A-1

3/3/88

® Rabon is the registered trademark of Shell Oil Co., U.S. Pat. 3,102,842