

USE DIRECTIONS

Use 2 ounces *ILF Fabric Softener Sanitizer (Powder)* per 100 lbs. of fabric in last cycle (rinse or sour) to impart light softening (250 ppm active softener) and bacteriostasis (125 ppm active quaternaries) against *S. aureus* and *B. ammoniagenes**

OR

Use 4 ounces *ILF Fabric Softener Sanitizer (Powder)* per 100 lbs. of fabric in last cycle (rinse or sour) to impart more softening (500 ppm active softener) and residual self-sanitization (250 ppm active quaternaries).

Fabric treated at 2 ounces per 100 lbs. dry weight (125 ppm active quaternaries) will develop bacteriostatic activity against *Staphylococcus aureus* and *Brevibacterium ammoniagenes**

Fabric treated to 4 ounces per 100 lbs. dry weight (250 ppm active quaternaries) will develop residual self-sanitizing activity against Gram positive organisms (*S. aureus* and *B. ammoniagenes**).

ILF Fabric Softener Sanitizer (Powder) softens clothes, thus eliminating chafing, "static cling" from nylons and other synthetics. Improves ironing time.

*Under conditions of high relative Humidity or wet contamination.

(This product is not effective as a bacteriostat against Gram negative organisms.)

ILF FABRIC SOFTENER SANITIZER

(POWDER)

FOR INDUSTRIAL AND COMMERCIAL
LAUNDRY USE ONLY

Active Ingredients:

n-Alkyl (60% C₁₄, 30% C₁₆, 5% C₁₂

5% C₁₈ dimethyl benzyl ammonium

chlorides 10.0%

Inert Ingredients 90.0%

100.0%

NET CONTENTS: GALS.

EPA Reg. No. 6609-6

EPA Est. Reg. No. 6609IL01

INDUSTRIAL LAUNDRY FORMULATIONS, INC.

10421 Franklin Avenue

Franklin Park, Illinois 60131

DANGER

KEEP OUT OF REACH OF CHILDREN

See right side panel for additional
precautionary and first aid statements.

DANGER

KEEP OUT OF REACH OF CHILDREN. Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin, or on clothing. Protect eyes and skin when handling product. Harmful if swallowed. Avoid contamination of food.

FIRST AID

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution, or, if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN:

Probable mucosal damage may contra indicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

Rinse empty container thoroughly with water and discard it.

Revised 9/25/74