

Residual Bacteriostatic and Residual Self-Sanitizing Activity.*

Fabric treated with *ILF 50 QUAT* as directed below will develop residual bacteriostatic and residual self-sanitizing activity against Gram Positive (*Staph aureus*) and Gram negative *K pneumoniae* organisms*.

Fabric treated with *ILF 50 QUAT* which has become contaminated or soiled through use will inhibit the growth* of many organisms not only at the time of exposure, but also through the handling and transportation procedures until the garments are again laundered and treated with *ILF 50 QUAT*.

ILF 50 QUAT is particularly effective in inhibiting the growth of ammonia producing organisms such as *Brevibacterium ammoniagenes** which acts on urine in the diapers to produce ammonia.

USE DIRECTIONS

ILF 50 QUAT is recommended for use in the final rinse cycle of the laundry operation. Introduce *ILF 50 QUAT* into the wash wheel at the rate of 1-1/8 ounce of *ILF 50 QUAT* per 100 pounds of dry laundry. It is recommended that *ILF 50 QUAT* be added halfway through the final rinse cycle filling operation. Rinse empty container thoroughly with water and discard container.

*Under conditions of high relative humidity or wet contamination.

ILF 50 QUAT

CONCENTRATED GERMICIDE

INDUSTRIAL LAUNDRY FABRIC USE

Active Ingredients:

n-Alkyl (C₁₂ 40%, C₁₄ 50%
C₁₆ 10%) Dimethyl Benzyl
Ammonium Chlorides . . . 50%


Inert Ingredients 50%

Total 100%

Net Contents:

EPA Reg. No. 6609-8

MANUFACTURED FOR:



INDUSTRIAL LAUNDRY FORMULATIONS, INC.

10421 Franklin Avenue

Franklin Park, Illinois 60131

DANGER

KEEP OUT OF REACH OF CHILDREN
See right side panel for First Aid and precautionary statements.

DANGER

KEEP OUT OF REACH OF CHILDREN. Corrosive. Causes severe eye and skin damage. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful or fatal if swallowed. Avoid contamination of food.

FIRST AID

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN:

Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock as well as oxygen and measures to support breathing manually or mechanically may be needed. If persistent, convulsions may be controlled by the cautious intravenous injection of a short-acting barbiturate drug.

ACCEPTED WITH COMMENTS

ACCEPTED