

FRONT PANEL

INDUSTRIAL AEROSOL
with 7% VAPONA* insecticide and Ronnel
FOR TOTAL RELEASE OR SPOT SPRAYING

(14 point type) DANGER: KEEP OUT OF REACH OF CHILDREN


Picture of skull and POISON
cross bones.

(10 point type) See antidote statement and other cautions on side
panel.

Kills exposed stages of a wide variety of insects including: Roaches,
Ticks, Fleas, Flies and Silverfish in Homes and Buildings.

Net Weight 6 Oz. Avoir.

CENTRAL EXTERMINATING & CHEMICAL CO.
365 CENTRAL PARK AVENUE
YONKERS, N. Y. 10704



IF WARNING SYMPTOMS APPEAR: Administer two 1/100 grain Atropine tablets immediately. NEVER ADMINISTER ATROPINE UNLESS WARNING SYMPTOMS APPEAR. (See warning symptoms below). Keep patient prone and quiet. Start artificial respiration immediately if patient is not breathing. Transport the patient immediately to the nearest physician.


NOTE TO PHYSICIAN:

WARNING SYMPTOMS: Symptoms include weakness, headache, tightness in chest, blurred vision, non-reactive pinpoint pupils, sweating, nausea, vomiting, diarrhea and abdominal cramps.

TREATMENT: Atropine-is the specific therapeutic antagonist of choice against parasympathetic nervous stimulation. If there are signs of parasympathetic stimulation, atropine-sulfate should be injected at 10 minute intervals, in doses of 1 to 2 milligrams until complete atropinization has occurred. Morphine is contraindicated. Clear chest by postural drainage. Observe patient continuously for 48 hours. Repeated exposure to cholinesterase inhibitors may, without warning, cause prolonged susceptibility to very small doses of any cholinesterase inhibitor. Allow no further exposure until time for cholinesterase regeneration has been allowed as determined by blood test.

PRECAUTIONS IN USING: This product is poisonous, therefore precautions must be observed when handling. (See antidote and warnings.) Store in a safe place. Children or animals must not be able to reach or handle. Care should be used in handling empty containers. Do not contaminate food, water or foodstuffs.


DISPOSAL OF CONTAINERS: Completely empty container by removing container to an open air area away from pets and animals. Press release button until no more vapor escapes. Puncture can to prevent reuse. Bury can 18 inches deep in an isolated location away from water supplies.



Comments

BACK PANEL

U. S. D. REG. NO.



SOLD TO PROFESSIONAL PEST CONTROL OPERATORS ONLY

GENERAL DIRECTIONS: To be used by licensed pest control operators only. Use one 6 oz. can for each 5,000 cubic feet of unobstructed closed area to be treated. During fogging and to release contents, use a mask or respirator of a type passed by the U. S. Dept. of Agriculture for DDVP insecticide protection. Shut off air conditioners and fans before using.

All occupants must be removed from house or building before treatment is started and should not re-occupy building for at least 2 hours. Open doors and windows and allow to be ventilated before re-occupying. Unwrapped or unpackaged foods should be removed or placed in refrigerator during fogging. Food handling equipment and dishes should be washed after application. Do not contaminate water, food or foodstuffs. Remove all pets. Cover fish tanks or remove. Cover house plants within 8 feet of point of release with newspaper to prevent possible burning of tender plants.

Place canister on table in center of area to be treated, or near point of greatest infestation. Place newspapers under canister. Use one or more canisters, depending upon the size of the area to be treated.

Snap lock-open nozzle over lip of container. Leave building at once. The entire contents will release automatically.

WARNING: This product is poisonous if swallowed, inhaled or absorbed through skin or eyes. Do not get on skin or eyes. Do not breathe fumes. Wear natural rubber gloves, protective clothing and goggles when handling. Wear a mask or respirator of a type passed by the U. S. Dept. of Agriculture for DDVP (Vapona) insecticide protection. Keep all unprotected persons out of operational area. In case of contact wash immediately with soap and water. Wash hands, arms and face thoroughly with soap and water before smoking or eating. Wash all contaminated clothing with soap and hot water before reuse.

Contents under pressure. High temperature may burst container. Do not expose to direct sunlight, radiators, stove, hot water or other sources of heat.

ANTIDOTE AND FIRST AID TREATMENT: ATROPINE is the emergency antidote for Vapona insecticide poisoning. Consult your physician about an adequate supply of 1/100 grain Atropine tablets for emergency use. CALL A PHYSICIAN IMMEDIATELY IN ALL CASES OF SUSPECTED POISONING.

INTERNAL: If the material has been swallowed induce vomiting immediately. This may be done by introducing a finger into the throat or by giving warm salty water (1 tablespoonful of salt to a glass of water). NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.


EXTERNAL: If the material has been spilled on the skin, immediately remove patient from the vicinity of the insecticide, remove all contaminated clothing and wash skin with soap and running water. If the material gets into eyes, wash immediately with running water for at least ten minutes.

ACCEPTED WITH COMMENTS

ACTIVE INGREDIENTS:

2,2-Dichlorovinyl Dimethyl Phosphate (DDVP)*	6.5%
Related Compounds*	0.5%
Ronnel (O,O-dimethyl O-(2,4,5-trichlorophenyl) Phosphorothioate	2.00%
Petroleum Distillates	4.00%
Xylol	2.00%
INERT INGREDIENTS:	<u>85.00%</u>
	100.00%

*Equivalent to 7% VAPONA insecticide



ACCEPTED WITH COMMENTS