



# Pyrenone® AQUEOUS Space Spray

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

### INDOOR APPLICATION

The use of this product in food processing establishments should be confined to time periods when the plant is not in operation. Food should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. De-energize electrical equipment prior to application. Allow ample time for material to evaporate and dry before starting motors or placing switches in "ON" position.

For maximum effectiveness, a combination of spot surface treatment and fogging is recommended.

**SPACE SPRAYING:** Close doors and windows and shut off ventilating systems. Apply with a good machine adjusted to deliver a fine mist or fog. For rapid control of house flies, fruit flies, gnats, mosquitoes, skipper flies, wasps, hornets, Anopheles grain moths, Mediterranean flour moths, India meal moths, tobacco moths and exposed or accessible stages of the crawling insects listed under **SURFACE SPRAYING:** Direct the spray at an upward angle distributing it uniformly throughout the entire area at a rate of 3-6 ounces per 1000 cubic feet of space (one gallon to 20,000 to 40,000 cubic feet of space).

**SURFACE SPRAYING:** To control cockroaches, ants, silverfish, spiders, crickets, clover mites, cheese mites, granary weevils, rice weevils, confused flour beetles, rust red flour beetles, sawtoothed grain beetle, spider beetle, cigarette beetles, drug-store beetles, meal worms, grain mites and caddis: Use a good sprayer adjusted to a coarse wet spray. Direct the spray into hiding places, cracks and crevices, under pallets, around containers of stored foods, around the base of machinery, behind shelves and drawers. For silverfish spray book-cases. For ants spray trails, nests and points of entry. If surface application only is to be used, spray floors and other surfaces applying at a rate of 1 gallon to 750 square feet of surface.

To control carpet beetles: Spray edges of carpeting and under carpeting and rugs. Make localized applications to floor and baseboards. Spray directly into cracks, closets and infested areas of shelving. Repeat treatment as needed.

To control bedbugs: Spray mattresses lightly particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, moldings and floors. Repeat treatment as needed.

To control flea and brown dog ticks: Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, moldings, window and door frames, and entire areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Concurrent treatment of animals with an approved insecticide is recommended. Repeat treatment as needed.

To control clothes moths: Remove any infested articles from storage, brush thoroughly and air for several hours in sunlight, if possible. Apply spray liberally to empty chests and closets, bureaus and other storages directing the spray into cracks, joints and crevices. After airing, infested articles may be treated lightly keeping nozzle at least three feet from fabric to avoid staining.


### OUTDOOR APPLICATION

**ADULT MOSQUITO CONTROL:** Apply as a fine mist to thoroughly treat the entire area. For best results, treat when the wind velocity does not exceed 5 miles per hour. Treat shrubbery or vegetation where mosquitoes may rest, low marshy places, stagnant pools or ponds and shore lines. Apply at a rate of 13 fluid ounces per acre (0.3 fluid ounces per 1000 square feet).

Using truck-mounted conventional misting equipment traveling at 5 m.p.h. to treat a 300 foot swath calibrate equipment to deliver 20 fluid ounces per minute.

SEE REVERSE SIDE

BATCH



ACCORDANCE

FEB 15 1985

6626-445

STO

Do not consume storage or PESTICIDE Pesticide, that cannot instruction to applica procedures CONTAINER Triple rin for recycl punctur. a fill, or o procedures

PRE HAZARDS T

Harmful if skin. Avoid Avoid contact of plenty of if irritat

Do not app food handl any food a during app processing ing in bake plants, wa ment, bench exposed for all food p should be thoroughly

Remove peti aquariums

This produ apply direc 75% mersl estimate w disposal of

Buyer assu handling of accordance

CONT

EPA Reg. No. 6626-44

EPA Est. No. 6626 NY-1

A Concentrated PYRENONE® Micro Emulsion

Effective, Fast Acting Insecticide

For Use in Home - Institutions, Restaurants, Food Processing Plants, and Outdoors for Mosquito Control

Apply With Conventional Mechanical or Compressed Air Equipment

### ACTIVE INGREDIENTS

|                               |        |
|-------------------------------|--------|
| Pyrethrins                    | 0.3%   |
| Piperonyl butoxide, Technical | 3.0%   |
| INERT INGREDIENTS             | 96.7%  |
|                               | 100.0% |

\*Equivalent to min. 2.4% (butylcarbityl) (6-propylpiperonyl) ether and 0.6% related compounds.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See reverse side for additional precautions.

DO NOT TRANSPORT OR STORE UNDER 32° F

PYRENONE - Registered trademark of Fairfield American Corporation

MANUFACTURED BY

ATLANTIC RESEARCH LABORATORIES CORPORATION

29-05 - 40th ROAD

Stillwell 4-8825

LONG ISLAND CITY, N. Y. 11101

BEST AVAILABLE COPY