



AQUEOUS Pyrenone[®] FOOD PLANT FOGGING INSECTICIDE

70/1
6626-45
PM 17

EPA Reg. No. 6626-4-5

EPA Est. No. 6626-NY-7

A Concentrated PYRENONE[®] Micro Emulsion
- Clear

- Stable

- Leaves No Oily or Objectional Residue

Designed for Use in Edible Produce Areas of Official Establishments Operating Under the Meat, Poultry, Shell Egg Grading and Egg Products Inspection Programs and in Other Food Processing Plants.

Apply with Conventional Mechanical or Compressed Air Equipment Designed and Adjusted to Deliver a True Aerosol Mist.

ACTIVE INGREDIENTS:

Pyrethrins.....	0.5%
*Piperonyl Butoxide, Technical.....	5.0%
INERT INGREDIENTS:	92.5%
Petroleum Distillate.....	2.0%
	<u>100.0%</u>

*Equivalent to 4.0% (butylcarbityl)(6-propyl-piperonyl) ether and 1.0% related compounds

PYRENONE - Registered trademark of Fairfield American Corporation

**KEEP OUT OF REACH OF CHILDREN.
CAUTION**

See side panel for additional precautionary statements

DO NOT TRANSPORT OR STORE UNDER 32°F

NET CONTENTS _____

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

The use of this product in food processing establishments should be confined to time periods when the plant is not in operation. Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using.

Close windows and doors and shut off ventilating systems. Use a non-thermal fogger or vaporizer adjusted to deliver an aerosol spray (no droplets over 50 microns in diameter and 80% less than 30 microns). Apply at the rate of 1 ounce per 1000 cubic feet of space. Direct the spray toward the ceiling and upper corners of the area and behind obstructions. Keep the area closed for at least 15 minutes. Vacate the treated area and ventilate before reoccupying. Repeat treatment as necessary.

The above described application will kill accessible, exposed stages of flies, including fruit flies, mosquitoes, silverfish, flying moths, gnats, gnats, cockroaches, spiders, scorpions, wasps, hornets, clover mites, cheese mites, cheese skippers and these common stored product pests: granary weevils, rice weevils, confused flour beetles, sawtoothed grain beetles, spider beetles, cigarette beetles, drugstore beetles, Angoumois grain moths, Mediterranean flour moths, Indian meal moths, tobacco moths, meal moths and grain mites.

IN MUSHROOM PRODUCTION AND PROCESSING:

To control adult Mushroom flies, fungus gnats and Cecid flies in growing houses, apply at a rate of 1 ounce per 1000 cubic feet of space. Fogger should be operated 8 to 10 feet away from beds and directed slightly upward over them. Circulating fans should be left in operation, but for best results refrigeration should be turned off. In packing and canning plants use at a rate of 1 ounce to 2000 cubic feet of space.

DISPOSAL

Do not use empty container. Wrap container and put in trash collection.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Avoid contamination of feed and foodstuffs.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply only as specified on the label.

Buyer assumes all risks of use, storage or handling of this product not in strict accordance with directions given herewith.

BEST AVAILABLE COPY

ACCEPTED
APR 1 1985
U.S. the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the product's registered under EPA Reg. No. 6626-45

MANUFACTURED BY
ATLANTIC RESEARCH LABORATORIES CORPORATION

29-05 - 401K ROAD
STUYVESANT 4-0025
LONG ISLAND CITY, N. Y. 11101