


1/4 6662-10 PM-17



CAUTION

Harmful if swallowed. Food should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before use. When using the product in these areas, apply only when the facility is not in operation.

Remove pets, birds and cover fish aquariums before spraying.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

1. PROHIBITION
DO NOT CONTAMINATE WATER, FOOD, OR FEED BY STORAGE OR DISPOSAL. OPEN DUMPING IS PROHIBITED.

2. PESTICIDE DISPOSAL
PESTICIDE, SPRAY MIXTURE, OR RINSAF THAT CANNOT BE USED OR CHEMICALLY PROCESSED SHOULD BE DISPOSED OF IN A LANDFILL APPROVED FOR PESTICIDES OR BURIED IN A SAFE PLACE AWAY FROM WATER SUPPLIES.

3. CONTAINER DISPOSAL - MORE THAN 10 GALLON DRUMS
(a) RESEAL CONTAINER AND OFFER FOR RECONDITIONING OR
(b) TRIPLE RINSE (OR EQUIVALENT) AND OFFER FOR RECYCLING, RECONDITIONING, OR DISPOSAL IN APPROVED LANDFILL OR BURY IN A SAFE PLACE.

CONTAINER DISPOSAL - LESS THAN 10 GALLON DRUMS
TRIPLE RINSE (OR EQUIVALENT) AND OFFER FOR RECYCLING, RECONDITIONING, OR DISPOSAL IN APPROVED LANDFILL OR BURY IN A SAFE PLACE.

4. GENERAL
CONSULT FEDERAL, STATE OR LOCAL DISPOSAL AUTHORITIES FOR APPROVED ALTERNATE PROCEDURES.

FOR INDUSTRIAL USE ONLY



CONCENTRATES

ACTIVE INGREDIENTS:

Pyrethrins	0.50%
Piperonyl butoxide, technical*	2.50%
Petroleum distillate	97.00%
TOTAL	100.00%

*Equivalent to 2.00% (butylcarbyl) (6-propylpiperonyl) ether and 0.50% related compounds.

Dalco Concentrate S has been developed with the aid of some of the leading entomologists in the country to produce a vastly superior insecticide to satisfy the demands of all types of food processing and manufacturing plants. This is an exceptionally concentrated and potent insecticide, adapted especially for use in aerosol producing equipment, fogging devices, thermal units as well as electric and hand spraying equipment. Due to its high concentration, requiring the use of less material and time per job, this proves to be, in all reality an extremely economical product. For use in all types of food processing plants such as Canneries, Producers of Milk and Dairy Products, Beverage and Brewery plants, Meat Packing Industry and Sausage Industry, Baking and Milling Industry, Nut Processors as well as Industrial Plants, Catereries, Institutions, Warehouses and all places where effective insect control is desirable. Harmless to equipment and utensils; Will not impart any color, odor or taste to products and equipment when used as directed; Kills Roaches, Water Bugs, Silver Fish, Flies, Spiders, Vinegar and Fruit Flies, Fleas, Gnats, Ants, Carpet Beetles, accessible Grain Beetles, Flour Weevils, Meal Moths, Rice Weevils, Gallees, Flat Grain Beetles and Chocolate Moths when used as directed.

KEEP OUT OF REACH OF CHILDREN

CAUTION

IF SWALLOWED DO NOT INDUCE VOMITING. CALL A PHYSICIAN IMMEDIATELY.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS



SMALL FLYING INSECTS

Directions for use with space spraying devices:

Close all doors, windows and other spaces through which fog may escape. Fog space treated, using one ounce DALCO CONCENTRATE S for each 2,000 cubic feet of space. Allow fog to remain in closed room for a period of not less than ten to twenty minutes, longer where possible. Air out room, sweep up and destroy fallen insects.

IN MUSHROOM PRODUCTION AND PROCESSING:

To control adult Phorids, Sciarids, and cecidomyiids in growing houses, apply at a rate of 1 ounce per 1,000 cubic feet of space. Fogger should be operated 8 to 10 feet away from beds and directed slightly upward over them. Circulating fans should be left in operation, but for best results, refrigeration should be turned off. In packing and canning plants, use at a rate of 1 ounce to 2,000 cubic feet of space.

IMPORTANT: Effective control by space spraying requires treatment as frequently as conditions on premises necessitate. Frequency of treatment must be based on prevailing conditions, such as, initial infestation, type and amount of product attracting various insects, prevailing temperature, humidity and amount of incoming traffic.

- Directions for Contact spraying with hand or electric sprayer:**
- 1. Roaches, Water Bugs, Silver Fish and Spiders:** Apply at rate of 1-4 ounces per 1,000 cubic feet. Forcefully and thoroughly spray inside all cracks and crevices where moisture collects such as drains, water pipes and sinks. Mouldings and wood work should also be treated when necessary. Wherever possible, spray directly onto insects as they emerge from hiding places which are being sprayed. Repeat treatment as often as necessary.
 - 2. For all accessible stages of Granary insects, such as Flour Beetles, Granary Weevils, Saw-Toothed Grain Beetles, Lesser Grain Borer, Larger Grain Borer, Angoumois Grain Moth, Mediterranean Flour Moth, Indian Meal Moth, accessible Grain Beetles:** Close all doors and windows of area treated. Spray floors, side walls, corners, and cross beams. Stack goods with as much space between stacked merchandise. Grain bins should be emptied as frequently as possible so that walls, floors, corners, etc., are covered with thick spray. Spray weekly in warm weather and at least monthly during winter. Spray liberally after hot periods when migrations start.
 - 3. Flies, Fruit and Vinegar Flies, Manquitos:** Apply at rate of 1 ounce per 2,000 cubic feet. Close doors and windows, direct spray toward ceiling and fill entire room with fog. Keep room closed for ten to twenty minutes. Sweep up and destroy fallen insects. Repeat treatment as often as required.
 - 4. Ants, Flies, Carpet Beetles:** Spray floors, rugs and side walls for several feet up from the floor. Also spray animal quarters. Do not spray animals and pets. Repeat as required.

ACCEPTED

JUN 16 1981

U.S. Environmental Protection Agency, Registration Division, Washington, D.C. 20460

EPA Reg. No. 6662-17