

[6662-20]

DIRECTIONS FOR USE

FOR FLOORING: Use approximately 1 ounce of material for each 1,000 cubic feet of space. Keep doors and windows shut for a minimum of 10 minutes. Sweep up and destroy fallen insects. Follow recommendations of equipment manufacturers for proper positioning of sprayers. Repeat as necessary. DO NOT apply the product in poultry houses.

To obtain RESIDUAL REPELLENCY: Appl. 1 to 2 ounces Barn Spray and Repellent as a mist to each animal for a three-day period. Apply with a good hand sprayer so that the animal's body has been uniformly covered. Special care should be used in spraying the legs, belly and shoulders. Spray in the same manner again after 2 days have elapsed and every second day thereafter, or as necessary. Wash teats and udder with warm water before milking. Calves should be sprayed very lightly daily, instead of attempting to build up a residual repellency by spraying more heavily. Avoid areas around the nose, eyes and mouth.

FOR CONTROL OF LICE ON LIVESTOCK: Spray animal with 2 ounces of spray as a mist or immediately rub spray against lay of hair, paying

ACCEPTED
6-20-72
6662-20

DALCO

BARN SPRAY AND REPELLENT



For use on horses, ponies, donkeys, mules and cattle to kill and repel horn flies, face flies, stable flies, deer flies, mosquitoes and gnats. When properly applied, the repellent effect against above insects will last two days.

ACTIVE INGREDIENTS 100%

- Petroleum Distillates
- N-octyl bicycloheptene dicarboximide
- Di-n-propyl isocinchomeronate
- Piperonyl Butoxide technical
- Pyrethrin

*Consists of (butylcarbityl) (6-propylpiperonyl) ether and related compounds.

CAUTION: Keep Out Of Reach Of Children. Harmful if swallowed. Avoid contact with eyes and mouth. See side panel for additional cautions.

*Animal Products Division
Dalco Corporation*

CLEVELAND, OHIO

NET CONTENTS

particular attention to infested areas. Repeat as necessary, but not more frequently than once every 3 weeks.

TO PROTECT LIVESTOCK FROM FLEAS: Spray bedding and walls of pen, particularly cracks and crevices. Repeat treatment frequently until adequate control is obtained.

Horse sense for animal health - have animals examined periodically by a veterinarian.

CAUTION

DO NOT SPRAY ON OR NEAR OPEN FLAME. DO NOT USE WHILE SMOKING. Store in a cool place. Keep container tightly closed. Avoid contamination of milk and milk hauling equipment. Avoid overspraying, which will result in wetting animals' hides.

DIRECTIONS FOR CONTAINER DISPOSAL

Pints, quarts, 1-and 5-gallon cans. Do not re-use empty container. Destroy it by perforating or crushing. Bury or discard in a safe place.

30-and 55-gallon drums. Do not re-use empty drum. Return to drum conditioner or destroy by perforating or crushing and burying in a safe place.

EPA REG. NO. 6662-20