

PM 23

6665-8

Revised  
FILE 10/16

United States Environmental Protection Agency  
Washington, D.C. 20460

Office of  
Prevention, Pesticides and  
Toxic Substances

OCT 15 1993

John L. White  
GRO GREEN PROD INC  
717 ELK ST  
BUFFALO NY 14210


Subject: Label Amendment Submission of 06/21/93  
in Compliance with the PR Notice 93-7  
EPA Reg Nr. 6665-8  
GRO GREEN WEED 'N FEED 60

Dear Registrant:

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, REMOVES THE PRODUCT FROM THE SCOPE OF THE REQUIREMENTS CONTAINED IN 40 CFR PART 156, SUBPART K. Five copies of the finished labeling must be submitted prior to releasing the product for shipment.

The Agency is accepting only those changes to your labeling submitted in connection with your amendment application dated 06/21/93 which are necessary to comply with the requirements of 40 CFR part 156, subpart K. Any proposed changes to the labeling submitted in connection with this amendment application that were not directly related to compliance with the requirements of PR Notice 93-7 were not considered and have not been accepted by the Agency. If you wish to make such changes, you must submit a separate amendment application proposing such changes. If your product is currently suspended, the acceptance of this labeling amendment does not affect in any way the suspension.

Sincerely,

  
Jim Tompkins, Deputy Chief  
Registration Support Branch

BEST AVAILABLE COPY

2076

(FRONT PANEL)

8/93

ACCEPTED  
with COMMENTS  
in EPA Letter Dated

OCT 15 1993

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

66658

**GRO GREEN**

**WEED'n FEED 60**

- \*APPLY ANYTIME DURING GROWING SEASON SPRING - SUMMER - FALL
- \* CONTROLS OVER 60 TYPES OF UGLY WEEDS
- \* KILLS DANDELIONS AND OTHER BROADLEAF WEEDS COMPLETELY - ROOTS AND ALL
- \* FEEDS YOUR LAWN WITH SUPERTURF TO PRODUCE A DEEP GREEN, VIGOROUS COMPETITIVE TURF.

**26-3-4**

**ACTIVE INGREDIENTS**

2,4-DICHLOROPHOXYACETIC ACID*	0.87%
ISOCTYL ESTER OF 2-(2-MENTHYL-4-CHLORO-PHOXY)	
PROPLONIC ACID**	0.90%
INERT INGREDIENTS.....	98.23%
TOTAL.....	100.00%
CHLORINE (Cl) MAXIMUM.....	2.5%

**GUARANTEED ANALYSIS**

TOTAL NITROGEN (N).....	26.0%
1.0% AMMONIACAL NITROGEN	
25.0% UREA NITROGEN	
AVAILABLE PHOSPHORIC ACID (P2O5).....	3.0%
SOLUBLE POTASH (K2O).....	4.0%

**PRIMARY PLANT FOOD DERIVED FROM UREA, DIAMMONIUM PHOSPHATE AND MURIATE OF POTASH**

**FEEDS 5,000 SQ. FT.  
(50 X 100)**

**KEEP OUT OF REACH OF CHILDREN**

**NET WT. 17 LBS.  
(7.8 kg)**

**CAUTION**

**SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS  
AND STATEMENT OF PRACTICAL TREATMENT**

**EPA REG. NO. 6665-8**

**EPA EST. NO. 6665-NY-01**

**WEED'N FEED 60**

**BEST AVAILABLE COPY**

3046

(REAR PANEL)

GRO GREEN

FEEDS 5,000 SQ. FT.  
(50 X 100)

NET WT. 17 LBS.  
(7.8 kg)

**WEED'N FEED 60**

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION**

AVOID CONTACT WITH THE SKIN, EYES OR CLOTHING. AVOID BREATHING DUST. WASH THOROUGHLY AFTER HANDLING. CAUSES MODERATE EYE INJURY.

WHEN USING THIS PRODUCT, WEAR LONG SLEEVED SHIRT, LONG PANTS, SOCKS AND SHOES, AFTER USING THIS PRODUCT, REMOVE CLOTHING AND LAUNDRY SEPARATELY BEFORE REUSE AND PROMPTLY AND THOROUGHLY WASH HANDS AND EXPOSED SKIN WITH SOAP AND WATER. THE MAXIMUM NUMBER OF BROADCAST APPLICATIONS PER TREATMENT SITE IS 2 PER YEAR. DO NOT ALLOW PEOPLE (OTHER THAN APPLICATOR) OR PETS ON TREATMENT AREA. DURING APPLICATION, DO NOT ENTER TREATMENT AREA UNTIL DUST HAS SETTLED.

**STATEMENT OF PRACTICAL TREATMENT**

- IF SWALLOWED: GET MEDICAL ATTENTION.
- IF ON SKIN: WASH SKIN WITH SOAP AND WATER.
- IF IN EYES:: IMMEDIATELY FLUSH EYES WITH WATER FOR 15 MINUTES.. IF IRRITATION DEVELOPS, GET MEDICAL ATTENTION.

**ENVIRONMENTAL HAZARDS**

THIS PRODUCT IS TOXIC TO AQUATIC INVERTEBRATES. DRIFT OR OFFGAS MAY ADVERSELY AFFECT AQUATIC INVERTEBRATES AND NON TARGET PLANTS. DO NOT APPLY DIRECTLY TO WATER OR TO AREAS WHERE SURFACE WATER IS PRESENT OR TO INTERTIDAL AREAS BELOW THE MEAN HIGH WATER MARK. DO NOT CONTAMINATE WATER WHEN DISPOSING OF EQUIPMENT WASHWATERS.

NOT A COPY

4076

(REAR PANEL)

**DIRECTIONS FOR USE**

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

GRO-GREEN WEED'N FEED 60 IS A STABLE MIXTURE OF TWO HERBICIDES. IT IS FOR USE ON HOME LAWNS, TURF (EXCLUDING PASTURES, SOO FARMS, AND TURF GROWN TO PRODUCE SEED) AND GOLF FAIRWAYS, APRONS, TEES AND ROUGHS, PARKS, CEMETERIES AND SIMILAR NON-CROP AREAS.

THIS PRODUCT MAY BE USED ON BLUEGRASS, FESCUE, RYE, BERMDUDA, BAHIA, ZOYSIA, CENTIPEDE AND ST. AUGUSTINE GRASS. DO NOT USE ON DICHONDRA AND BENTGRASS.. DO NOT USE THIS PRODUCT WHERE DESIRABLE CLOVERS ARE PRESENT OR ALLOW THE PRODUCT TO "WIND DRIFT" ONTO FLOWERS, VEGETABLES, ORNAMENTALS, SHRUBS, TREES OR OTHER DESIRABLE PLANTS

THIS PRODUCT IS DESIGNED TO PROMOTE LUSH GREEN GROWTH AND CONTROL MANY BROADLEAF WEEDS IN LAWNS. SOME OF THESE WEEDS ARE:

BLACK MEDIC, BUCKHORN, BURDOCK, CHICORY, CHICKWEED, CLOVER, DOCK, DANDELION, ENGLISH DAISY, GROUND IVY, HAEALALL, LAMB'S QUARTER, MALLOW, MORNING GLORY, PIGWEED, PLANTAIN, POISON IVY, POISON OAK, PURSLANE, RAGWEED, RED CLOVER, RED SORREL, STITCHWORT, SPEEDWELL, SPRUGE, WHITE CLOVER, WILD CARROT, WILD GARLIC, WILD ONION, WILD LETTUCE, YARROW AND OTHER BROADLEAF WEEDS.

**WHEN TO APPLY:** BEST RESULTS ARE OBTAINED WHEN WEEDS ARE ACTIVELY GROWING SPRING, OR EARLY FALL.

**IMPORTANT:** APPLY WHEN GRASS IS MOIST (AFTER A RAIN, SPRINKLING, OR DEW) **DO NOT** water lawn after material is applied. For most effective broadleaf weed killing action, avoid unnecessary disturbances, including watering of treated areas for 48 hours after application. Watering will wash off weed killing material. Avoid contact with valuable shrubs, plants or flowers.. Do not over-apply. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings.

**NOTE:** When applying to Southern Grasses - under conditions of inadequate moisture, high temperature and high humidity, overapplication may cause minor transitory discoloration on Centipede and St. Augustine grasses. It is recommended that during hot, dry conditions, application rate should be reduced 50% to avoid possible dehydration or browning of the grass.

**BEST AVAILABLE COPY**

(Rear panel contd)

**HOW TO APPLY:** This 17 pound bag should be used on 5,000 square feet. Apply evenly with a lawn spreader as a one pass application. A two pass application, crisscrossing the area at half the rate each time is also acceptable. Lawns that are heavily overrun with weeds and that are in poor condition may require more than one application of the product. A suggested second application should be made in approximately 30 days.

The following settings are approximate for applying contents of this package at a suggested rate of 3.4 pounds per 1,000 square feet. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 250 sq. ft. and adjust to apply at the rate of .85 lb. per 250 sq. ft. Clean spreader thoroughly after use.

**SPREADER SETTINGS-**

Quick & Easy - We recommend the Cyclone (roto, broadcast) spreaders.

1. They cover 5,000 sq. ft. (50' x 100') in about 15 min.
2. Overlap 6-8" and you will get professional results without streaking.

On standard models set all makes about 1/5 of total opening.

Cyclone (Roto) Type.....	4
Scotts-Standard Models Only.....	5
Central, Greenfield Cross Country.....	6

These settings are furnished as a guide ONLY - Age, condition of spreader, speed of operation and roughness of terrain can cause variations in rates applied.

Sweep all fertilizer materials from concrete surfaces immediately after application. Do not wash material from concrete surfaces because fertilizers which contain iron may cause discoloration upon oxidation.

**STORAGE:** Always Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals. Do not store near foodstuffs or feed. To prevent cross contamination, do not store near other fertilizers, or near herbicides, insecticides, or fungicides.

**PRODUCT DISPOSAL:** To dispose of bags which contain unused product, securely wrap bag in several layers of newspaper and discard in trash.

**CONTAINER DISPOSAL:** Do not reuse empty bag. Fold empty bag and discard in trash.

**BEST AVAILABLE COPY**

6076

(Rear panel contd)

The Manufacturer and Seller makes no warranties, express or implied, concerning this product or its use, which extend beyond the description on the face hereof. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

Manufactured by: **GRO-GREEN PRODUCTS, 717 ELK ST., BUFFALO NEW YORK 14210**

(ERP, WPS, WET) (080293 JLW)

**BEST AVAILABLE COPY**