

DEC 15 1993

Mr. John L. White
 Gro Green
 717 Elk Street
 Buffalo, New York 14210

Dear Mr. White:

SUBJECT: Gro Green
 EPA Registration No. 6665-8
 Commercial and Homeowner Label Amendments
 Your Resubmission Dated November 17, 1993

The labelings referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable with the following provisions:

1. On the Homeowner labeling make the following changes:
 - a. Under the Environmental Hazards section, change the third sentence to read as follows: "Do not apply directly to water."
 - b. Under the Directions for Use section, add the following heading at the end of the first paragraph: "For Homeowner Use Only."
2. On the Commercial labeling make the following change:
 - a.. To remove this product from the scope of the Worker Protection Standards, add the following statement: "Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes."

A stamped copy is enclosed for your records. Please submit five (5) final printed copies for the referenced label, incorporating the above changes, before releasing the product for shipment.

Sincerely yours,

BEST AVAILABLE COPY

Joanne I. Miller
 Product Manager (23)
 Fungicide-Herbicide Branch
 Registration Division (H7505C)

CONCURRENCES							
SYMBOL	Enclosure						
SURN/ME	JSM						
DATE	12/15/93						

(FRONT PANEL)

PG 1

GRO GREEN

WEED'n FEED 60

- *APPLY ANYTIME DURING GROWING SEASON SPRING - SUMMER - FALL
- * CONTROLS OVER 60 TYPES OF UGLY WEEDS
- * KILLS DANDELIONS AND OTHER BROADLEAF WEEDS COMPLETELY - ROOTS AND ALL
- * FEEDS YOUR LAWN WITH **SUPERTURF** TO PRODUCE A DEEP GREEN, VIGOROUS COMPETITIVE TURF.

28-3-4

ACTIVE INGREDIENTS

2,4-DICHLOROPHENOXYACETIC ACID*	0.87%
ISOOCTYL ESTER OF 2-(2-MENTHYL-4-CHLORO-PHENOXY)	
PROPLONIC ACID**	0.90%

INERT INGREDIENTS

TOTAL	100.00%
-------	---------

CHLORINE (Cl) MAXIMUM	0.5%
-----------------------	------

GUARANTEED ANALYSIS

TOTAL NITROGEN (N)	28.0%
1.0% AMMONIACAL NITROGEN	
27.0% UREA NITROGEN	
AVAILABLE PHOSPHORIC ACID (P2O5)	3.0%
SOLUBLE POTASH (K2O)	4.0%

PRIMARY PLANT FOOD DERIVED FROM UREA, DIAMMONIUM PHOSPHATE AND MUURIATE OF POTASH

FEEDS 5,000 SQ. FT.
(50 X 100)

KEEP OUT OF REACH OF CHILDREN

NET WT. 15 LBS

(67.25 kg)

CAUTION

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

AND STATEMENT OF PRACTICAL TREATMENT

EPA REG. NO. 6665-8

WEED'N FEED 60

ACCEPTED
with COMMENTS
In EPA Register Dated
EPA REG. NO. 6665-8W-01
15 DEC 1995
Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
6665-8

BEST AVAILABLE COPY

(REAR PANEL)

Pg 2

GRO GREEN

FEEDS 5,000 SQ FT
(50 X 100)

NET WT 16 LBS
(7.25 kg)

WEED'N FEED 60

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

AVOID CONTACT WITH THE SKIN, EYES OR CLOTHING. AVOID BREATHING DUST. WASH THOROUGHLY AFTER HANDLING. CAUSES MODERATE EYE IRRITATION.
WHEN USING THIS PRODUCT, WEAR LONG SLEEVED SHIRT, LONG PANTS, SOCKS AND SHOES. AFTER USING THIS PRODUCT, REMOVE CLOTHING AND LAUNDRY SEPARATELY BEFORE REUSE AND PROMPTLY AND THOROUGHLY WASH HANDS AND EXPOSED SKIN WITH SOAP AND WATER. THE MAXIMUM NUMBER OF BROADCAST APPLICATIONS PER TREATMENT SITE IS 2 PER YEAR. DO NOT ALLOW PEOPLE (OTHER THAN APPLICATOR) OR PETS ON TREATMENT AREA DURING APPLICATION. DO NOT ENTER TREATMENT AREA UNTIL DUST HAS SETTLED.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED:	GET MEDICAL ATTENTION
IF ON SKIN:	WASH SKIN WITH SOAP AND WATER
IF IN EYES:	IMMEDIATELY FLUSH EYES WITH WATER FOR 15 MINUTES IF IRRITATION DEVELOPS, GET MEDICAL ATTENTION

ENVIRONMENTAL HAZARDS

THIS PRODUCT IS TOXIC TO AQUATIC INVERTEBRATES. DRIFT OR RUNOFF MAY ADVERSELY AFFECT AQUATIC INVERTEBRATES AND NON-TARGET PLANTS. **DO NOT** APPLY DIRECTLY TO WATER OR TO AREAS WHERE SURFACE WATER IS PRESENT OR TO INTERTIDAL AREAS BELOW THE MEAN HIGH WATER MARK. **DO NOT** CONTAMINATE WATER WHEN DISPOSING OF EQUIPMENT WASHWATERS.

BEST AVAILABLE COPY

GROUNDWATER CONTAMINATION

MOST CASES OF GROUNDWATER CONTAMINATION INVOLVING HERBICIDES SUCH AS 2,4-D HAVE BEEN ASSOCIATED WITH MIXING/LOADING AND DISPOSAL SITES. CAUTION SHOULD BE EXERCISED WHEN HANDLING 2,4-D PESTICIDES AT SUCH SITES TO PREVENT CONTAMINATION OF GROUNDWATER SUPPLIES. USE OF CLOSED SYSTEMS FOR MIXING OR TRANSFERRING THIS PESTICIDE WILL REDUCE THE PROBABILITY OF SPILLS. PLACEMENT OF THE MIXING/LOADING EQUIPMENT ON AN IMPERVIOUS PAD TO CONTAIN SPILLS WILL HELP PREVENT GROUNDWATER CONTAMINATION.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

GPO-GREEN WEED'N FEED 60 IS A STABLE MIXTURE OF TWO HERBICIDES. IT IS FOR USE ON HOME LAWNS, TURF (INCLUDING PASTURES, SOO FARMS) AND GOLF FAIRWAYS, APPRONS, TEES AND ROUGHS.

THIS PRODUCT MAY BE USED ON BLUEGRASS, FESCUE, RYE, BERBERIDA, BAHIA, COYSLA, CENTIPEDE AND ST. AUGUSTINE GRASS. **DO NOT USE** ON DICHOLOPA AND BENTGRASS. **DO NOT USE** THIS PRODUCT WHERE DESIRABLE CLOVERS ARE PRESENT OR ALLOW THE PRODUCT TO "WIND DRIFT" ONTO FLOWERS, VEGETABLES, ORNAMENTALS, SHRUBS, TREES OR OTHER DESIRABLE PLANTS.

THIS PRODUCT IS DESIGNED TO PROMOTE LUSH GREEN GROWTH AND CONTROL MANY BROADLEAF WEEDS IN LAWNS. SOME OF THESE WEEDS ARE:

BLACK MEDIC, BUCKHORN, BURDOCK, CHIRUPY, CHICK WEED, CLOVER, DOCK, DANDELION, ENGLISH DAISSY, GROUND IVY, HAEALALL, LAMB'S QUARTER, MALLOW, MORNING GLORY, PIGWEED, PLANTAIN, POISON IVY, POISON OAK, PURSLANE, PIGWEED, RED CLOVER, RED SOPPEL, STITCHWORT, SPEEDWELL, SPRUCE, WHITE CLOVER, WILD CAPPOT, WILD GARLIC, WILD ONION, WILD LETTUCE, YARROW AND OTHER BROADLEAF WEEDS.

WHEN TO APPLY: BEST RESULTS ARE OBTAINED WHEN WEEDS ARE ACTIVELY GROWING. SPRING, OR EARLY FALL.

IMPORTANT: APPLY WHEN GRASS IS MOIST (AFTER A RAIN, SPRINKLING, OR DEW). **DO NOT** water lawn after material is applied. For most effective broadleaf weed killing action, avoid unnecessary disturbances, including watering of treated areas for 48 hours after application. Watering will wash off weed killing material. Avoid contact with valuable shrubs, plants, etc.

BEST AVAILABLE COPY

flower: Do not over-apply. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings.

NOTE: When applying to Southern Grasses - under conditions of inadequate moisture, high temperature and high humidity, overapplication may cause minor transitory discoloration on Centipede and St. Augustine grasses. It is recommended that during hot, dry conditions, application rate should be reduced 50% to avoid possible dehydration or browning of the grass.

HOW TO APPLY: This 16 pound bag should be used on 5,000 square feet. Apply evenly with a lawn spreader as a one pass application. A two pass application, crisscrossing the area at half the rate each time is also acceptable. Lawns that are heavily overrun with weeds and that are in poor condition may require more than one application of the product. A suggested second application should be made in approximately 30 days.

The following settings are approximate for applying contents of this package at a suggested rate of 3.2 pounds per 1,000 square feet. Settings are based on an average walking speed. Reduce setting for slower speed, raise setting for higher speed. Check your spreader on 250 sq. ft. and adjust to apply at the rate of .08 lb. per 250 sq. ft. Clean spreader thoroughly after use.

SPREADER SETTINGS-

Quick & Easy - We recommend the Cyclone (Roto, broadcast) spreaders.

- 1. They cover 5,000 sq. ft. (50' x 100') in about 15 min!
- 2. Overlap 6-8" and you will get professional results without streaking.

On standard models set all makes about 1/5 of total opening.

Cyclone (Roto) Type	4
Scotts-Standard Models Only	5
Central, Greenfield Cross Country	6

These settings are furnished as a guide ONLY - Age, condition of spreader, speed of operation and roughness of terrain can cause variations in rates applied.

Sweep all fertilizer materials from concrete surfaces immediately after application. Do not wash material from concrete surfaces because fertilizers which contain iron may cause discoloration upon oxidation.

STORAGE: Always Store in original container and keep any bags containing unused product in a

locked storage area inaccessible to children and domestic animals. Do not store near foodstuffs or feed. To prevent cross contamination, do not store near other fertilizers, or near herbicides, insecticides, or fungicides.


PRODUCT DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinseate, is a violation of Federal Law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA regional office for guidance.

CONTAINER DISPOSAL: Completely empty bag into application equipment, then dispose of empty bag to a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

The Manufacturer and Seller makes no warranties, express or implied, concerning this product or its use, which extend beyond the description on the face hereof. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

Manufactured by **GRO-GREEN PRODUCTS, 717 ELK ST., BUFFALO NEW YORK 14210**

(EPP, WPS, WET, 16# CUR III) (11/93 JLW)



BEST AVAILABLE COPY