

NEW FORMULA FIESTAR

We have improved the effectiveness of FIESTAR for Crabg. ss control and therefore the following instructions replace those printed on this bag:

CAUTION:

"Keep out of reach of children. Keep children and pets off the treated areas until this material has been thoroughly washed into the soil and grass completely dried." See other cautions below:

LAWN CARE

To prevent crabgrass--fertilize the soil--and as a soil conditioner-- It improves the soil to assure a better lawn-growth by increasing the humus content of the soil.

ESTABLISHED LAWNS

Broadcast FIESTAR at rate of 1 lb. per 15 sq. ft. before crabgrass seeds have germinated--In late Winter to early Spring. (50 pounds will cover 750 sq. ft.)

NEW LAWNS--Broadcast FIESTAR at rate of 1 lb. per 15 sq. ft. after third mowing of newly established lawns.

FOR CRABGRASS CONTROL

For pre-emergence CRABGRASS control, apply in EARLY SPRING or LATE WINTER. Do not reseed until one month after treating with FIESTAR.

FERTILIZER CONTENT
Guaranteed Analysis

Nitrogen 2%
Phosphoric Acid 2%
Potash 0

INSECTICIDE CONTENT

ACTIVE INGREDIENT
*Technical chlordane 2.3%
INERT INGREDIENTS 97.7% (including wetting agents)
100.0%

*Equivalent to 1.3% octachloro-4,7-
Methanotetrahydroindane and .92% of
related compounds

APPLICATION:

Can be reseed one month after application.
Do not reseed with CRABGRASS when reseeding.

CAUTION:

AVOID inhalation, avoid skin contact with solution. In case of contact with skin, wash with soap and water. Wash hands after use. Do not contaminate drinking water or use near fish ponds.

Do not reuse container when empty and dispose of bag promptly. Do not let the bag lay around, but place same in a closed metal container (garbage can).

Contamination of waters by application, cleaning of equipment, or disposal of waste will kill fish.

USDA, REG. NO. 6699-2

