



# FIELD MOUSE BAIT



## POISON

E.P.A. Registration No. 6704-20  
E.P.A. EST. 6704-IN-1

**WARNING:** Keep out of reach of children. See additional warnings below:

<b>Active Ingredient:</b>	
Zinc Phosphide .....	2.0%
<b>Inert Ingredients:</b> .....	98.0%
<b>Total</b>	<b>100.0%</b>

**DIRECTIONS FOR USE:** For the control of mice in orchards and other crops during fall-winter months, except deer or white-footed mice (*Peromyscus*).


Place teaspoonful quantities in active runways under cover of grass. Two or four bait spots should be made near the base of each infested tree. Baiting should be conducted at the rate of 2-3 lbs. per acre. In broadcast operations, apply 6-10 lbs. per acre in heavy vegetation. Do not apply to bare ground, nor broadcast over growing crops. Wear rubber gloves when handling bait. To reduce the hazard to beneficial wildlife, scatter bait evenly or hide under cover in runways.

**ANTIDOTE:** Give a teaspoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Have victim lie down and keep warm. Call a physician immediately.

**WARNING:** May Be Fatal If Swallowed! Do not breathe dust or fumes. Avoid contact with skin. Wash hands after using. Do not contaminate feed and foodstuffs. Keep away from children and domestic animals.

Do not reuse bag. Dispose of bag by deep burial or burning. Stay out of smoke.

U. S. Department of the Interior  
Fish and Wildlife Service  
Washington, D. C. 20240



Net Weight (25 lbs. or 50 lbs.)

PLANT DOCUMENT AVAILABLE