

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS (4 DOMESTIC ANIMALS)

DANGER

For use by trained operators only. May be fatal if swallowed or absorbed through skin. Avoid contact with skin and eyes. May produce severe burns: care must be exercised in handling the concentrated form of LAMPRECID®. Protective clothing, rubber gloves, and face masks are necessary for the minimum protection of handlers. Wash thoroughly after handling.

ENVIRONMENTAL HAZARDS

Directions for use must be strictly followed to minimize hazard to non-target organisms. Do not contaminate water by cleaning of equipment or disposal of wastes.

Local, State, and Provincial Fish and Game Agencies must be contacted before product is applied. Municipalities that use streams requiring treatment as potable water sources must be notified of the impending treatment at least 24 hours prior to application. Agricultural irrigators that use streams requiring treatment as a source of irrigation water must be notified of the impending treatment at least 24 hours prior to application. Agricultural irrigators must turn off their irrigation system for a 24-hour period during and after treatment.

Not to be used by unauthorized personnel. Nr. 2039

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. SEE RIGHT PANEL FOR DIRECTIONS.

CATEGORY OF APPLICATOR

Aquatic pest control.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

STORAGE: Containers of LAMPRECID® should be stored in an area where spill or leakage of the material can be contained.

DISPOSAL: LAMPRECID® spray mixture or residue that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.

Note Area (or equivalent) and offer container for recycling, reconditioning, or disposal in a safe place.

Consult Federal, provincial, state or local authorities for approved alternative procedures.

RESTRICTED USE PESTICIDE

ONLY FOR SALE TO AND APPLICATION BY CERTIFIED APPLICATORS OF THE U.S. FISH AND WILDLIFE SERVICE, FISHERIES AND OCEANS CANADA, AND PROVINCIAL AND STATE FISH AND GAME EMPLOYEES OR PERSONS UNDER THEIR DIRECT SUPERVISION.

SEA LAMPREY LARVICIDE LAMPRECID®

ACTIVE INGREDIENT:

α, α, α -TRIFLUORO-4-NITRO-m-CRESOL, SODIUM SALT:

EQUIVALENT TO

FREE CRESOL

INERT INGREDIENT:

38.8%

61.2%

TOTAL: 100.00%

THIS PRODUCT CONTAINS

LBS. OF TFN
GMS.

PER GALLON
LITRE

KEEP OUT OF REACH OF CHILDREN DANGER-POISON



STATEMENT OF PRACTICAL TREATMENT

NO KNOWN EFFECTIVE ANTIDOTE:

IF SWALLOWED, CALL PHYSICIAN IMMEDIATELY. INDUCE VOMITING by giving milk or white of egg, beaten in water, then a tablespoon of salt in glass of warm water and repeat until vomit fluid is clear. Repeat milk or white of egg beaten with water.

IF ON SKIN, flush liberally with water for at least 15 minutes and wash with soap if possible.

IF IN EYES, flush liberally with water for at least 15 minutes; get medical attention.

SEE LEFT PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

Filed for: U.S. Fish and Wildlife Service
Washington, DC 20540
EPA Registration No. 0704-48
Fisheries and Oceans Canada, Ottawa, Ontario
Canadian Registration No. 11763 Pest Control
Products Act

MFG. by: Hoechst Aktiengesellschaft
Tonn, Country: 6730 Frankfurt am Main 90, Germany
Establishment No. 0300-970-4

Hoechst

Net Contents _____ Lbs.
Batch No. _____ Kgs.

DIRECTIONS FOR USE

Pretreatment surveys are always made to determine of sea lamprey larvae (*Petromyzon marinus*) in the Great Lakes Basin and the Lake Champlain selected for treatment must first be analysed or mine both the minimum concentration of required to kill sea lamprey larvae and the maximum that can be applied without causing undue non-target organisms. "Analysis" constitutes live say, or the use of a multiple regression relating to on-site determination of total alkalinity and the body of water.

The concentration of LAMPRECID® needed to kill larvae may vary depending upon water chemistry. Measure volume or flow-rate and add 1 LAMPRECID® necessary at rates based on the analysis. Dispense LAMPRECID® by application, clearly accurate to maintain predetermined concentration in the body of water must be made by colorimetric analysis, gas chromatography, or manometric chromatography.

LAMPRECID® may be used by itself in the treatment in the Great Lakes Basin and the Lake Champlain times, however, BAYLUSCIDE 70% Wettable REG NO. 3125-136 may be used in combination CID® (EPAREG NO. 6704-45) for control of sea lamprey. Application of BAYLUSCIDE 70% Wettable Powder simultaneous addition with LAMPRECID® on land reduce the amount of LAMPRECID® required or addition downstream to enhance LAMPRECID® activity. Prior to using BAYLUSCIDE-LAMPRECID® surveys must be made to determine larval density. When using BAYLUSCIDE 70% Wettable Powder with LAMPRECID®, mix in proportions so that final concentration of BAYLUSCIDE 70% Wettable not more than 2% of LAMPRECID® by weight 70% Wettable Powder may be added to LAMPRECID® ways:

1. One method of application is to add a slurry of 70% Wettable Powder pumped into the air pump while the LAMPRECID® is fed separately conventional fuel pump feeder in amount deliver the desired ratio of BAYLUSCIDE to LAMPRECID® is more easily mixed as a slurry mixing with LAMPRECID® and more uniform result.
2. A second application method is used on the farms with multiple tributaries. The number sites on these large rivers precludes the use of a pump because of the number of feeder pumps need for a 110-volt power source at each site and the often difficult access to sites. On farms, LAMPRECID® alone is fed into the tributaries join to form the main stem of the river. BAYLUSCIDE is introduced into the chemical bank calculated to produce the desired LAMPRECID® ratio. The LAMPRECID® applications are timed so that the individual chemical banks in the main stream that approximate concentrations in the tributaries. Since diluted by ground water, swamp seepage, or rain, occasional rain, or other conditions included when the application rates for the tributaries, the toxicity of the bank in the main stream raised by the addition of LAMPRECID® or BAYLUSCIDE can be used in place of LAMPRECID® increased toxicity of the BAYLUSCIDE-LAMPRECID® alone to sea lamprey.

BEST AVAILABLE COPY

Under
Hoechst Akt. & Chem. Ges.
Pesticide Reg. No. 0704-48
EPA Reg. No. 0704-48
JUL 06 1990
6704-45