

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
(& DOMESTIC ANIMALS)
DANGER**

Harmful if swallowed, inhaled, or absorbed through the skin. Avoid contact with eyes, skin, or clothing. Handle only with protective gloves, clothing, and face mask, or respirator. Wash hands with soap and water after handling.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to birds. Do not expose in areas accessible to waterfowl, poultry, and other non-target birds. Keep out of lakes, ponds, streams, tidal marshes, and estuaries. Do not apply where runoff is likely to occur. Do not contaminate water by the cleaning of equipment or disposal of waste.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

STORAGE—Keep dry. Store in an isolated, well ventilated room. Do not contaminate water, food, or feed by storage or disposal.

DISPOSAL—Open burning and dumping of this product is prohibited. Do not re-use empty container. Consult local, state, and federal disposal authorities for approved procedure for disposing of excess or unused concentrate and container.

GENERAL: Use 1339 Gull Toxicant 98% Concentrate for preparing bread baits to control herring gulls (*Larus argentatus*) and great black-backed gulls (*Larus marinus*) only. Apply only within the coastal area of Northeastern United States (Delaware, New York,

SEE RIGHT PANEL FOR ADDITIONAL
GENERAL, PREPARATION,
AND APPLICATION

**RESTRICTED USE
PESTICIDE**

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's Certification.

For use only by U.S. Fish and Wildlife Personnel trained in Bird Control or persons under their Direct Supervision.

**1339 GULL TOXICANT 98%
CONCENTRATE**

Active Ingredient: 3-Chloro-4-Methyl Benzenamine Hydrochloride	98%
Inert Ingredients:	2%
Total	100%

Net Weight 16 ozs. (1 lb.)

KEEP OUT OF REACH OF CHILDREN

ACCEPTED

MAR 30 1979

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 6704-77

DANGER—POISON



POISON

Statement of Practical Treatment

IF SWALLOWED: Induce vomiting and immediately call a physician.

IF INHALED: Move patient from contaminated area and immediately call a physician.

IF ON SKIN OR IN EYES: Immediately flush eyes or skin with water. Call a physician if irritation develops and persists.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

MANUFACTURED FOR:
U.S. DEPARTMENT OF THE INTERIOR
FISH AND WILDLIFE SERVICE
WASHINGTON, D.C. 20240

ESTABLISHMENT NO. 6704-IN-1
EPA REGISTRATION NO. 6704-77

New Jersey, Connecticut, Rhode Island, Massachusetts, New Hampshire, and Maine) in breeding areas or colonies within predation radius of important nesting colonies of terns, puffins, laughing gulls, and other colonial nesting seabirds from March 1 to June 30 each year.

BAIT PREPARATION: Blend 0.2 oz. (6 gms.) 1339 Gull Toxicant 98% Concentrate into 1 lb. (454 gms.) melted, stick oleomargarine. Spread ½ oz. (15 gms.) of blended mixture (1 tablespoon) on a slice of standard sandwich bread and cover with another slice. Immediately cut each sandwich into 9 equally-sized cubes. Prepared baits must be placed in a plastic bag for transportation or distribution and must be used within 12 hours.

BAIT APPLICATION

Pre-Treatment—Each site destined to be treated will be prebaited with untreated bread cubes to insure rapid bait acceptance.

Treatment—Treatments will be made by hand only in or near nesting colonies of the target species. Treated bread cubes will be broadcast or placed only in the same areas where bread cubes were accepted during prebaiting. Initial applications will be broadcast; however, no broadcast application will be made after April 20. Application after April 20 will be made at or in gull nests. The number of bait applications will be determined by the degree of control provided by previous applications; however, no more than 10 bait applications should be made in or near individual colonies. The number of baits exposed at an individual site must not exceed 5 times the total number of gulls to be controlled at that location.

Post-Treatment—Baits regurgitated or not accepted must be retrieved within 12 hours after each application and disposed of by burial or other adequate means. A search must be conducted within 48-72 hours after application to remove and dispose of bird carcasses, except for those areas where disturbances of eiders may adversely affect their breeding efforts.

**DO NOT USE TREATED BREAD AS
FOOD OR FEED**

6704-77

3-30-1979

1/1