

6720-9

### CAUTION

Harmful if swallowed. Avoid prolonged breathing of dust or spray mist. Avoid prolonged or repeated contact with skin; wash thoroughly after using. Avoid contamination of feed and foodstuffs. Keep out of reach of children.

### NOTICE

Seller's guaranty shall be limited to the terms of the label and subject thereto. The buyer assumes any risk of damage to persons or property arising out of use or handling, and accepts the product on these conditions.

#### ANIMALS

**BEEF CATTLE: LICE** — 1 gallon to 100 gallons of water or 6 1/2 oz. per 5 gallons of water.

**TICKS** — 1 - 2 gallons per 100 gallons of water or 6 1/2 - 13 oz. per 5 gallons of water. Repeat application only as needed at 2 week intervals.

**HORN FLY** — 1 - 1 1/2 gallons to 100 gallons of water or 6 1/2 - 13 oz. per 5 gallons.

Apply complete coverage spray. One treatment may be sufficient. Repeat application only as need at 2 week intervals.

DO NOT use on animals less than 1 month old. Avoid contamination of feed or food.

**DAIRY CATTLE (Non-Lactatin)** use 1 - 1 1/2 gallons per 100 gallons of water for dip and spray. DO NOT treat during lactation or spray within 2 weeks of freshening. DO NOT dip dairy animals within 2 months of freshening. DO NOT apply to calves under one month of age. DO NOT apply to lactating dairy cattle.

**SWINE: LICE** — 1 gallon to 100 gallons of water or 6 1/2 oz. per 5 gallons of water. Apply as a complete spray to animals, pens and litter. One treatment may be sufficient. Repeat application only as needed.

#### POULTRY

**CHICKENS, DUCKS, GEESE, TURKEYS** — For Northern Fowl Mite, Poultry Lice, Chicken Red Mite use at the rate of 2 tbs. per 1 gallon of water per 100 - 150 Birds as direct application. Repeat application in 4 - 8 weeks or when necessary.

As a Premise Treatment use 4 tbs. per 1 gallon of water. Apply liberally to litter, walls, ceilings, roosts, nests and adjacent areas, taking care to force spray in cracks and crevices. Repeat as necessary. Avoid contamination of feed, feeding troughs, and water fountains. Provide clean nesting material after spraying.

As a Roost Pait use 4 tbs. per 1 gallon of water at a rate of 1 pint per 150 ft. of roost, as an aid in the control of Chicken rec mite and Poultry lice. Brush on top sides of cleared perches or roost poles before birds go to roost. If necessary, repeat in 8-10 days.

#### DOGS & CATS DIRECT APPLICATION Sprays

For control of fleas apply spray of 1 oz. of Malathion 57% Emulsifiable Liquid per 1 gal. of water. Wet the animal thoroughly using a hand sprayer.

If the animal has free run, treatments may have to be made every second or third week. To avoid unnecessary applications, animal should be closely examined for fleas before retreating. If the animal is confined to an area free of fleas, one treatment may be sufficient.

#### ANIMAL QUARTERS Sprays

For control of fleas in kennels, pens, yards, lawns and under houses. Remove manure or debris before spraying.

Use spray of 5 ozs. of 57% Emulsifiable Liquid per 1 gal. of water at rate of 1 gal per 1000 sq. ft. Use garden sprayer.

#### ORNAMENTALS

For Oyster Shell Scale and Lace Bugs, use 1 pint. For Spider Mites, Aphids, White Flies, Mealybugs, Thrips, Japanese Beetle Adults, European Pine Shoot Moth, Four-Lined Plant Bug, Tarnished Plant Bug, Rose Leafhopper, Potato Leafhopper, Scurfy Scale and Eucnymus Scale use 1 1/2 pints. For Bagworm, Tent Caterpillar, Boxwood Leaf Miner, Birch Leaf Miner, Oak Kermes, Florida Red Scale, Fletcher Scale, Juniper, Azalea and Magnolia Scales, use 2 pints. For Black Scale crawler, Soft Brown and Monterey Scales, use 2 1/2 pints. Apply thorough full coverage sprays as needed. Note: DO NOT apply to petunias, certain ferns (including Boston, Maidenhair, Pteris), and some species of Crassula, since plant injury may occur.

# MA

ACTIVE INGRE

Malathion  
diethy

INERT INGRE

# CAU

U.S





mist. Avoid prolonged or repeated  
of feed and foodstuffs. Keep out of

thereto. The buyer assumes any risk  
cepts the product on these conditions.

**DOGS & CATS  
DIRECT APPLICATION  
Sprays**

apply spray of 1 oz. of Malathion 57% Emulsi-  
gal. of water. Wet the animal thoroughly using a

free run, treatments may have to be made every  
ek. To avoid unnecessary applications, animal should  
d for fleas before retreating. If the animal is con-  
free of fleas, one treatment may be sufficient.

**ANIMAL QUARTERS  
Sprays**

in kennels, pens, yards, lawns and under houses.  
debris before spraying.  
ozs. of 57% Emulsifiable Liquid per 1 gal. of water  
per 1000 sq. ft. Use garden sprayer.

**ORNAMENTALS**

Scale and Lace Bugs, use 1 pint. For Spider Mites,  
, Mealybugs, Thrips, Japanese Beetle Adults, European  
Four-Lined Plant Bug, Tarnished Plant Bug, Rose Leaf-  
hopper, Scurfy Scale and Eucynimus Scale, use 1 1/2  
m, Tent Caterpillar, Boxwood Leaf Miner, Birch Leaf  
s, Florida Red Scale, Black Scale, Juniper, Azalea  
les, use 2 pints. For Black Scale crawler, Soft Brown  
les, use 2 1/2 pints. Apply thorough full coverage  
Note: DO NOT apply to petunias, certain ferns  
Maidenhair, Pteris, and some species of Crassula,  
may occur.

# MALATHION EM-5

An Emulsifiable Concentrate containing  
5 lbs. of Actual Malathion per Gallon

**ACTIVE INGREDIENTS:**

Malathion (0,0-dimethyl dithiophosphate of  
diethyl mercaptosuccinate)  
Aromatic petroleum derivative solvent

**INERT INGREDIENTS:**

TOTAL

BY WEIGHT

57.0%  
32.0%  
11.0%  
100.0%

**CAUTION: KEEP OUT OF REACH OF CHILDREN**  
SEE SIDE PANEL FOR OTHER CAUTIONS.  
U.S.D.A. REGISTRATION NUMBER 6720-9

MANUFACTURED BY  
**Southern MILL CREEK PRODUCTS COMPANY, INC.**  
TAMPA FLORIDA

M-5

**HOUSEHOLD INSECTS**

For the control of ants, clover mites, crickets, earwigs, pantry pests\*, roaches, scorpions, silverfish and spiders in and about homes, dairy and food-processing plants, use Malathion 57% Emulsifiable Liquid at the rate of **6½ fluid ounces per gallon** of water. Apply by means of a coarse spray or paint brush to window sills, baseboards, under sinks, drainboards, stoves, cracks and crevices and to other areas frequented by insects.

For control of bedbugs in dwellings use Malathion 57% Emulsifiable Liquid at a rate of **1 to 2 fluid ounces per gallon** of deodorized kerosene. Apply lightly to all mattress surfaces in sufficient quantity to "mist" the fabric and generously to beds and woodwork with special care taken to wet all possible hiding places. Remove fish, birds and pets before treating.

Avoid contamination of food, utensils, milk, milk equipment and water.

Water emulsions may stain finished surfaces such as wallpaper, furniture, baseboards, etc.

Repeat applications as necessary.

\*Pantry pests such as the exposed stages of: saw-toothed grain beetle, flour beetle, rice weevil, cigarette beetle, drug store beetle and Indian meal moth.

This product is toxic to fish.

Keep out of lakes, streams and ponds.

**MOSQUITOES AND SMALL FLYING INSECTS**

**OUTDOORS**

Use 2% - 5% Malathion fog, aerosol or space spray. As a spray dilute 57% Emulsifiable Liquid **1 to 28 parts** of water, fuel oil or diesel oil. When using kerosene type solvent as carrier, dilute **1 part** of 57% Emulsifiable Liquid in **28 parts** of a mixture consisting of **4 parts** kerosene type solvent and **1 part** aromatic hydrocarbon type solvent. For 5% dilute 1-11 using similar solvents. Repeat applications as necessary.

Avoid applying oil-based formulations to valuable ornamental plants as injury may occur. Malathion may be toxic to certain species of fish, particularly in shallow water.

**Mosquito larvae in standing water**

Apply 57% Malathion Emulsifiable Liquid at rate of **13 fluid ounces** (approximately ½ pound actual Malathion) per acre. Mix in sufficient water or oil to obtain even coverage when applied by air or ground equipment.

**INDOORS**

Use a 2% Malathion liquid space and contact spray or a 2% Malathion low pressure spray bomb. A knockdown agent of a type commonly used in space sprays should be added to the spray.

Rooms should be thoroughly misted to envelop insects present and should be kept closed for **5 to 10 minutes**. Sweep up and destroy fallen insects. Repeat applications as necessary. **DO NOT** contaminate food, dishes or utensils. **DO NOT** apply oil-based sprays directly to flowers or other ornamental plants in the home.

BY WEIGHT

57.0%  
32.0%  
11.0%  
100.0%

CHILDREN

20--9

, Inc.