

HOUSEHOLD INSECTS

control of ants, clover mites, crickets, earwigs, pantry pests, flies, scorpions, silverfish and spiders in and about home, dairy food-processing plants, use Malathion 50% Emulsifiable Liquid at a rate of 8 fluid ounces per gallon of water. Apply by means of coarse spray or paint brush to window sills, baseboards, under doors, drainboards, stoves, cracks and crevices and to other areas infested by insects.

control of bedbugs in dwellings use Malathion 50% Emulsifiable Liquid at a rate of 1 to 2 fluid ounces per gallon of deodorized kerosene. Apply lightly to all mattress surfaces in sufficient quantity to "mist" the fabric and generously to beds and woodwork with special care taken to wet all possible hiding places.

to prevent contamination of food, utensils, milk, milk equipment and other applications as necessary.

to destroy pests such as the exposed stages of: saw-toothed grain beetle, flour beetle, rice weevil, cigarette beetle, drug store beetle Indian meal moth.

MOSQUITOES AND SMALL FLYING INSECTS OUTDOORS

2% - 5% Malathion fog, aerosols or mist sprays. As a spray use 50% Emulsifiable Liquid 1 to 24 parts of water, fuel oil or kerosene. When using kerosene type solvent as carrier, dilute 1 part of 50% Emulsifiable Liquid in 28 parts of a mixture consisting of 1 part kerosene type solvent and 1 part aromatic hydrocarbon solvent. For 5% dilute 1 to 10 using similar solvents. Repeat applications as necessary.

to prevent injury to valuable ornamentals as injury may occur.

to prevent injury to valuable ornamentals as injury may occur. Fog spray should be directed around areas having puddles, stagnant water, or accumulated debris, and should not be used close to dwellings, or vehicles where residues could be troublesome.

to prevent injury to valuable ornamentals as injury may occur. Fog spray should be directed around areas having puddles, stagnant water, or accumulated debris, and should not be used close to dwellings, or vehicles where residues could be troublesome.

to prevent injury to valuable ornamentals as injury may occur. Fog spray should be directed around areas having puddles, stagnant water, or accumulated debris, and should not be used close to dwellings, or vehicles where residues could be troublesome.

Mosquito larvae in standing water

50% Malathion Emulsifiable Liquid at rate of 15 fluid ounces

(approximately 1/2 pound actual Malathion) per acre only in intermittently flooded areas, stagnant water, temporary rain pools, sloughs and log ponds. Mix in sufficient water or oil to obtain even coverage when applied by air or ground equipment.

INDOORS

Use a 2% Malathion liquid space and contact spray or a 2 1/2 Malathion low pressure bomb.

Rooms should be thoroughly misted to envelop insects present and should be kept closed for 5 to 10 minutes. Sweep up and destroy fallen insects. Repeat applications as necessary. DO NOT contaminate food, dishes or utensils. DO NOT apply oil-based sprays directly to flowers or other ornamental plants in the home.

If used as a crack and crevice spray, do not use in commercial food preparation areas or in edible products areas of food processing plants.

If used as a space spray in food processing plants, foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment. Thoroughly cleaned before using. When using the product as a space spray in these areas, apply only when plant is not in operation.

CAUTION

Harmful if swallowed. Avoid prolonged breathing or spray mist. Avoid contact with skin; wash thoroughly after handling. Keep out of reach of children. Avoid contamination of feed and food stuffs.

This product is toxic to fish. Keep out of lakes, streams or ponds. DO NOT apply where runoff is likely to occur. DO NOT contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

DO NOT reuse empty drum. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place.

NOTICE: Seller's guaranty shall be limited to the terms of the label and subject thereto the buyer assumes any risk to persons or property arising out of use or handling, and accepts the products on these conditions.

6720-17

2/21/71

1 of 3

ANIMALS

BEEF CATTLE: LICE — 1 1/4 gallons to 100 gallons of water or 8 oz. per 5 gallons of water.

TICKS — 1 - 2 gallons per 100 gallons of water or 8 - 16 oz. per 5 gallons of water.

HORN FLY — 1 - 1 1/2 gallons to 100 gallons of water or 8 - 16 oz. per 5 gallons.

Apply complete coverage spray. One treatment may be sufficient. Repeat application only as needed at 2 week intervals.

DAIRY CATTLE (Non-Lactating) use 1 - 1 1/2 gallons per 100 gallons of water for dip or spray. DO NOT spray during lactation or within two weeks of freshening. DO NOT apply to calves under one month of age. DO NOT apply to lactating dairy cattle.

SWINE: LICE — 1 1/4 gallon to 100 gallons of water or 8 oz. per 5 gallons of water. Apply as a complete spray to animals, pens and litter. One treatment may be sufficient. Repeat application only as needed.

POULTRY

CHICKENS, DUCKS, GEESE, TURKEYS — For Northern Fowl Mite, Poultry Lice, Chicken Red Mite use at the rate of 2 1/2 tbs. per 1 gallon of water per 100 - 150 birds as direct application. Repeat application in 4 - 8 weeks or when necessary.

As a Roost Paint for control of poultry lice use 5 tbs. per gallon of water at rate of 1 pint per 150 ft. of roost. Repeat as needed.

As a premise treatment for control of poultry lice use 5 tbs. per 1 gallon of water. Apply liberally to all surfaces. Repeat as needed.

DOGS & CATS

DIRECT APPLICATION

Sprays

For control of fleas apply spray of 1 1/2 oz. of Malathion 50% Emulsifiable Liquid per 1 gal. of water. Wet the animal thoroughly using a hand sprayer.

If the animal has free run, treatments may have to be made every second or third week. To avoid unnecessary applications, animals should be closely examined for fleas before retreating. If the animal is confined to an area free of fleas, one treatment may be sufficient.

ANIMAL QUARTERS

Sprays

For control of fleas in kennels, pens, yards, lawns and under houses. Remove manure or debris.

Use spray of 8 ozs. of 50% Emulsifiable Liquid per 1 gal. of water at rate of 1 gal. per 1000 sq. ft. Use garden sprayer.

ORNAMENTALS

For Oyster Shell Scale, and Lace Bugs, use 1 1/4 pints. For Spider Mites, Aphids, White Flies, Mealybugs, Thrips, Japanese Beetle, European Pine Shoot Moth, Four-Lined Plant Bug, Tarnished Plant Bug, Rose Leafhopper, Potato Leafhopper, Scurfy Scale and Euonymus Scale use 1 3/4 pints. For Bagworm, Tent Caterpillar, Boxwood Leaf Miner, Birch Leaf Miner, Oak Kermes, Florida Red Scale, Fletcher Scale, Juniper, Azalea and Magnolia Scales, use 2 1/2 pints. For Black Scale Crawler, Soft Brown and Moterey Scales, use 3 pints. Apply thorough full coverage sprays as needed. Note: DO NOT apply to petunias, certain ferns (including Boston, Maidenhair, Pteris), and some species of Crassula, since plant injury may occur.

All above rates per 100 gallons of water

2043

ACCEPTED
JUL 21 1971
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR A DANGEROUS POISON REGISTERED
UNDER NO. SUBJECT
TO ATTACHED COMMENTS.

ACCEPTED
JUL 21 1971
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR A DANGEROUS POISON REGISTERED
UNDER NO. SUBJECT
TO ATTACHED COMMENTS.

ACCEPTED
JUL 21 1971
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR A DANGEROUS POISON REGISTERED
UNDER NO. SUBJECT
TO ATTACHED COMMENTS.

ACCEPTED
JUL 21 1971
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR A DANGEROUS POISON REGISTERED
UNDER NO. SUBJECT
TO ATTACHED COMMENTS.

ACCEPTED
JUL 21 1971
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR A DANGEROUS POISON REGISTERED
UNDER NO. SUBJECT
TO ATTACHED COMMENTS.

ACCEPTED
JUL 21 1971
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR A DANGEROUS POISON REGISTERED
UNDER NO. SUBJECT
TO ATTACHED COMMENTS.