



RAT-PEL WP-³/₈


KILLS RATS AND MICE

ACTIVE INGREDIENTS:

Warfarin: 30-A-Acetonil benzyl 4-hydroxycoumarin
Sulfaquinoxaline N'-Z-quinoxaliny sulfanilamide

INERT INGREDIENTS:

0.025%
0.025%
99.950%
100.000%



IMPORTANT NOTICE

The antibacterial agent increases the effectiveness of the product and assures thorough control of rats and mice. Especially recommended for rat and mouse infestations which are difficult to control.

DIRECTIONS FOR USE

RATS: Place 4 to 6 ounces of bait in dry containers next to burrows, in runways, or where rats are accustomed to feed. Assure an uninterrupted supply of bait for a period of not less than 12 days and continue baiting until all signs of feeding have stopped. Replace stale, damp or moldy baits with fresh bait.

MICE: Place tablespoonful amounts of bait in shallow containers where mice or signs of mice have been seen. Bait stations at intervals of 5 to 12 feet are recommended. Assure an uninterrupted supply of bait for a period of not less than 12 days.

CAUTION: Protect children, pets, and domestic animals from bait. If swallowed by humans, domestic animals, or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such case, intravenous and oral administration of vitamin K combined with blood transfusions are indicated as in the case of hemorrhage caused by overdoses of bishydroxycoumarin.

This rodenticide contains Warfarin, an anticoagulant chemical that reduces the clotting ability of the blood and, upon repeated feedings, causes fatal hemorrhage in rats.

CAUTION: Keep Out of Reach of Children
NET WEIGHT 50 POUNDS



RAT-PEL WP-³/₈


KILLS RATS AND MICE

ACTIVE INGREDIENTS:

Warfarin: 34.0% Acetonyl benzyl 4-hydroxycoumarin
Sulfaquinoxaline N'-Z-quinoxalinyll sulfanilamide

INERT INGREDIENTS:

0.025%
0.025%
99.950%
100.000%



IMPORTANT NOTICE

The antibacterial agent increases the effectiveness of the product and assures thorough control of rats and mice. Especially recommended for rat and mouse infestations which are difficult to control

DIRECTIONS FOR USE

RATS: Place 4 to 6 ounces of bait in dry containers next to burrows, in runways, or where rats are accustomed to feed. Assure an uninterrupted supply of bait for a period of not less than 12 days and continue baiting until all signs of feeding have stopped. Replace stale, damp or moldy baits with fresh bait

MICE: Place tablespoonful amounts of bait in shallow containers where mice or signs of mice have been seen. Bait stations at intervals of 8 to 12 feet are recommended. Assure an uninterrupted supply of bait for a period of not less than 12 days.

CAUTION: Protect children, pets and domestic animals from bait. If swallowed by humans, domestic animals, or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such case, intravenous and oral administration of vitamin K combined with blood transfusions are indicated as in the case of hemorrhage caused by overdoses of bishydroxycoumarin

This rodenticide contains Warfarin, an anticoagulant chemical that reduces the clotting ability of the blood and, upon repeated feedings, causes fatal hemorrhage in rats

CAUTION: Keep Out of Reach of Children
NET WEIGHT 50 POUNDS