

KILLS RATS AND MICE

ACTIVE INGREDIENTS:
Warfarin (3 (aplha-acetonylbenzyl)-4hydroxycoumarin)
INERT INGREDIENTS:

0.025% 99.975%

100.000%

DIRECTIONS FOR USE

RATS Place 4 to 6 ounces of bait in dry containers next to burrows, in runways, or where rats are accustomed to feed. Assure an uninterrupted supply of bait for a period of not less than 12 days and continue baiting until all signs of feeding have stopped. Replace stale, damp or moldy baits with fresh bait MICE: Place tablespoonful amounts of bait in shallow containers where mice or

MICE: Place tablespoonful amounts of bait in shallow containers where mice or signs of mice have been seen. Bait stations at intervals of 8 to 12 feet are recommended. Assure an uninterrupted supply of bait for a period of not less than 12 days.

CAUTION: Protect children, pets and domestic animals from bait. If swallowed by humans, domestic animals, or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such case, intravenous and oral administration of Vitamin K combined with blood, transfusions are indicated as in the case of hemorrhage caused by overdoses of bishydroxycoumarin.

DO NOT reuse container. Destroy when empty.

This rodenticide contains **an** anticoagulant chemical that reduces the clotting ability of the blood and, upon repeated feedings, causes fatal hemorrhage in rats

NOTICE: Seller's guaranty shall be limited to the terms of the label and subject thereto. The buyer assumes any risk blackage to persons or property arising out the sellor bundling, and accepts the product on these conditions.

CAUTION: KEEP OUT OF REACH OF CHILDREN
NET WEIGHT 50 POUNDS

USAG AFC NA

ACCEPTED

JAN 1 8 1003

UNDE A FICIDE
FUNGION AND CIDE ACT
FOR ECOLON BLANCE REGISTER
LE UNDES AND TAN AEGISTER
TO ATTACLED COMMENTS.

4726 44