



DSMA-30

LIQUID

FOR DALLIS GRASS CONTROL IN BERMUDA TURF LAWNS

ACTIVE INGREDIENTS:	
Disodium Methanearsonate	18.90%
INERT INGREDIENTS:	
TOTAL	100.00%
Total Arsenic, all in water soluble form	7.70%
*Equivalent to 30% Disodium Methanearsonate Hexahydrate	

CAUTION: KEEP OUT OF REACH OF CHILDREN.
SEE SIDE PANEL FOR OTHER CAUTIONS.

ACCEPTED

MAR-11-1968

UNDER THE FEDERAL INSECTICIDE, FUNGICIDE AND ROUENTICIDE ACT OR ECONOMIC POISON REGISTERED UNDER NO. 6720-59

Comment

USE IN ACCORDANCE WITH DIRECTIONS,
WARNING AND CAUTION

WARNING

DO NOT spray or allow to drift onto desirable flowers, trees, shrubs, vegetables or into lakes or streams. Coarse sprays are less likely to drift. DO NOT store near fertilizer, fungicides, insecticides or

CAUTION

DO NOT get in eyes or on the skin. If spilled on skin, wash off immediately with warm water and soap. Flush eyes with clear water. Avoid any inhalation of spray mist. Avoid contamination of feed, foodstuffs, streams or other water sources. Keep beyond the reach of children and domestic animals. Keep children and domestic animals off treated areas. Wash hands after using. Harmful if swallowed. DO NOT graze treated areas or feed livestock treated forage.

ANTIDOTE

If taken internally, induce vomiting and call a physician at once.

NOTICE

Seller's guaranty shall be limited to the terms of the label and subject thereto. The buyer assumes any risk of damage to persons or property arising out of use or handling, and accepts the product on these conditions.

MANUFACTURED BY
Southern MILL CREEK PRODUCTS COMPANY, Inc.
TAMPA FLORIDA

For control of Wild Mustard, Dandelion, Goldenrod, Ragweed, Plantains, Thistles, Docks, Wild Morning Glory, Shepherd's-purse, Wild Carrot, Mallow and certain other broad-leaved weeds and woody plants.

DIRECTIONS

"APPLY THIS PRODUCT ONLY AS SPECIFIED ON THIS LABEL"

Where sensitive species such as young Wild Mustard Predominate, the lower rates given may be satisfactory. For less sensitive species and where conditions for kill are difficult, use the higher rates of application. Apply at low pressures (30 to 50 pounds) during warm weather when weeds are young and growing actively.

MIXING: Add half the required amount of water to the spray tank, then add the ESTER WEED KILLER with agitation, and finally the balance of the water with continued agitation. **WARNING:** If ESTER WEED KILLER is to be used in straight oil mixtures, DO NOT let water get into the ESTER WEED KILLER itself nor into the finished mixture. **NOTE:** ESTER WEED KILLER in water forms an emulsion — not a solution — which tends to separate on standing. Provide agitation to prevent such separation and ensure uniformity of spray mixture.

WEED CONTROL IN PASTURES AND OTHER GRASSY AREAS: Use 2 to 3 pints per acre in sufficient water for adequate coverage. Clover and other legumes are usually injured by 2,4-D, especially if application is made when weather is hot. DO NOT use on Bent, St. Augustine, or Carpet Grasses, Dichondra, Clover, or new turf of any variety. Apply in warm weather when weeds are young and growing actively. Protect ornamentals from spray drift. For wild onion or garlic, apply in early spring each year for 2 or 3 years. DO NOT apply until grass is well established.

GENERAL WEED CONTROL: For weed control along roads, right-of-ways, fence rows, industrial yards and similar non-crop areas, use 2 quarts in 100 gallons of water. Make applications when foliage is fully expanded and plants are growing actively. Thoroughly wet plants with spray mixture. For weeds which are hard to control, repeat as needed.

WEED CONTROL IN CORN: Pre-emergence: Use 1 to 2 quarts per acre in sufficient water for adequate coverage. Use higher rate on heavy soil and lower rate on light soil. Make application within 3 days after planting. Heavy rains following application may cause injury to corn. **NOTE:** DO NOT make pre-emergence applications to corn in sandy soil.

Post-emergence: Use 1/3 to 1/2 pint per acre in sufficient water for adequate coverage. Application should be made only between the time of emergence and the time when corn reaches a height of 10 inches. Apply when most of the weeds have germinated but are still in the seedling stage. DO NOT treat corn grown for the production of hybrid seed with any 2,4-D formulation.

WEED CONTROL IN SMALL GRAINS (Wheat, Barley, Oats) Use 1/4 to 3/4 pint in sufficient water for adequate coverage. Use the lower rate for susceptible weeds such as mustard, and the higher rate for the resistant weeds such as thistle and wild buckwheat. Make application when grain is in full tiller stage (4 to 8 inches high). DO NOT use on flax, sorghum or rice, or grains underseeded to legumes unless injury to legumes can be tolerated.

WOODY PLANT CONTROL: For certain woody plants such as willow, elderberry, wild cherry, sumac and wild grape, use 3 quarts in 100 gallons of water for brush up to 6 feet tall. Make applications when foliage is fully expanded and plants are growing rapidly. Use a coarse spray and thoroughly wet foliage, branches and stems with spray mixture. Best results will usually be obtained from applications made soon after maximum foliage development in the spring. Applications may be made up to 3 weeks before frost under good growing conditions.



STANDARD ESTER WEED KILLER

Contains 3.34 Pounds of 2,4-D Acid Equivalent Per Gallon

ACTIVE INGREDIENTS:	
Isopropyl Ester of 2,4-Dichlorophenoxyacetic Acid*	47.8%
INERT INGREDIENTS:	52.2%
TOTAL	100.0%
*2,4-D Acid Equivalent	40.2%

DO NOT USE OR STORE NEAR HEAT OR OPEN FLAME

CAUTION: KEEP OUT OF REACH OF CHILDREN.

SEE SIDE PANEL FOR OTHER CAUTIONS.
U.S.D.A. Registration Number 6720-74

MANUFACTURED BY

Southern MILL CREEK PRODUCTS COMPANY, Inc
TAMPA FLORIDA

6720-74

RECEIVED
Aug 5 1970
UNDER THE FEDERAL INSECTICIDE
AND RODENTICIDE ACT
ECONOMIC POISON REGISTER-
SUBJECT