







5.000

3.25

91.75%

100.00°

SLUG AND SNAIL BANT PELLETS

ACTIVE INGREDIENTS: Tricalcium Arsenate Metaldehyde INERT INGREDIENTS

Total

Arsenic expressed as metallic 1.85 Arsenic in water soluble form expressed as metallic 0.035

DIRECTIONS

For use in nurseries and commercial ornamental plantings.

stugs, snalls, cutworms — Scatter pellets throughout infested area at a rate of ½ to ½ cupful per 100 sq. ft Ground should be wet. Sprinkle with water after application. Repeat as necessary

Make first application at planting time, before pests cause damage. Repeat at 2 to 3 week intervals.

Always wear protective gloves. DO NOT apply to foliage Bait should be conthe ground.

STRAWBERRY ROOT WEEVIL LARVAE ON ORNAMENTALS — Place several pellets around the base of each plant when adults begin to appear. Avoid touching plants with pellets Do not use on strawberries

NOTICE: Seller's guaranty shall be limited to the terms of the label and subject thereto. The buyer assumes any risk to persons or property arising out of use or handling, and accepts the product on these conditions.

ANTIDOTE STATEMENT:

Give one tablespoonful of salt in a glass of warm water and repeat until vomit is clear. Then give two tablespoonsful of Epsom Salts or Milk of Magnesia in water and plenty of milk and water. Have victim lie down and keep quiet.

CALL A PHYSICIAN IMMEDIATELY.

WARNING

MAY BE FATAL IF SWALLOWED. Avoid breathing dust. Avoid contact with skin, eyes or clothing. Wash thoroughly after using. Avoid contamination of feed and foodstuffs.

NOT FOR USE OR STORAGE IN OR AROUND THE HOME.

Keep out of reach of children and domestic animals.

May be attractive to dogs. Keep dogs off treated area

This product is toxic to wildlife. Birds and other wildlife in treated areas may be harmed Apply this product only as specified on this label. Do not contaminate water by cleaning of equipment, or disposal of wastes.

Destroy empty container DC NOT re-use

WARNING: KEEP OUT OF REACH OF CHILDREN