

6720-78

9-11-1970

1/3



SEVIN[®] 5

DUST

FOR GARDENS & PETS

ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N-methylcarbamate) . . . 5.0%

INERT INGREDIENTS 95.0%

TOTAL 100.0%

© Reg. T. M. of Union Carbide Corp. U. S. Patent No. 2-903-478

CAUTION: Keep Out Of Reach of Children.

SEE OTHER INFORMATION AND CAUTIONS ON BACK PANEL

U.S.D.A. Registration Number 6720-78

2/3

X-CEL SEVIN R 5 DUST is a new pesticide which will control many of the common insects infesting vegetables and ornamentals

VEGETABLE INSECT CONTROL

INSECTS	ON
Mexican Bean Beetle, Bean Leaf Beetle, Cucumber Beetle, Japanese Beetle, Leafhoppers	Green Beans, Lima Beans, Snap Beans, Cowpeas
Flea Beetles, Harlequin Bugs, Leafhoppers, Cabbage Worms, Corn Earworms, Tarnished Plant Bugs, Stink Bugs & Small Cabbage Loopers	Carrots, Collards, Corn, Lettuce, Spinach, Turnips
Cucumber Beetles, Flea Beetles, Leafhoppers, Squash Bugs	Cucumbers, Melons, Squash, Pumpkins, Canteloupes
Colorado Potato Beetle, Flea Beetles, Fall Armyworms, Lace Bugs, Tomato Fruitworm, Stink Bugs, Tarnished Plant Bugs	Potatoes, Tomatoes, Eggplant, Peppers, Okra
Flea Beetles, Harlequin Bugs, Cabbage Worms, Small Cabbage Loopers	Cabbage, Broccoli, Brussels Sprouts, Cauliflower

HOW TO USE

Apply thoroughly to infested plants on upper and lower leaf surfaces at first sign of damage and repeat as necessary. DO NOT apply within 3 days of harvest of head lettuce, cabbage, broccoli, brussels sprouts, cauliflower & turnips. DO NOT apply within 14 days of harvest on collards, leaf lettuce, and spinach. Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee populations.

ORNAMENTAL INSECT CONTROL

X-CEL SEVIN R 5 DUST will control Bagworms, Boxelder Bug, Lace Bugs, Leafhoppers, Rose Aphids, Leaf Rollers & Thrips (exposed).

On: Carnation, Gladioli, Zinnia, Chrysanthemum and other herbaceous and annual plants. Shrubs, trees and woody plants (Rose, Lilac, Juniper, Azalea, Elm, Maple, Dogwood, Birch, Pine, etc.)

HOW TO USE

Apply dust thoroughly to infested plants at first sign of damage and repeat at 7 day intervals or as necessary for good insect control.

PETS

Use on dogs and cats as follows: To control Brown Dog Ticks and Fleas apply liberally to animal and rub thoroughly into skin. Apply to feet between toes and all area where ticks and fleas occur. Repeat weekly if needed. Change bedding in sleeping quarters after treatment. DO NOT treat kittens or puppies less than 4 weeks old.

CAUTION

Harmful if swallowed or inhaled. Avoid breathing dust. Avoid contact with skin. Avoid contamination of feed or foodstuffs. Wash thoroughly after handling. Change clothing if contaminated.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your cooperative Agricultural Extension Service.

NOTICE: Seller's guaranty shall be limited to the terms of the label and subject thereto. The buyer assumes any risk to persons or property arising out of use or handling, and accepts the product on these conditions.

ACCEPTED

SEP 11 1970

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
ED UNDER NO. ~~6729-2~~ SUBJECT
TO ATTACHED COMMENTS