

6720-78

10-24-1975

1/2

NET CONTENTS:



ACTIVE INGREDIENTS:	
Carbaryl (1-naphthyl N-methylcarbamate)	5.0%
INERT INGREDIENTS	95.0%

Sevin is a Registered Trademark of Union Carbide Corp.
 U. S. Patent No. 2-903-478

Pest Control in & around Domestic Dwellings, Commercial & Industrial Areas, Medical Facilities, Schools: To control **Bees, Hornets and Wasps** - Locate nest in wall, bore 1/4 inch hole in wall at nest site, insert nozzle and inject dust. If hole is undesirable, sprinkle dust around entrance and pump dust into entrance. Do not eat honey from treated bee hives.

For **Fleas** - Sprinkle lightly on floors, rugs and furniture. Remove with a vacuum cleaner after an hour or two.

For control of **Earwigs, Fleas and Millipedes** around domestic dwellings:

Earwigs - Apply 10 to 15 lbs. per 5,000 sq. ft. Sevin 5% Dust to soil surface, mulch and debris around moist shady places.

Fleas - Apply 5 to 10 lbs. per 5,000 sq. ft. Sevin 5% Dust and apply liberally to pens, yards, lawns and under buildings around home. Force dust into cracks and crevices. Repeat as needed.

Millipedes - Apply 10 to 15 lbs. per 5,000 sq. ft. Sevin 5% Dust to soil surface, mulch and debris around moist shady places.

Mosquitoes (adults) - Apply 5 - 10 lbs. per 5,000 sq. ft. Sevin 5% Dust to lawns, shrubbery and other vegetation around the home. Repeat as needed.

Ticks - Apply 2 1/2 to 5 lbs. per 5,000 sq. ft. Sevin 5% Dust liberally to ground, debris and vegetation to a height of about 2 feet on lawns, vacant fields and wooded areas around the home. Repeat as needed.

DO NOT USE IN EDIBLE PRODUCTS AREAS OF FOOD PROCESSING PLANTS, RESTAURANTS OR OTHER AREAS WHERE FOOD IS COMMERICALLY PREPARED OR PROCESSED. DO NOT USE IN SERVING AREAS WHILE FOOD IS EXPOSED.

ORNAMENTAL INSECT CONTROL

SMCP SEVIN 5% DUST will control: Japanese, blister, flea and June beetles; boxelder, mealy, lace and plant bugs; leafhoppers, leaf roller, psyllid, rose aphid and exposed thrips on flowers and herbaceous plants. SMCP SEVIN 5% DUST will also control Gypsy moth, scale insects, bagworm, birch, boxwood and oak leaf miners, elm leaf aphid, Japanese, June, elm leaf and willow leaf beetles, mimosa webworm, tent and puss caterpillars, rose aphid and slug, periodical cicada, leafhopper, leaf roller, boxelder, lace mealy, plant and thorn bugs, apple aphid, orange tortox and exposed thrips on shrubs, trees and woody plants.

Apply dust thoroughly to infested plants at first sign of damage and repeat at 7 day intervals or as necessary for good insect control.

2/2

VEGETABLE INSECT CONTROL

INSECTS	ON
Mexican Bean Beetle, Bean Leaf Beetle, Cucumber Beetle, Japanese Beetle & Leafhoppers	Green Beans, Lima Beans, Snap Beans, Cowpeas
Flea Beetles, Harlequin Bugs, Leafhoppers, Cabbage Worms, Corn Earworms, Tarnished Plant Bugs, Stink Bugs & Small Cabbage Loopers	Carrots, Collards, Corn, Lettuce, Spinach, Turnips
Cucumber Beetles, Flea Beetles, Leafhoppers, Squash Bugs	Cucumbers, Melons, Squash, Pumpkins, Cantaloupes
Colorado Potato Beetle, Flea Beetles, Fall Armyworms, Lace Bugs, Tomato Fruitworm, Stink Bugs, Tarnished Plant Bugs	Potatoes, Tomatoes, Eggplant, Peppers, Okra
Flea Beetles, Harlequin Bugs, Cabbage Worms, Small Cabbage Loopers	Brussels Sprouts, Cauliflower, Cabbage, Broccoli

HOW TO USE: Apply thoroughly to infested plants on upper and lower leaf surfaces at first sign of damage and repeat as necessary. DO NOT apply within 3 days of harvest on head lettuce, cabbage, broccoli, Brussels sprouts, cauliflower and turnips. DO NOT apply within 14 days of harvest on collards, leaf lettuce and spinach. Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee populations.

**PETS
DOGS & CATS**

For **Fleas, Lice and Ticks:** Dust liberally over animal and rub thoroughly in to skin. For brown dog tick be sure to also apply on legs, feet and between toes. Comb out dead ticks a few hours after treatment. Keep dust out of eyes. Apply in and around sleeping quarters, kennels and dog houses. Force dust into cracks and crevices. Repeat at weekly intervals or as needed. DO NOT treat kittens or puppies less than 4 weeks old.

CAUTION

Harmful if swallowed or inhaled. Avoid breathing dust. Avoid contact with skin. Avoid contamination of feed or foodstuffs. Wash thoroughly after handling. Change clothing if contaminated. Wear regular long sleeved clothing.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your cooperative Agricultural Extension Service.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on label.

Do not reuse empty container. Destroy when empty.

NOTICE: Seller's guaranty shall be limited to the terms of the label and subject thereto. The buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

CAUTION: KEEP OUT OF REACH OF CHILDREN
E.P.A. Reg. No. 6720-78 E.P.A. Est. No. 6720-FL-1

MANUFACTURED BY

Southern Mill Creek

PRODUCTS COMPANY, INC.

TAMPA, FLORIDA 33601