

GENERAL INFORMATION

Methoxychlor 2E Emulsifiable Insecticide contains 2 pounds technical methoxychlor per gallon, and is to be used as a spray with water (forming a stable emulsion) for the control of certain insects. It is suitable for high- or low-volume application.

DIRECTIONS

Methoxychlor 2E should be used only in accordance with recommendations on this label.

FORAGE CROPS — (Alfalfa, clover, grass, soybean cowpea and peanut forage) — Use the specified dosages of Methoxychlor 2E in a minimum of 100 gallons of water per acre on the listed forage crops and potatoes by ground equipment. Start applications at first sign of infestation and repeat at 7- to 14-day intervals as needed. DO NOT apply later than 7 days before harvest or grazing.

Leafhopper, spittlebug 1½ to 2 qts. per acre
Alfalfa caterpillar, flea beetles 2 qts. per acre

*Alfalfa weevil (larvae), alfalfa webworm, fall armyworm, clover leaf weevil, Mexican bean beetle, pea weevil, soybean caterpillar, velvet bean caterpillar 2 to 3 qts. per acre

Cowpea curculio, Japanese beetle 3 qts. per acre

Armyworm 3 to 4 qts. per acre


*For pea aphid and as an alternate program for control of alfalfa weevil, add malathion (emulsifiable liquid) at the rate of **1 to 1¼ lbs. per acre.**

POTATOES — Use **2 qts. per acre** for control of Colorado potato beetle, leafhoppers, and flea beetles. Re-treat as necessary.

LIVESTOCK — (except dairy animals) — For control of certain external parasites. --

Use as spray. DO NOT use in dipping vats.

Cattle — Use **1 qt.** Methoxychlor 2E to **12½ gals.** of water. Apply **2 qts.** of spray per mature animal for control of hornflies; for control of short-nosed and long-nosed cattle lice, apply as a drenching spray.



ACTIVE INGREDIENTS
*Methoxychlor
Methylparathion
INERT INGREDIENTS
T
*24 methoxychlor, technical
3% other isomers

IF EXPOSED TO
STORAGE (50 lb. container)

CAUTION



ACCENT

July 14 1970

UNDER THE FEDERAL INSECTICIDE
AND RODENTICIDE ACT
AND POISON REGISTER
REGISTRATION NUMBER 6720-94
REGISTERED COMMENTS



6720-94

METHOXYCHLOR 2E

EMULSIFIABLE CONCENTRATE

Contains 24% Technical Methoxychlor

| | |
|--------------------------|------|
| ACTIVE INGREDIENTS | 93% |
| *Methoxychlor, Technical | 24% |
| Methylated Naphthalenes | 69% |
| INERT INGREDIENTS | 7% |
| TOTAL | 100% |

*24 methoxychlor, technical is equivalent to 21% 2, 2-bis (p-methoxyphenyl) 1,1, 1-trichloroethane and 3% other isomers and reaction products.

IF EXPOSED TO PROLONGED COLD, PLACE IN WARM STORAGE (50 to 80° F.) FOR SEVERAL HOURS AND AGITATE BEFORE USING.

CAUTION: KEEP OUT OF REACH OF CHILDREN.

SEE SIDE PANEL FOR ADDITIONAL CAUTIONS.
U.S.D.A. Registration Number 6720-94

Hogs — For control of lice, use 1 gal. of 2E to 12½ gals. of water applied as spray.

SELF APPLICATOR FLY CONTROL (except dairy animals) — For control dilute 1 gal. Methoxychlor 2E with 4 gal. fuel or furnace type oil. DO NOT use kerosene type of oil. Use 1 gal. of solution to 10 gal. of water on "back rubber"; soak burlap every 2-3 weeks.

GRAIN STORAGE BINS — To clean out empty storage bins, use 1 gal. M in 10 gals. of water applied at the rate of 1 spray per 1,000 sq. ft. before production. This aids in control of cadelle, flat grain beetle, toothed grain beetle, lesser grain beetle, long-headed flour beetle, rice weevil, confused flour beetle, and hairy fungus. NOTE - DO NOT use in bins for at least 24 hours or until well out thoroughly.

CONTACT AND SPACE SPRAY FOR reduce populations of houseflies and other flies in farm buildings, in barnyards, in alleys, and other outdoor locations, dilute 1 gal. of 2E with 2½ gals. of water and apply by mist blower or directly to flies.

MANUFACTURED BY
Southern MILL CREEK PRODUCTS COMPANY, Inc.
TAMPA FLORIDA

Hogs — For control of lice, use **1 qt.** Methoxychlor 2E to **12½ gals.** of water applied as a drenching spray.

SELF APPLICATOR FLY CONTROL ON CATTLE — (except dairy animals) — For control of hornflies dilute **1 gal.** Methoxychlor 2E with **4 gals.** of Diesel fuel or furnace type oil. **DO NOT** use any other type of oil. Use **1 gal.** of solution to saturate burlap on "back rubber"; soak burlap every two to three weeks.

GRAIN STORAGE BINS — To clean up infestations in empty storage bins, use **1 gal.** Methoxychlor 2E in **10 gals.** of water applied at the rate of **2 gals. of spray per 1,000 sq. ft.** before products are stored. This aids in control of cadelle, flat grain beetle, saw-toothed grain beetle, lesser grain borer, red flour beetle, long-headed flour beetle, rice weevil, granary weevil, confused flour beetle, foreign grain beetle and hairy fungus. **NOTE** — **DO NOT** add grain to bin for at least 24 hours or until walls have dried out thoroughly.

CONTACT AND SPACE SPRAY FOR FLIES — To reduce populations of houseflies and stable flies in farm buildings, in barnyards, in alleys, on dumps and other outdoor locations, dilute **1 pt. of** Methoxychlor 2E with **2½ gals.** of water and apply in space by mist blower or directly to flies by hydraulic

sprayer, hand sprayer or cylindrical pressure sprayer. Care should be taken to avoid contamination of milk handling equipment, feedstuffs, feed troughs and watering receptacles. Dairy animals should be present while spraying.

COMPATIBILITY — **DO NOT** use this product with any spray materials which are incompatible with it. **DO NOT** apply within 14 days of sulfur or sulfur product applications.

CAUTION!

Avoid prolonged or repeated contact with skin. Avoid prolonged or repeated breathing of spray mist.

Avoid contamination of feed and foodstuffs. Harmful if swallowed.

Keep out of reach of children.

IMPORTANT: Users should observe the usual cautions concerning application of oil-containing products to crops during high temperatures, since crops are then more susceptible to oil injury. This product is toxic to fish. Keep out of any body of water. Do not contaminate water by cleaning equipment, or disposal wastes.

Destroy empty container. **DO NOT** re-use.

NOTICE: Seller's guaranty shall be limited to terms of the label and subject thereto. The buyer assumes any risk to persons or property arising out of use or handling, and accepts the product on these conditions.

2E

93%
24%
69%
7%
100%

1,1, 1-trichloroethane and

**WARM
E BEFORE USING.**

CHILDREN.

PANY, Inc.