

SMCP ZINEB

An Effective Fungicide

ACTIVE INGREDIENTS:

Zineb (Zinc Ethylenebisdithiocarbamate).....75%
INERT INGREDIENTS:.....25%
TOTAL.....100%

ORNAMENTAL GRASSES

Directions for use on ornamentals:

SMCP Zineb will control Helminthosporum and Curvularia diseases on Bermuda, Zoysia and Bahia grasses. Apply at the rate of 2 to 4 ounces (7 to 14 tablespoonsful) in 3 to 5 gallons of water per 1000 sq. ft. Begin application before diseases are expected to occur and repeat at 7 to 10 day intervals as necessary.

Do not graze treated areas or feed clippings to livestock.

DIRECTIONS FOR USE ON ORNAMENTALS

Table with columns for Diseases and FLOWERS. Includes handwritten notes like 'DEC 16, 1974' and '6720-108'.

ROOFS

Infested areas should be cleaned prior to treatment. To control mildew on Asbestos, Burma, Composition, Marble Chip, Slate, Stone or Tile Roofs use 2 to 4 tablespoonsful per gallon of water. 1 gallon of solution will cover 200 to 300 sq. ft. of surface. Repeat at first sign of mildew.

CAUTION: Combustible mixture. May cause irritation of nose and throat. DO NOT breathe dust of spray mist.

Do not reuse container. Destroy when empty.

NOTICE: It is recommended that... (small text)

CAUTION: KEEP OUT OF REACH OF CHILDREN

E.P.A. Reg. No. 6720-108



RECOMMENDATIONS

The dosage rates on this label indicate pounds of Zineb per 100 gal. of dilute spray. Use 100 to 125 gal. spray per acre for low crops such as beans, carrots and onions. Use 150 to 200 gal. spray per acre for large, heavier crops such as cucumber, and tomato vines. Make thorough, full coverage applications.

FRUIT AND NUTS

APPLE: Apple Blotch, Black Rot, Botryosphaeria Fruit Rot, Brood Spot, Cedar Rust, Fly Speck, Frogeye, Quince Rust, Scab, Sooty blotch - Use 1 to 2 lbs. Begin applications in delayed dormant and continue through cover sprays. Use 1/2 to 1 lb. when combined with Captan. DO NOT use within 30 days of harvest.

APPLE, PEAR: Fire Blight - Use 2 lbs. at 2 to 3 day intervals during periods of bloom-opening and wet weather. Maintain recommended sanitary pruning practices. DO NOT apply within 7 days of harvest.

APRICOT, PEACH: Coryneum Blight (Shot Hole), Peach Leaf Curl - Use 2 lbs. in postharvest or dormant sprays and in popcorn to jacket stages according to local spray program.

CITRUS: Rust Mite and associated Russetting and Greasy Spot - Use 1 lb. Apply with thorough coverage. Make application from late June through early August, or earlier if rust mite begins to show.

SOUR CHERRY: Leaf Spot or Shot Hole - Use 1 lb. Apply at petal fall and 10 and 20 days after petal fall and, immediately harvest. DO NOT apply within 7 days of harvest.

CURRENT: Leaf Spot - Use 2 lbs. Apply 2 to 3 weeks after bloom. DO NOT apply within 7 days of harvest.

GRAPE: Black Rot, Downy Mildew, Ripe Rot - Use 1 1/2 lbs. just before bloom, 10 days after bloom, and 2 to 3 weeks later. DO NOT apply within 7 days of harvest.

STRAWBERRY: Leaf Spot - Use 1 1/2 to 2 lbs. Begin applications when new growth starts in spring and repeat at 7 to 10 day intervals until fruit begins to form. Leaf Scorch - Use 1 1/2 to 2 lbs. in new bed in late summer. Make 2 to 3 applications at 7 day intervals. DO NOT apply with 7 days of harvest.

VEGETABLES

Use 1 1/2 to 2 lbs. per 100 gal. of dilute spray. Begin applications at first signs of disease and repeat at 7 to 10 day intervals or as needed to maintain control of the disease stated.

BEANS: Anthracnose - Begin when first blossoms appear. Downy Mildew, Rust - Begin when disease threatens. DO NOT apply within 7 days of picking.



BEETS, KALE, MUSTARD GREENS, SWISS CHARD, TURNIP: Cercospora Leaf Spot, Downy Mildew - DO NOT apply within 7 days of harvest. Do not apply on Kale within 10 days of harvest.

CARROT: Alternaria, Cercospora, Leaf Spot - DO NOT apply within 7 days of harvest if tops are to be used as food or feed.

CELERY: Early Blight, Late Blight - Begin applications when plants emerge in the plant beds. Repeat at 3 to 5 day intervals until plants are set in field. Continue applications at 7 day intervals. DO NOT apply within 14 days of harvest.

CUCURBITS: (Cantaloupe, Cucumber, Melons, Squash) - Alternaria (Macrosporium) Blight, Anthracnose, Downy Mildew, Gummy Stem Blight - Direct spray to underside of leaves. DO NOT apply within 5 days of harvest.

LETTUCE: Downy Mildew - DO NOT apply within 7 days of harvest on head lettuce or 10 days on leaf lettuce.

ONION: Botrytis Leaf Blight (Blast), Downy Mildew, Purple Blotch - DO NOT apply within 7 days of harvest on green or spring onions.

SPINACH: Downy Mildew (Blue Mold), White Rust - DO NOT apply within 10 days of harvest.

TOMATO: Anthracnose, Cladosporium Gray Mold, Early Blight, Late Blight, Septoria Leaf Spot, Stemphyllium Leaf Spot. DO NOT apply within 5 days of harvest.