



REGISTERED  
 3-26-70  
 UNDER THE FEDERAL INSECTICIDE,  
 FUNGICIDE AND RODENTICIDE ACT  
 FOR A DANGEROUS PESTICIDE  
 TO WHICH 6720-112

# SPECIAL ESTER LV D&T

Contains the Equivalent of 2 Pounds of 2,4-D Acid  
 and 2 Pounds of 2,4,5-T Per Gallon

ACTIVE INGREDIENTS:	
Isooctyl Ester of 2,4-Dichlorophenoxyacetic Acid	33.7%
Isooctyl Ester of 2,4,5-Trichlorophenoxyacetic Acid	32.1%
INERT INGREDIENTS:	34.2%
TOTAL	100.0%
2,4-Dichlorophenoxyacetic Acid equivalent	22.3%
2,4,5-Trichlorophenoxyacetic Acid equivalent	22.3%

**CAUTION: KEEP OUT OF REACH OF CHILDREN.**

SEE SIDE PANEL FOR OTHER CAUTIONS.  
 U.S.D.A. Registration Number 6720-112



MANUFACTURED BY:

**Southern MILL CREEK PRODUCTS COMPANY, Inc.**  
 TAMPA FLORIDA

ACCEPTED

3-16-70

UNDER THE FEDERAL INSECTICIDE, FUNGICIDE AND RODENTICIDE ACT FOR THE CONTROL OF PESTS REGISTERED UNDER NO. 6720-112

**STUMP AND BASAL TREATMENTS:**

These treatments are also known as dormant treatments. They may be used effectively at any time of the year but best results are usually obtained with late winter applications.

**STUMP TREATMENT:**

Use 1 to 2 gallons in 25 gallons of diesel oil. Spray or brush freshly cut surfaces and sides of stumps thoroughly until run-off occurs. Foliage treatment should be used for sprout growth.

**BASAL TREATMENT:**

Use 1 to 2 gallons in 25 gallons of diesel oil or kerosene. Spray bark of plant from soil surface to about 2 feet up and completely around stem or trunk. Bark, especially at the ground line, should be thoroughly sprayed to point of run-off. For plants with trunks more than 6 inches in diameter, overlapping cuts should be made through the bark around the trunk and spray should be directed into the cuts when the bark is treated.

**NOTICE**

Seller's guaranty shall be limited to the terms of the label and subject thereto. The buyer assumes any risk of damage to persons or property arising out of use or handling, and accepts the product on these conditions.

**WARNING!**

DO NOT allow this product to contact in any manner Legumes, Cotton, Grapes, Fruit Trees, Vegetables, Flowers, Ornamentals or other desirable plants either in the growing or dormant periods by direct application, spray drift, or vapors. Susceptible plants growing in the immediate vicinity may be injured by vapors from this product under certain conditions. Use coarse sprays as they are less likely to drift. DO NOT use on lawns. Spraying of this product may be subject to restrictions imposed by state or local laws. Seed germination or plant growth may be temporarily inhibited by excessive amount of this product in the soil.

DO NOT permit this product or spray drift to contaminate irrigation ditches or domestic water supplies.

Equipment used for the application or handling of any SPECIAL ESTER LV D&T spray material should not be used for applying other agricultural chemicals due to the difficulty of removing all residues from such equipment. This container cannot be satisfactorily decontaminated and must not be reused or sold for reuse.

**NOTE:** If this material has been exposed to below-freezing temperatures it should be warmed to at least 40°F. and thoroughly mixed before it is used.

**DO NOT STORE NEAR FERTILIZERS, SEEDS, INSECTICIDES, FUNGICIDES, FEED OR FOODSTUFFS**

**CAUTION!**

**HARMFUL IF SWALLOWED**

**MAY CAUSE SKIN IRRITATION. AVOID CONTACT**

**WITH EYES, SKIN AND CLOTHING.**

**AVOID INHALATION.**

**USES:** For use on brush, deciduous woody plants and tree sprouts in range lands, right-of-ways, stump lands and similar land areas where such growth is not wanted. SPECIAL ESTER LV D&T is not intended for weed control in growing crops and should not be used where damage to desirable plants might occur from vapors, spray drift or direct application.

**For Use on the Following Woody Plants:**

- |                |                |              |
|----------------|----------------|--------------|
| ALDER          | ELDERBERRY     | PINE*        |
| ASH*           | ELM            | POISON IVY   |
| ASPEN          | GREEN BRIAR*   | POISON OAK   |
| AM. CRAB APPLE | HACKBERRY*     | POPLAR       |
| BARBERRY       | HAWTHORN       | RASPBERRY    |
| BIRCH*         | HAZEL          | ROSE         |
| BLACKBERRY     | HERCULES' CLUB | SAGEBRUSH    |
| BLACK GUM*     | HICKORY*       | SALT CEDAR*  |
| BLACK LOCUST   | HONEY LOCUST   | SASSAFRAS    |
| BOX ELDER      | HONEYSUCKLE    | SERVICEBERRY |
| BUCKBRUSH      | HORNBEAM       | SILVERBERRY  |
| CATALPA        | HORSE CHESTNUT | SKUNKBRUSH   |
| CHESTNUT       | JUNEBERRY      | SHADBUSH     |
| CHINABERRY     | MAPLE*         | SPICEBUSH    |
| CHOKE CHERRY   | MULBERRY       | SUMAC        |
| COTTONWOOD     | OAK*           | SWEETFERN    |
| CURRANT        | OSAGE ORANGE   | SWEET GUM    |
| DOGWOOD*       | PERSIMMON*     | WILD CHERRY  |
|                |                | WILLOW       |

**DIRECTIONS**

**MIXING:** Add the recommended amount of SPECIAL ESTER LV D&T to about one-half the volume of water or oil to be used in spraying. Mix well, then add remaining water or oil and mix until spray mixture is uniform. Continue mixing while spraying. DO NOT let water get into brush holes or spray used in stump or basal treatment.

**FOLIAGE TREATMENT**

For plants less than 6 feet tall, use 3 to 4 quarts in 100 gallons of water (approximately 1 cupful in 5 gallons of water). Make applications, using a coarse spray, when leaves are fully developed and plants are growing rapidly. Thoroughly wet foliage branches and stems. Repeat applications in 4 to 6 weeks if regrowth makes it necessary. Foliage treatment is not recommended for plants more than 6 feet tall.

\*For best results on these plants use only the higher rates given in the directions.



SPECIMEN LABEL



3-26-70

6720-112

# SPECIAL ESTER LV D&T

Contains the Equivalent of 2 Pounds of 2,4-D Acid  
and 2 Pounds of 2,4,5-T Per Gallon

ACTIVE INGREDIENTS		
Isooctyl Ester of 2,4-Dichlorophenoxyacetic Acid		33.7%
Isooctyl Ester of 2,4,5-Trichlorophenoxyacetic Acid		32.1%
INERT INGREDIENTS		
TOTAL		100.0%
2,4-Dichlorophenoxyacetic Acid equivalent*	22.3%	
2,4,5-Trichlorophenoxyacetic Acid equivalent	22.3%	

**CAUTION: KEEP OUT OF REACH OF CHILDREN.**

SEE SIDE PANEL FOR OTHER CAUTIONS  
U.S.D.A. Registration Number 6720-112

MANUFACTURED BY  
**Southern MILL CREEK PRODUCTS COMPANY, Inc.**  
TAMPA FLORIDA

ACCEPTED

3-16-70

UNION ...

6720-112

**STUMP AND BASAL TREATMENTS:**

These treatments are also known as dormant treatments. They may be used effectively at any time of the year but best results are usually obtained with late winter applications.

**STUMP TREATMENT**

Use 1 to 2 gallons in 25 gallons of diesel oil. Spray or brush freshly cut surfaces and sides of stumps thoroughly until run-off occurs. Foliage treatment should be used for sprout growth.

**BASAL TREATMENT**

Use 1 to 2 gallons in 25 gallons of diesel oil or kerosene. Spray bark of plant from soil surface to about 2 feet up and completely around stem or trunk. Bark, especially at the ground line, should be thoroughly sprayed to point of run-off. For plants with trunks more than 6 inches in diameter, overlapping cuts should be made through the bark around the trunk and spray should be directed into the cuts when the bark is treated.

**NOTICE**

Seller's guaranty shall be limited to the terms of the label and subject thereto. The buyer assumes any risk of damage to persons or property arising out of use or handling and accepts the product on these conditions.

**WARNING!**

DO NOT allow this product to contact in any manner Legumes, Alfalfa, grapes, fruit trees, Vegetables, Flowers, Ornamentals or other desirable plants either in the growing or dormant periods by direct application, spray drift, or vapors. Susceptible plants growing in the immediate vicinity may be injured by vapors from this product under certain conditions. Use coarse sprays as they are less likely to drift. DO NOT use on lawns. Spraying of this product may be subject to restrictions imposed by state or local laws. Seed germination or plant growth may be temporarily inhibited by excessive amount of this product in the soil. DO NOT permit this product or spray drift to contaminate irrigation ditches or domestic water supplies.

Equipment used for the application or handling of any SPECIAL ESTER LV D&T spray material should not be used for applying other agricultural chemicals due to the difficulty of removing all residues from such equipment. This container cannot be satisfactorily decontaminated and must not be reused or sold for reuse. NOTE: If this material has been exposed to below freezing temperatures it should be warmed to at least 40 F. and thoroughly mixed before it is used.

**DO NOT STORE NEAR FERTILIZERS, SEEDS, INSECTICIDES, FUNGICIDES, FEED OR FOODSTUFFS**

**CAUTION!**

**HARMFUL IF SWALLOWED**

**MAY CAUSE SKIN IRRITATION. AVOID CONTACT**

**WITH EYES, SKIN AND CLOTHING.**

**AVOID INHALATION.**

**USES:** For use on brush, deciduous woody plants and tree sprouts in range lands, right-of-ways, stump lands and similar land areas where such growth is not wanted. SPECIAL ESTER LV D&T is not intended for wood control in growing crops and should not be used where damage to desirable plants might occur from vapors, spray drift or direct application.

**For Use on the Following Woody Plants:**

- |                |                |              |
|----------------|----------------|--------------|
| ALDER          | ELDERBERRY     | PINE*        |
| ASH*           | ELM            | POISON IVY   |
| ASPEN          | GREEN BRIAR*   | POISON OAK   |
| AM. CRAB APPLE | HACKBERRY*     | POPULAR      |
| BARBERRY       | HAWTHORN       | RASPBERRY    |
| BIRCH*         | HAZEL          | ROSE         |
| BLACKBERRY     | HERCULES-CLUB  | SAGEBRUSH    |
| BLACK GUM*     | HICKORY*       | SALT CEDAR*  |
| BLACK LOCUST   | HONEY LOCUST   | SASSAFRAS    |
| BOX ELDER      | HONEYSUCKLE    | SERVICEBERRY |
| BUCKRUSH       | HORNBEAM       | SILVERBERRY  |
| CATALPA        | HORSE CHESTNUT | SKUNKBRUSH   |
| CHESTNUT       | JUNEBERRY      | SHADBUSH     |
| CHINABERRY     | MAPLE*         | SPICEBUSH    |
| CHOKE CHERRY   | MULBERRY       | SUMAC        |
| COTTONWOOD     | OAK*           | SWEETFERN    |
| CURRANT        | OSAGE ORANGE   | SWEET GUM    |
| DOGWOOD*       | PERSIMMON*     | WILD CHERRY  |
|                |                | WILLOW       |

**DIRECTIONS**

**MIXING:** Add the recommended amount of SPECIAL ESTER LV D&T to about one half the volume of water or oil to be used in spraying. Mix well then add remaining water or oil and mix until spray mixture is uniform. Continue mixing while spraying. DO NOT let water get into brush killer oil spray used in stump or basal treatment.

**FOLIAGE TREATMENT**

For plants less than 6 feet tall, use 3 to 4 quarts in 100 gallons of water (approximately 1 cupful in 5 gallons of water). Make applications using a coarse spray, when leaves are fully developed and plants are growing rapidly. Thoroughly wet foliage, branches and stems. Repeat applications in 4 to 6 weeks if regrowth makes it necessary. Foliage treatment is not recommended for plants more than 6 feet tall.

\*For best results on these plants use only the higher rates given in the directions.