



SEVIN* 10 DUST

ACCEPTED

APR 7 1971

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC PEERON REGISTERED
UNDER NO. SUBJECT
TO ATTACHED COMMENTS.

6720-136

FOR USE BY PROFESSIONAL PEST CONTROL OPERATORS

ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N-methylcarbamate) 10.0%

INERT INGREDIENTS: 90.0%

TOTAL 100.0% ✓

CAUTION: KEEP OUT OF REACH OF CHILDREN

DIRECTIONS

For use by Pest Control Operators outside of buildings for the control of Ants, Brown Dog Ticks, Earwigs and Millipedes.

Dust ant trails, cracks and crevices outside of buildings. Also, try to locate nest and dust it. For centipedes and millipedes dust under stones, flower pots, boards and other debris. For the control of earwigs dust during late afternoon or early evening in a band around premises or structures. Also treat base of mulched shrubbery and flower beds. This band of dust may also prevent Brown Dog Ticks from entering buildings.

RECOMMENDATIONS

FOR INSECT CONTROL ON PLANTS

Dosages given are in amount of Sevin 10% Dust per acre. Apply when insects first threaten damage to crop and repeat at 5 to 7 day intervals as required to maintain control.

COTTON: Boll Weevil, Bollworm, Cotton Fleahopper, Lygus Bugs, Pink Bollworm, Thrips. For Early Season control use 15 lbs. per acre. Mid-Season use 15 to 20 lbs. per acre. Late Season use 20 to 25 lbs. per acre.

PEANUTS: Corn Earworm, Fall Armyworms, Potato Leafhoppers, Velvet Bean Caterpillars - Use 15 lbs. per acre. Apply when larvae are young on Corn Earworms, and Fall Armyworms. Apply when feeding by larvae is noted on Velvet Bean Caterpillars. Apply about July 10 and then at three week intervals on leafhoppers.

TOBACCO: (Plant Bed): Tobacco Flea Beetles, Green June Beetle Larvae - use 2½ lbs. per 100 sq. yd. Apply when plants appear. Repeat as needed at 10 to 14 day intervals and just before transplantings.

TOBACCO: (Field): Hornworms, Budworms, Flea Beetles, June Beetles, Japanese Beetles - use 25 lbs. per acre. Good coverage is essential. Repeat as needed. Allow 3 days before priming or cutting.

USE THIS PRODUCT ONLY AS DIRECTED ON THIS LABEL

CAUTION

Harmful if inhaled or swallowed. DO NOT breathe dust. DO NOT take internally. Avoid breathing dust.

Avoid contact with skin or eyes. Wash hands thoroughly after handling and before eating or smoking. Take shower or bath after work. Wear regular long sleeved work clothing. Wash clothing in hot water and soap before re-using.

OBSERVE FOLLOWING USE PRECAUTIONS

Toxic to honey bees. DO NOT apply during bloom when honey bees or other pollinators are active.

Avoid drift to adjoining food and forage crops.

Consult State Agricultural Extension Service for additional information, as the timing, number and rates of applications needed will vary with local conditions. Apply only as specified on label.

NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

NOTICE: Seller's guaranty shall be limited to the terms of the label and subject thereto. The buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

* Trademark of Union Carbide Corp.
U.S. Pat. No. 2-903-478

U.S.D.A. Registration Number 6720-136

NET WEIGHT:

POUNDS



MANUFACTURED BY

Southern MILL CREEK PRODUCTS COMPANY, Inc.

TAMPA, FLORIDA