


6720-154

08/01/1974

1/3

CENTER PANEL

Best Available copy



FOR USE BY PEST
For the
A Product

D/227
INSECTICIDE
POISON

OPERATE
each
and

ACTIVE INGREDIENTS:

Chlorpyrifos (O,O-diethyl O-(3,5,6-trichloro-2-pyridyl) phosphorothionate).....	12.0%
2,2-dichlorovinyl dimethyl phosphate.....	11.2%
Related Compounds.....	0.8%
Aromatic Petroleum Derivative Solvents.....	58.0%
INERT INGREDIENTS.....	18.0%
TOTAL.....	100.0%

DANGER: KEEP OUT OF REACH OF CHILDREN

See side panels for warnings, and first aid statement and other information
E.P.A. Registration Number 6720-154

COCKROACH CONTROL

RESISTANT AND NON-RESISTANT COCKROACHES

Dilute the required amount of D-217 in water. Mix according to the following table to obtain the desired concentration of 0.25% or 0.50% D/V 217 insecticide. Use higher concentration (0.50%) to quickly reduce heavy cockroach infestations and the lower concentration (0.25%) on light infestations.

Amount of D/V 217 to use

Amount of Spray Mixture Desired	for 0.25% spray	for 0.50% spray
1 gallon	2 1/4 ounces	5 1/4 ounces
3 gallons	8 ounces	16 ounces
5 gallons	12 ounces	24 ounces
10 gallons	24 ounces	48 ounces

SPOT APPLICATION (NON-FOOD AREAS)

Use as a spot treatment by applying as a coarse spray or with a paint brush, to areas where cockroaches normally are found. These areas include dark corners of rooms and closets, at the base of walls, in cracks in the wall, behind baseboards, behind and beneath sinks, refrigerators, stoves and cabinets, and around drainpipes. Apply in a manner to form spray or mist contacting food, foodstuffs and water supplies. Dishes and food handling utensils should be thoroughly washed in soap and water if they have become contaminated by the application of this product. Repeat treatment as needed.

APPLICATIONS WITHIN FOOD AREAS OF FOOD HANDLING ESTABLISHMENTS

Limited to Crack and Crevice Treatment Only - Apply a small amount of material directly into cracks and crevices such as expansion joints and the floor, wall voids, motor housing, junction boxes or switch boxes, conduits or hollow equipment legs where cockroaches hide. Care should be taken to avoid depositing the product onto exposed surfaces or introducing the material into the air. Equipment that can deliver a pin stream of insecticide must be used for crack and crevice treatment. Use 1/2% spray. Avoid contamination of food or food processing surfaces.

Applications of this product in the food areas of food handling establishments, other than as a crack and crevice treatment, are not permitted.

NOTE: A period of 4 to 7 days of treatment is normally required for maximum effect on cockroaches.

*DURSBAN - Trademark of the Dow Chemical Company

DANGER

POISONOUS IF SWALLOWED, INHALED OR ABSORBED THROUGH SKIN AND EYES. DO NOT get on skin or in eyes. DO NOT breathe vapor or spray mist. Keep away from food, feedstuff and water supplies. Keep container closed. Keep away from heat or open flame.

Wear natural rubber gloves, protective clothing and goggles. In case of contact wash immediately with soap and water.

Wear mask or respirator of a type found by the U. S. Bureau of Mines to give adequate protection against this material.

Wash hands, arms and face with soap and water after handling and before eating and smoking.

DO NOT wear contaminated clothing. Wash all contaminated clothing with soap and hot water before reuse.

KEEP CHILDREN AND PETS OUT OF TREATED AREAS UNTIL SURFACES ARE DRY. This product is toxic to fish and wildlife. DO NOT contaminate water by cleaning of equipment, or disposal of wastes.

ANTIDOTE & FIRST AID TREATMENT

WARNING SYMPTOMS: Symptoms include weakness, headache, tightness in chest, blurred vision, non-reactive pin point pupils, salivation, sweating, nausea, vomiting, diarrhea, abdominal cramps.

Keep patient prone and quiet. Transport the patient immediately to the nearest physician.

Atropine is the emergency antidote. Consult your physician about obtaining an adequate supply of 1-10-grain atropine tablets for use in case of poisoning. 2-PAM is also antidotal and may be administered in conjunction with atropine. CALL A PHYSICIAN IMMEDIATELY IN ALL CASES OF SUSPECTED POISONING.

INTERNAL: Induce vomiting immediately. This may be done by introducing a finger into the throat or by giving warm salt water (1 tablespoon of salt to a glass of water). NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON. Keep patient prone and quiet. Start artificial respiration immediately if patient is not breathing.

EXTERNAL: If the material has been spilled on the skin immediately remove patient from the vicinity of the insecticide, remove all contaminated clothing and wash skin with soap and running water. If the material gets into the eyes, wash immediately with running water for at least 10 minutes.

NOTE TO PHYSICIAN: Active ingredients are cholinesterase inhibitors. Treat symptomatically. Atropine is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine. Morphine is contraindicated. Clear chest by postural drainage. Observe patient continuously for 48 hours. Repeated exposures to cholinesterase inhibitors may without warning cause prolonged susceptibility to very small doses of any cholinesterase inhibitor. Allow no further exposure until time for cholinesterase regeneration has been allowed as determined by blood test.

CONTAINER DISPOSAL

DO NOT re-use empty container. Decontaminate by repeated rinsing with water and detergent. Crush or perforate and bury in a safe place.

NOTICE: Seller's guaranty shall be limited to the terms of the label and subject thereto. The buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.