



RESIDUAL ROACH SPRAY

contains Dursban*

RESIDUAL SPRAY FOR INDUSTRIAL AND INSTITUTIONAL USE
KILLS RESISTANT ROACHES

A highly effective ready-to-use formulation for the control of Cockroaches, including those resistant to certain chlorinated hydrocarbons. This product drives insects from their hiding places, gives rapid kill and long residual activity.

FOR USE BY PEST CONTROL OPERATORS ONLY

ACTIVE INGREDIENTS:

O,O-Diethyl O-(3,5,6-trichloro-2-pyridyl) phosphorothioate	0.25%
Pyrethrins	0.05%
Technical Piperonyl Butoxide**	0.25%
INERT INGREDIENTS	99.45%
TOTAL	100.00%

**Equivalent to 0.20% (butylcarbityl (6-propyl-piperonyl) ether and 0.05% related compounds

DURSBAN - T. M. of The Dow Chemical Company, U. S. Pat No. 3,244,586

CAUTION: KEEP OUT OF REACH OF CHILDREN

See side panels for additional cautions and information

U.S.D.A. Registration Number 6720-158



MANUFACTURED BY
Southern MILL CREEK PRODUCTS COMPANY, Inc.

TAMPA, FLORIDA

ACCEPTED
3-11-71

RESTRICTIONS

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC PEST REGISTRATION
ED UNDER NO. 6720-128

FOR SPOT APPLICATION ONLY

DO NOT use as a space spray. Provide adequate ventilation of area being treated. Avoid breathing of spray. DO NOT allow children to contact treated area until surfaces are thoroughly dry. DO NOT store near feed or food products. Avoid contamination of feed, food, food processing equipment, and food handling surfaces. DO NOT apply to animals. Remove pets and cover fish bowls before spraying. DO NOT apply to plants.

CAUTION

May be harmful if swallowed, inhaled or absorbed through skin. DO NOT get in eyes, on skin or clothing. In case of contact wash immediately with soap and water. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and water before reuse. DO NOT allow children to contact treated surfaces until completely dry.

NOTE TO PHYSICIAN: Active ingredient is cholinesterase inhibitor. Treat symptomatically. Atropine is an antidote.

NOTICE: Seller's guaranty shall be limited to the terms of the label and subject thereto. The buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

DIRECTIONS FOR USE

Apply with a low pressure sprayer equipped with a coarse spray nozzle. Spray surfaces only until wet. Avoid excessive run-off. Apply to cockroach breeding sites and harborages. Also, treat adjacent exposed surfaces where cockroaches may crawl when they come out of hiding. Repeat treatment as

May be harmful if swallowed, inhaled or absorbed through skin. DO NOT get in eyes, on skin or clothing. In case of contact wash immediately with soap and water. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and water before reuse. DO NOT allow children to contact treated surfaces until completely dry.

NOTE TO PHYSICIAN: Active ingredient is cholinesterase inhibitor. Treat symptomatically. Atropine is an antidote.

NOTICE: Seller's guaranty shall be limited to the terms of the label and subject thereto. The buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

DIRECTIONS FOR USE

Apply with a low pressure sprayer equipped with a coarse spray nozzle. Spray surfaces only until wet. Avoid excessive run-off. Apply to cockroach breeding sites and harborages. Also, treat adjacent exposed surfaces where cockroaches may crawl when they come out of hiding. Repeat treatment as necessary.

Cockroaches are normally found in dark corners of rooms and closets, at base of walls, in cracks in the wall, behind baseboards, behind and beneath sinks, refrigerators, stoves and cabinets, around drain pipes, in and around water containers, floor drains and switch panels.

Dishes and food handling utensils should be thoroughly washed in soap and water if they become contaminated by the application of this product. Repeat treatment as needed.

NOTE: A period of 4 to 7 days after treatment is normally required for maximum effect on cockroaches.

Meat Packing and Food Processing Plants: Use only in inedible product areas of meat packing plants, including those which are federally inspected. Any exposed food products, equipment and packaging materials must be removed or covered before spraying. Rinse all equipment with water before resuming operation.

Treat food processing plants early in the season before canning or other processing begins. Repeat application, if necessary, when plant is not in operation. Spraying unloading docks and outside areas where insects congregate will aid in preventing reinfestation. Good sanitation practices and proper disposal of wastes are essential in effective insect control.