

NET  
WEIGHT

5%

ACTIVE INGREDIENT  
Malathion (0,0 dimethyl dithio phosphoric acid diethyl mercaptosuccinate)  
INERT INGREDIENTS

5  
95  
100

#### DIRECTIONS FOR USE

Do not use in edible products areas of food processing plants, restaurants or other areas where food is commercially prepared or processed. Do not use in serving areas while food is exposed.

#### PEST CONTROL OPERATORS

For the control of Ants, Crickets, Clover Mites, Earwigs, Pantry Pests, Saw-toothed Grain Beetle, Flour Beetle, Rice Weevil, Cigarette Beetle, Drugstore Beetle and Indian Meal Moth, Scorpions, Silverfish and Spiders in Homes, Dairies and Food Processing Plants apply Malathion 5% Dust liberally around window frames, cracks and crevices, around doors, baseboards, storage areas, bookcases, under sinks and to other areas frequented by these pests. It will also control Fleas, Brown Dog Ticks and Roaches.

Malathion is not  
a for Copy

Do not treat kittens or puppies under 4 weeks of age.

## PEST CONTROL OPERATORS

For the control of Ants, Crickets, Clover Mites, Earwigs, Pantry Pests, Saw-toothed Grain Beetle, Flour Beetle, Rice Weevil, Cigarette Beetle, Drugstore Beetle and Indian Meal Moth, Scorpions, Silverfish and Spiders in Homes, Dairies and Food Processing Plants apply Malathion 5% Dust liberally around window frames, cracks and crevices, around doors, baseboards, storage areas, bookcases, under sinks and to other areas frequented by these pests. It will also control Fleas, Brown Dog Ticks and Roaches.

Do not treat kittens or puppies under 4 weeks of age.

Dust Bath Boxes: For the control of Poultry Lice apply 1 lb. of dust per box (18"x12"x3") per 30 hens. Remove boxes when dust has been used. For the control of Sticktight Fleas on birds place 12 lbs. of dust in box (2"x3'x1') per 100 birds, or apply in natural soil wallows at the rate of 6 lbs. of dust per 50 bird. In range pens apply 3/4 lb. per 20 sq.ft. of surface. Repeat in 28 days and then as often as necessary. In brooder houses distribute dust evenly in each corner of the brooder house at the rate of 1 1/2 lbs. per 100 chicks.