



ST. AUGUSTINE GRASS
BROADLEAF HERBICIDE
 Contains TRIMEC (R) Herbicide

ACTIVE INGREDIENTS:

| | |
|---|--------|
| *Dimethylamine salt of 2,4-dichlorophenoxyacetic acid | 3.23% |
| **Dimethylamine salt of 2-(2-methyl-4-chlorophenoxy) propionic acid | 10.59% |
| ***Dimethylamine salt of Dicamba (3,6-dichloro-o-anisic acid) | 1.28% |

INERT INGREDIENTS

TOTAL 100.00%

*2,4-dichlorophenoxyacetic acid equivalent 0.235 pounds per gallon or 2.69%
 **2-(2-methyl-4-chlorophenoxy) propionic acid equivalent 0.764 pounds per gallon or 8.76%
 ***3,6-dichloro-o-anisic acid equivalent 0.094 pounds per gallon or 1.07%

CAUTION: KEEP OUT OF REACH OF CHILDREN

See side panels for additional cautions and information

U. S. Patent No. 3,284,186

EPA Reg. No. 6720-178

WEEDS CONTROLLED

St. Augustine Grass Broadleaf Herbicide is especially effective against weeds as SHEEP SORREL, KNOT WEED, SPOTTED SPURGE, CHICKWEED, OXALIS AND HENBIT. In addition to these, it kills virtually all broadleaf weeds, some of which are listed below.

| | | |
|--------------|---------------|------------------|
| Bedstraw | Heal-all | Ragweed |
| Black Medic | Henbit | Sheep Sorrel |
| Buckhorn | Knotweed | Shepherd's Purse |
| Bull Thistle | Lambsquarter | Speedwell |
| Burdock | Lespedeza | Tansy Mustard |
| Buttercup | Mallow | Thistle |
| Chicory | Morning Glory | Spurge |
| Chickweed | Mustard | Velvet Leaf |
| Clover | Oxalis | Wild Carrot |
| Cockleburr | Peppergrass | Wild Garlic |
| Dandelion | Pigweed | Wild Onion |
| Dock | Plantains | Wild Lettuce |
| Galinsoga | Poison Ivy | Yarrow |
| Ground Ivy | Poison Oak | And many other |
| Hawkweed | Purslane | Broadleaf Weeds |

CAUTION

Harmful if swallowed. Avoid contact with skin, eyes or clothing.

Avoid breathing spray mist. If contact with eyes occurs, flush with plenty of water.

Avoid drift of spray mist onto vegetables, flowers, or namental plants, shrubs, trees and other desirable plants. Do Not contaminate domestic or irrigation water.

Do not store in contact with fertilizers, seeds, insecticides, or fungicides. Use only lawn sprayers for this use. Coarse sprays are less likely to drift.

Do not apply to areas underlaid by roots of desirable trees and shrubs.

When container is empty, wash thoroughly and destroy.

Never re-use. Destroy by perforating or crushing, and burying in a safe place. Do not burn.

NOTICE: Seller's guaranty shall be limited to the terms of the label and subject thereto. The buyer assumes any risk of damage or property arising out of use of handling, and accepts the product on these conditions.



MANUFACTURED BY

Southern MILL CREEK PRODUCTS COMPANY, Inc.

TAMPA, FLORIDA

**DIRECTIONS
FOR SPOT TREATMENT ONLY**

For general vegetation control in sand traps only. Will not damage adjacent Bermuda turf when used as directed. Apply to moist sand. Do not water for 24 hours after application.

To Kill and Prevent Growth of most Vegetation

Apply at the rate of 1 quart of TRIM-TRAP in 5 gallons of water to each 1000 sq. ft. For smaller areas, mix 4 oz. in 1 gallon of water and apply to 100-150 sq. ft.

**DO NOT USE OR STORE IN OR AROUND THE HOME.
DANGER IN USING**

Sodium Arsenite is poisonous. Therefore, precautions must be observed when handling (See antidote and warnings on side panel.)

Store the solution in a safe place. Children must not be able to reach or handle the solution. It is recommended that only the amount of diluted solution be prepared from the concentrate that will be used immediately.

Mix the solution into water outdoors or in a well ventilated room. Wash off the outside of the bottle of solution in running water before storing away.

Avoid spray drift to desirable plants.

Animals must not be able to reach the solution. This product is toxic to fish and wildlife. Birds and other wildlife in treated areas may be killed. DO NOT apply where runoff is likely to occur. DO NOT contaminate water by cleaning of equipment, or disposal of wastes.

DO NOT contaminate water used for irrigation or domestic purposes.

DO NOT contaminate feed, water or foodstuffs.

DO NOT apply to areas underlaid by roots of desirable trees and shrubs.



NET
CONTENTS:

TRIM-TRAP

HERBICIDE

A NON-SELECTIVE HERBICIDE FOR THE CONTROL OF WEEDS IN
GOLF COURSE SAND TRAPS



POISON



ACTIVE INGREDIENTS:

Sodium Arsenite

41.7%

INERT INGREDIENTS

58.3%

TOTAL

100.0%

Arsenic expressed as metallic all in water soluble form: 22.1
Trioxide As₂O₃ per gallon 3.5 pounds Arsenic

DANGER: KEEP OUT OF REACH OF CHILDREN

SEE SIDE PANELS FOR ANTIDOTE STATEMENT, WARNINGS AND DIRECTIONS

E.P.A. Reg. No. 6720-181

E.P.A. Est. 6720-FL-1

MANUFACTURED BY
 Southern MILL CREEK PRODUCTS COMPANY, Inc.
TAMPA, FLORIDA

KILLS Dandelion, Chickweed, Knotweed, Plantain,
Henbit, Spurge, Crabgrass, and other broadleaved weeds

HOW TO MIX

1. For use on St. Augustine grass, bentgrass, buffalo grass and grass seedlings, mix 1 ounce in 1 gallon of water and uniformly spray 400 sq. ft. (for example, an area 10 feet wide and 40 feet long). If necessary, repeat treatment in 2 to 3 weeks if hard to kill weeds persist.
2. For use on certain other grasses such as Bermuda, Bahia, bluegrass, centipede, fescue and rice. Mix 1 1/2 ounce in 1 gallon of water and spray uniformly over 400 sq. ft. On hard to kill species or under poor growing conditions or on older weeds, mix 2 ounces in 1 gallon of water and spray on 400 sq. ft. with special attention to thorough coverage of the hard to kill weeds.

DIRECTIONS FOR USE

GENERAL INFORMATION St. Augustine Grass Broadleaf Herbicide is specially formulated for maximum safety to seedling grasses and 2,4-D sensitive grasses such as St. Augustine grass, bentgrass and buffalo grass. Sprays should be applied when broadleaf weeds are small and tender for best results. If weeds are older and hard to kill, the higher rates recommended should be used. Avoid spraying St. Augustine grass under conditions of high temperature and moisture when the St. Augustine is growing rapidly. Under these conditions or in cases of accidental overdose, some slight temporary discoloration of the St. Augustine foliage might occur.

To insure weeds are growing and able to absorb the herbicide, irrigate the lawn 2 or 3 times prior to spraying. Do not water for 24 hours after treatment. Although spray works faster with temperatures above 70 F, spray can be applied at any time plants are growing. Spray when it is calm to avoid drift. Spray will injure or kill clover, dandelion and most ornamental and vegetable crops.

Amount of water in spray is not important, except to evenly distribute the chemical. If directions call for 1 ounce in 1 gallon on 400 sq. ft. it may be more convenient to use 1 oz. in 2 gallon on 400 sq. ft. depending on the sprayer. Generally large volume of water result in more uniform coverage.

ST. AUGUSTINE GRASS BROADLEAF HERBICIDE

ACTIVE INGREDIENTS

- *Dimethylamine salt of 2,4-dichlorophenoxyacetic acid
- **Dimethylamine salt of 2-methyl-4-chlorophenoxyacetic acid
- ***Dimethylamine salt of 3,6-dichloro-2-naphthoic acid

INERT INGREDIENTS

- *2,4-dichlorophenoxyacetic acid
- **2-methyl-4-chlorophenoxyacetic acid
- ***3,6-dichloro-2-naphthoic acid

CAUTION: KEEP OUT OF REACH OF CHILDREN
See side panels for first aid instructions.
U.S. Patent No. 3,284,186



ACCEPTED
4/8/72
6720-178

6720-178



ST. AUGUSTINE GRASS
BROADLEAF HERBICIDE
 Contains TRIMEC (R) Herbicide

ACTIVE INGREDIENTS
 *Diquat dibromide (4-aminodipyridine) 25%
 **Diquat dibromide (4-aminodipyridine) 25%
 ***Diquat dibromide (4-aminodipyridine) 25%

INERT INGREDIENTS
 75%

1.4 lb. per gallon (4.54 kg. per liter) equivalent to 25% Diquat
 2.12 lb. per gallon (6.25 kg. per liter) equivalent to 25% Diquat
 0.70 lb. per gallon (2.00 kg. per liter) equivalent to 25% Diquat
 ***3.12 lb. per gallon (8.90 kg. per liter) equivalent to 25% Diquat

CAUTION: KEEP OUT OF REACH OF CHILDREN

See directions for additional cautions and information

U.S. Patent No. 284181

EPA Reg. No. 6720-178

MANUFACTURED BY

MILL CREEK PRODUCTS COMPANY, Inc.
 TAMPA, FLORIDA

WEEDS CONTROLLED

Abutilon
 Aeschynomene
 Amaranthus
 Aster
 Bidens
 Cirsium
 Conium
 Echinochloa
 Eragrostis
 Galium
 Geranium
 Ipomoea
 Lactuca
 Malva
 Mimulus
 Mollis
 Nymphaea
 Paspalum
 Portulaca
 Rotala
 Sida
 Solanum
 Stachys
 Taraxacum
 Verbena
 Xanthoxylum
 Zinnia

Abutilon
 Aeschynomene
 Amaranthus
 Aster
 Bidens
 Cirsium
 Conium
 Echinochloa
 Eragrostis
 Galium
 Geranium
 Ipomoea
 Lactuca
 Malva
 Mimulus
 Mollis
 Nymphaea
 Paspalum
 Portulaca
 Rotala
 Sida
 Solanum
 Stachys
 Taraxacum
 Verbena
 Xanthoxylum
 Zinnia

NOTICE
 See the label
 for full
 directions

2-178

KILLS. Dandelions, Chickweed, Knotweed, Plantains, Herb's Spruce and many other broadleaf weeds.

HOW TO MIX

1. For use on St. Augustine grass, bentgrass, buffalo grass and grass seedlings, mix 1 ounce in 1 gallon of water and uniformly spray 400 sq. ft. for example, an area 10 feet wide and 40 feet long. If necessary, repeat treatment in 2 to 3 weeks, if hard to kill weeds persist.
2. For use on certain other grasses such as Bermuda, Bahia, bluegrass, centipede, fescue and rye, mix 1 ounce in 1 gallon of water and spray uniformly over 400 sq. ft. On hard to kill species or under poor growing conditions or on older weeds, mix 2 ounces in 1 gallon of water and spray on 400 sq. ft. with special attention to thorough coverage of the hard to kill weeds.

DIRECTIONS FOR USE

GENERAL INFORMATION. St. Augustine Grass Broadleaf Herbicide is specially formulated for maximum safety to seedling grasses and 2,4-D sensitive grasses such as St. Augustine grass, bentgrass and buffalo grass. Sprays should be applied when broadleaf weeds are small and tender for best results. If weeds are older and hard to kill, the higher rates recommended should be used. Avoid spraying St. Augustine grass under conditions of high temperature and moisture when the St. Augustine is growing rapidly. Under these conditions, or in cases of accidental overdose, some slight temporary discoloration of the St. Augustine foliage might occur.

To insure weeds are growing and able to absorb the herbicide, irrigate the lawn 2 or 3 times prior to spraying. Do not water for 24 hours after treatment. Although spray works faster with temperatures above 70°F, sprays can be applied at any time plants are growing. Spray when air is calm to avoid drift. Spray will injure or kill clover, dandelion and most ornamental and vegetable crops.

Amount of water in spray is not important, except to evenly distribute the chemical. If directions call for 1 ounce in 1 gallon on 400 sq. ft. it may be more convenient to use 1 oz. in 2 gallons on 400 sq. ft. depending on the sprayer. Generally large volume of water result in more uniform coverage.



ST. AUGUSTINE BROADLEAF HERBICIDE Contains TRIMEC (R)

| ACTIVE INGREDIENTS: | TOTAL |
|---|---|
| *Dimethylamine salt of 2,4-dichlorophenoxyacetic acid | *2,4-dichlorophenoxyacetic acid equivalent 0.094 lb/gal |
| **Dimethylamine salt of 2-(2-methyl-4-chlorophenoxy) propionic acid | **2-(2-methyl-4-chlorophenoxy) propionic acid equivalent 0.764 pounds per gallon or 8.76% |
| ***Dimethylamine salt of Dicamba (3,6-dichloro-o-anisic acid) | ***3,6-dichloro-o-anisic acid equivalent 0.094 lb/gal |
| INERT INGREDIENTS | |

CAUTION: KEEP OUT OF REACH OF CHILDREN
See side panels for additional cautions
U. S. Patent No. 3,284,186

MANUFACTURED BY
Southern MILL CREEK PRODUCTS
TAMPA, FLORIDA

ACCEPTED
9/5/72
FOR LOCAL USE
ED IN ACCORDANCE WITH
TO APPLICABLE REGULATIONS

6720-178

26



ST. AUGUSTINE GRASS BROADLEAF HERBICIDE Contains TRIMEC (R) Herbicide

ACTIVE INGREDIENTS

| | |
|---|----------------|
| *Dimethylamine salt of 2,4-dichlorophenoxyacetic acid | 3.23% |
| **Dimethylamine salt of 2-(2-methyl-4-chlorophenoxy) propionic acid | 10.59% |
| ***Dimethylamine salt of Dicamba (3,6-dichloro-o-anisic acid) | 1.28% |
| INERT INGREDIENTS | 84.90% |
| TOTAL | 100.00% |

*2,4-dichlorophenoxyacetic acid equivalent 0.235 pounds per gallon or 2.69%
 **2-(2-methyl-4-chlorophenoxy) propionic acid equivalent 0.764 pounds per gallon or 8.76%
 ***3,6-dichloro-o-anisic acid equivalent 0.094 pounds per gallon or 1.07%

CAUTION: KEEP OUT OF REACH OF CHILDREN

See side panels for additional cautions and information

U.S. Patent No. 3,284,186

EPA Reg. No. 6720-178

WEEDS CONTROLLED

St. Augustine Grass Broadleaf Herbicide is especially effective against weeds as SHEEP SORREL, KNOT WEED, SPOTTED SPURGE, CHICKWEED, OXALIS AND HENBIT. In addition to these, it kills virtually all broadleaf weeds, some of which are listed below.

| | | |
|--------------|---------------|------------------|
| Bedstraw | Heal-all | Ragweed |
| Black Medic | Henbit | Sheep Sorrel |
| Buckhorn | Knockweed | Shepherd's Purse |
| Bull Thistle | Lambsquarter | Speedwell |
| Burdock | Lespedeza | Tansy Mustard |
| Buttercup | Mallow | Thistle |
| Chicory | Morning Glory | Spurge |
| Chickweed | Mustard | Velvet Leaf |
| Clover | Oxalis | Wild Carrot |
| Cocklebur | Peppergrass | Wild Garlic |
| Dandelion | Pigweed | Wild Onion |
| Dock | Plantains | Wild Lettuce |
| Galinsoga | Poison Ivy | Yarrow |
| Ground Ivy | Poison Oak | And many other |
| Hawkweed | Purslane | Broadleaf Weeds |

Harmful or clo...

Avoid bre...

Avoid d...

Do not...

Do not...

When co...

Never re...

NOTICE: ...

MANUFACTURED BY
Southern MILL CREEK PRODUCTS COMPANY, Inc.
TAMPA, FLORIDA

0-178



ST. AUGUSTINE GRASS

BROADLEAF HERBICIDE with TRIMEC (R) Herbicide

| | |
|--|----------------|
| 2,4-dichlorophenoxyacetic acid | 3.23% |
| 2-(2-methyl-4-chlorophenoxy)propionic acid | 10.59% |
| Dicamba (3,6-dichloro-o) | 1.28% |
| | 84.90% |
| TOTAL | 100.00% |
| acid equivalent 0.235 pounds per gallon or 2.69 pounds per 100 gallons | |
| propionic acid equivalent 0.094 pounds per gallon or 1.07 pounds per 100 gallons | |

OUT OF REACH OF CHILDREN
For additional cautions and information
EPA Reg. No. 6720-178

WEEDS CONTROLLED

St. Augustine Grass Broadleaf Herbicide is especially effective against weeds as SHEEP SORREL, KNOT WEED, SPOTTED SPURGE, CHICKWEED, OXALIS AND HENBIT. In addition to these, it kills virtually all broadleaf weeds, some of which are listed below.

| | | |
|--------------|---------------|------------------|
| Bedstraw | Heal-all | Ragweed |
| Black Medic | Henbit | Sheep Sorrel |
| Buckhorn | Knotweed | Shepherd's Purse |
| Bull Thistle | Lambsquarter | Speedwell |
| Burdock | Lespedeza | Tansy Mustard |
| Buttercup | Mallow | Thistle |
| Chicory | Morning Glory | Spurge |
| Chickweed | Mustard | Velvet Leaf |
| Clover | Oxalis | Wild Carrot |
| Cockleburr | Peppergrass | Wild Garlic |
| Dandelion | Pigweed | Wild Onion |
| Dock | Plantains | Wild Lettuce |
| Galinsoga | Poison Ivy | Yarrow |
| Ground Ivy | Poison Oak | And many other |
| | Purslane | Broadleaf Weeds |

CAUTION

Harmful if swallowed. Avoid contact with skin, eyes or clothing.

Avoid breathing spray mist. If contact with eyes occurs, flush with plenty of water.

Avoid drift of spray mist onto vegetables, flowers, or ornamental plants, shrubs, trees and other desirable plants. Do Not contaminate domestic or irrigation water.

Do not store in contact with fertilizers, seeds, insecticides, or fungicides. Use only lawn sprayers for this use. Coarse sprays are less likely to drift.

Do not apply to areas underlaid by roots of desirable trees and shrubs.

When container is empty, wash thoroughly and destroy. Never re-use. Destroy by perforating or crushing, and burying in a safe place. Do not burn.

NOTICE: Seller's guaranty shall be limited to the terms of the label and subject thereto. The buyer assumes any risk of damage or property arising out of use of handling, and accepts the product on these conditions.

MANUFACTURED BY
MILL CREEK PRODUCTS COMPANY, Inc.
TAMPA, FLORIDA