

[6720-185]

DIRECTIONS FOR USE

For Flies, Mosquitoes, and Gnats: Close windows and doors. Spray freely toward ceiling in all directions until room is filled with mist, then leave the room. Keep doors closed for at least 15 minutes. Ventilate the room when treatment is completed.

Wasps: Application should be made in the late evening when insects are at rest. Spray liberally into hiding and breeding places, contacting as many insects as possible. Repeat as necessary.

NET CONTENT



ACCEPT

MAR 8 1977

UNDER THE FEDERAL INSECTICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REG. NO. 6720-185. SEE ATTACHED COMMENTS.

SBP-1382 INSECTICIDE SPRAY 0.10%

FOR: Bakeries, Food Plants, Restaurants, Schools, Hotels, Motels, Office Buildings, Hospitals and Food Handling Plants.

ACTIVE INGREDIENTS:

*+ (5-Benzyl-3-furyl) methyl 2,2-dimethyl-3-(2-methylpropenyl) cyclopropanecarboxylate	0.100%
Related compounds	0.014%
Aromatic petroleum hydrocarbons	0.132%
Petroleum distillate	99.750%
INERT INGREDIENTS:	0.004%
TOTAL	100.000%

*Cis-trans isomers ratio: 30% (·) cis and min. 70% (+) trans.

U.S. Patent No. 3,465,007


CAUTION: KEEP OUT OF REACH OF CHILDREN

E. P. A. Registration Number 6720-185



MANUFACTURED BY

Southern MILL CREEK PRODUCTS COMPANY, Inc.
TAMPA, FLORIDA



SBP-1382 INSECTICIDE SPRAY 0.10%

FOR: Bakeries, Food Plants, Restaurants, Schools, Hotels, Motels, Office Buildings,
Hospitals and Food Handling Plants.

ACTIVE INGREDIENTS:

*+(5-Benzyl-3-furyl) methyl 2,2-dimethyl-3-(2-methylpropenyl) cyclopropanecarboxylate	0.100%
Related compounds	0.014%
Aromatic petroleum hydrocarbons	0.132%
Petroleum distillate	99.750%
INERT INGREDIENTS:	0.004%
TOTAL	100.000%

*Cis trans isomers ratio: 30% (+) cis and min. 70% (+) trans.

U. S. Patent No. 3,465,007

CAUTION: KEEP OUT OF REACH OF CHILDREN

E. P. A. Registration Number 6720-185



Southern MILL CREEK PRODUCTS COMPANY, Inc.
TAMPA, FLORIDA

MANUFACTURED BY

CAUTION
KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Cover or remove fishbowls. Do not store near heat of open flame.

Avoid contamination of feed and foodstuffs. Use in food processing plants only when the plant is not in operation. Foods should be removed or covered during treatment. All food processing surfaces should be covered or thoroughly cleaned before using. In hospitals, patients should be removed from room prior to treatment. Room should be ventilated for 2 hours after spraying. Do not return patients to room until after ventilation.

Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place away from water supplies.

(Drum disposal instructions for 30 & 55 gallon drums)

Do not reuse empty drum. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place away from water supplies.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME