


6720-193

03/20/1974

1/4



NET CONTENTS

CONTENIDO NETO

DIAZINON* AG 500

INSECTICIDE: For control of insects on fruits, vegetables, field crops and ornamentals.

INSECTICIDA: Para el control de los insectos en frutas, vegetales, campos de cosecha y ornamentales.

WARNING: Keep out of reach of children. See additional warning statements on back/side of container.

AVISO: Manténgase fuera del alcance de los niños. Ver avisos adicionales en los paneles del lado de la etiqueta.

ACTIVE INGREDIENTS:

0,0-diethyl 0-(2-isopropyl-4-methyl-6-pyrimidinyl).....	48%
Xylene.....	38%
INERT INGREDIENTS.....	14%
TOTAL.....	100%

ACCEPTED

MAR 20 1974

UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTERED UNDER NO. 6720-193 SUBJECT TO ANTIDOTE COMMENTS.

DIAZINON AG 500 contains 4 lbs. Diazinon per gallon.

*DIAZINON - Reg. Trademark of CIBA-GEIGY Corporation

E.P.A. Reg. No. 6720-193



MANUFACTURED BY

SOUTHERN MILL CREEK PRODUCTS COMPANY, INC.

TAMPA, FLORIDA

May be fatal if swallowed. May be absorbed through skin. Do not breath spray mist. Do not get in eyes, on skin, or on clothing. Wash thoroughly after handling. Avoid contamination of food. Do not use on humans or permit children to go on to sprayed grass until spray has completely dried. Fish and Wildlife Caution: This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not apply when weather conditions favor drift from areas treated or where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service. Do not reuse empty container. Destroy by perforating or crushing and burying in a safe place.


CAUTION: INFLAMMABLE MIXTURE. DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME. Minimum Flash Point 82°F.

General Information:

Diazinon AG500 is an emulsifiable solution which when diluted with water according to the directions below, is intended for the control of insect pests of agricultural crops. Do not use in barns or dwellings. Do not use on livestock.

In the directions for use given below, best control is obtained when application is made at first sign of insects. Application should be repeated as necessary to maintain control.

NOTE: The marketing of raw agricultural commodities having residues in excess of their permitted tolerances, or marketing those for which no tolerances have been set and bearing residues, will violate Federal Law when shipped in interstate commerce and may violate State Law.



USE OF DIAZINON AG-500

Vegetables

Crops	Insects	Dosage of Diazinon AG 500 per Acre	Time to wait after last application
Beans	leaf miners	1 - 1½ Pts.	7 days
	:	:	Bean forage or bean hay may
	:	:	be fed to dairy and beef cat-
	:	:	tle and sheep. Wait 4 days
Cabbage	aphids	½ - 1 Pt.	7 - 10 days
	diamond-back moths (Plutella)	:	:
	:	:	:
	:	:	:
Cucumber	Serpentine leaf miner	1 Pt.	7 days
	thrips	:	:
Squash	aphids	1 - 1½ Pt.	7 days
	leafhopper	:	:
	mites	:	:
	Serpentine leaf miners	:	:
Melons	aphids	1 - 1½ Pt.	7 days
	leafhoppers	:	:
	mites	:	:
Peppers	aphids	:	:
	Serpentine leaf miners	½ Pt.	5 days
Tomatoes	aphids	½ Pt.	1 day
	dipterous leaf miners	:	:
	vinegar flies	½ - 1½ Pt	:
Lettuce	aphids	:	:
	dipterous leaf miners	½ - 1 Pt.	10 days

Fruits

Crops	Insects	Dosage of Diazinon AG500 per acre	Time to wait after last application
Citrus	aphids	½ - 1 Pt. per	Do not apply during bloom
	:	100 gls. of water	Do not apply to citrus within
	thrips	1 pt. per 100 gls.	21 days of harvest
	:	of water	Shrimps and crabs may be
Pineapples	citrus snow scale	1 - 2 Pt. per	killed at application rates
	:	100 gls. water	recommended on this label.
	:	:	Do not apply when these are
	:	:	important resources.
Pineapples	scale ?	5 Qts. plus 2 gls.	
	:	of light or medium	
	:	horticultural oil	
	:	in 500 gls. of wa	
Pineapples	:	ter per acre to	
	:	the vegetative	
	:	slips of the plants	
	:	following harvest	
Pineapples	mealybugs	2 Qts.	May be applied on day of
	:	:	harvest



Sugar Cane

Crops	Insects	Dosage of Diazinon AG500 per Acre	Time to wait after last application
Sugar Cane	'yellow sugarcane aphid	'½ - 1 Pt.	'Allow 1 day between last application and harvest

Lawn Insect Control			
	Insects	Rate per 1000 square feet	Amount of water to use
Turf	'Lawn Chinch bug	'2 - 3 oz.	' 25 gls.
	'ants	'Spot spray ant hills	'3 gls.
	'crickets	'	'
	'cutworms	'4 oz.	'
	'armyworm	'	'
	'earwigs	'	'
	'Brown dog ticks	'1½ oz. Do not apply to animals. For brown dog ticks spray grass and under shrubbery, particularly near house.	' 3 gls.
	'Bermuda Mites	'	'
	'Fleas	'	'
	'Ticks	'	'

Ornamentals

To control certain insects on Azalea, Camellia, Carnation, Chrysanthemum, Elm Gladioli, Lilac, Maple, Oak, Pine and Spruce apply the recommended rates indicated below.

Insects	Rate per 3 gls. of water	Rate per 100 gls of water
'aphids	'	'
'dipterous leaf miners	'½ oz.	' 1 Pt. *
'thrips	'	'
'scale crawlers	'	'
'leafhopper	'	'
'white flies	'	'
'flea beetles	'	'

NOTE: Do not use Diazinon AG 500 on certain ferns, poinsettia, hibiscus, papaya, pilea and gardenia, since injury to the plant may occur.

(*) 1 Fl oz. = 2 tablespoons
 16 Fl. oz. = 1 Pint

