



MSMA H.C. 8
Liquid High Concentrate
 For Selective Post-Emergent Weed Control

| | | |
|--|--------|---------|
| ACTIVE INGREDIENT | | |
| Monosodium Acid Methanearsonate | 58.39% | |
| INERT INGREDIENTS | 41.61% | |
| | | TOTAL |
| | | 100.00% |
| Total Arsenic as elemental ar | | |
| in water solution | 27.91% | |
| *Product contains 5.0 lbs. MSMA per gallon | | |

Nov 7, 1974

6720-219

CAUTION: KEEP OUT OF REACH OF CHILDREN

See directions and other cautions on side panels.

EPA Reg. No. 6720-219

EPA Est. No. 6720-FL-1

NET CONTENTS



DISTRIBUTED BY

Southern MILL CREEK PRODUCTS COMPANY, Inc.
 TAMPA, FLORIDA

DIRECTIONS FOR USE:

COTTON: MSMA H.C. 8 Herbicide is useful as a directed application in cotton for the post-emergent control of weeds and grasses such as johnsongrass, nutsedge, dallisgrass, cocklebur, ragweed, sandbur and puncture vine. It should be applied at the rate of 1 quart per acre.

Add 5 pints of MSMA H.C. 8 Herbicide and 1 to 2 quarts of a suitable surfactant to 100 gallons of water and mix thoroughly. Apply spray solution at 40 gallons per acre. Weed control is most effective at temperatures above 70 F. For band application in 40 inch rows, apply 1 gallon of spray solution per inch of band width per acre. Direct the spray solution to the base of the cotton plant. Nozzles should be placed so as to avoid spraying the cotton foliage. A second application may be required during humid growing conditions to control perennials such as johnsongrass and nutsedge. Apply the second application 1 to 3 weeks later.

Slight burning and a reddish discoloration of the cotton leaf may occasionally occur following the recommended treatment, but the cotton plant will develop normally.

Apply only when cotton is in bloom. Do not apply after first bloom. Do not feed treated foliage to livestock, or graze treated areas.

NON-CROP: MSMA H.C. 8 Herbicide is useful for controlling johnsongrass, nutsedge, and certain other weeds on drainage ditches, banks, rights-of-way, storage yards, and similar non-crop areas.

Mix MSMA H.C. 8 Herbicide at a rate of 2 to 5 pints, plus 1 to 2 quarts of a suitable surfactant in 100 gallons of water. Spray the unwanted vegetation thoroughly to just short of run-off. Any spray equipment that gives good coverage may be used. If regrowth occurs, reapply as required.

LAWN AND ORNAMENTAL TURF: MSMA H.C. 8 Herbicide is useful for control of dallisgrass, sandbur, bahiagrass, nutsedge, crabgrass, and kikweed, and broad scurf with little or no injury to good lawn grasses. On new lawns, do not treat until after 3 to 4 mowings. Good grasses may be temporarily discolored. Zoysia, bluegrass, and bermuda are quite tolerant. Do not use on St. Augustine or centipede.

Mow lawns 1 to 1 1/2 inches high before treatment. Mix 1 fluid ounce (2 tablespoons) of MSMA H.C. 8 Herbicide and 1 fluid ounce (1 tablespoon) of a suitable surfactant in 5 gallons of water and apply to an area of 1,000 square feet. Spray thoroughly to wet all undesirable plants. Repeat applications 10 to 14 days apart, may be needed for good control.

9-25-74

CAUTION: KEEP OUT OF REACH OF CHILDREN
 Wash hands after using. All material has been washed.

ANTIDOTE: If taken internally, drink milk or water.

READ ENTIRE LABEL BEFORE USING.

NOTICE: Use of this material may result in excessive residues, or other effects.

GENERAL INFORMATION: This material controls growing weeds at air temperatures above 70 F.

MIXING INSTRUCTIONS: Mix with water and add the required amount of surfactant. After use, clean equipment.

Do not contaminate watersheds. Do not contaminate water by disposal.

CONTAINER DISPOSAL: Dispose of in a safe place.

NOTICE: Seller's warranty does not extend to use or handling not recommended on label.