DIRECTIONS FOR USE

A non-selective herbicide that can be applied to post-emergence world control it is recommended that Weedox C. R 250 be used in the areas such as industrial sites, around buildings, roadways, right-ofways, lumber yards, tank farms, cemeteries, driveways, parking lots, fence areas, recreational areas and airports. Weedox C/ R 250 is water emulsifiable and non-corrosive to equipment.

DILUTION RECOMMENDATIONS FOR WEED AND GRASS CONTROL: SPOT TREATMENT:

For control of annual and susceptible perennial weeds (such as downy bromegrass, oatgrass, goosegrass, puncturevine, goldenrod and plantain) thoroughly mix 2-3 pints of Weedox C R 250 in 3 gallons of water and spray up to 250 square feet to give thorough and uniform coverage.

LARGE AREA TREATMENT: Application rates for large areas depend on climatic conditions. soil type, the weeds present and stage of growth of the weeds. Mix one gallon Weedox C R 250 with 10 gallons water to cover 650 to 1000 square feet. The recommended dosages are equivalent to 43 to 65 gallons of Weedox C R 250 concentrate per acre

The higher rate of application should be used where longer residual control is desired in regions with a long growing season or for control of hard-to-kill perennial weeds and grasses such as Johnson Grass, bindweed and wild carrot. During seasons with excessive rainfail causing heavy weed growth repeat treatment may be required.

Fc+ faster top-kill of existing vegetation, apply Weedox C: R 250 at the above dilution rates in dieseFoil, fuel oil or weed oil.

Weedox C. R 250 should not be used on land to be cropped, or near desirable trees, shrubs or plants, as injury may occur. Do not apply to areas underlaid by roots of desirable trees and shrubs

UNIDER THE FEDERAT FUNGICIDE AND ME FOR ECONOMIC

WEEDOX C/R 250

NON-SELECTIVE HERBICIDE

DOUBLE ACTION — CONTACT & RESIDUAL, Works through foliage contact and root absorption — LONG TERM CONTROL — WATER EMULSIFIABLE

ACTIVE INGREDIENTS: Prometon (2.4-Bis(isopropylam no)-6-methoxy-s-triazine	3.73°c
Pentachlorophenol	
2,3,4,6-Tetrachlorophenol	
Petroleum Distillates	79.41° 5
INERT INGREDIENTS	
TOTAL	00.00° 5

* Pramitol: registered trademark C RA Geigy Corp.

WARNING: KEEP OUT OF REACH OF CHILDREN SEE SIDE PANEL FOR ADDITIONAL CAUTIONS

MANUFACTURED BY

outhsin MILL CREEK PRODUCTS COMPANY, Inc.



NON-SELECTIVE HERBICIDE

DOUBLE ACTION — CONTACT & RESIDUAL, Works through foliage contact and root absorption — LONG TERM CONTROL — WATER EMULSIFIABLE

ACTIVE INGREDIENTS:	
Prometon (2.4-Bis(isopropylam no)-6-methoxy-s-triazine	3.73°°
Pentachlorophenol	1.32°°
2,3.4.6-Tetrachlorophenol	.18°°
Petroleum Distillates	79.41%
INERT INGREDIENTS:	15. 36 %
TOTAL	100 00°°

* Pramitol registered trademark CBA Geigy Corp.

WARNING: KEEP OUT OF REACH OF CHILDREN SEE SIDE PANEL FOR ADDITIONAL CAUTIONS



•

WARNING Keep Out of Reach of Children

Harmful if swallowed. Avoid contact with eyes and skin. In case of skin contact, wash with soap and water. Avoid inhalation of spraymist. Avoid contamination of food and feed.

Do not use, pour, spill or store near heat or open flame. Store at temperatures above 32 degrees F.


PRECAUTIONARY STATEMENT: Do not contaminate water used for irrigation or domestic purposes. Do not store near fertilizers, seeds, insecticides or fungicides. Thoroughly clean equipment with a suitable cleaner before using for other purposes.

FISH AND WILDLIFE CAUTION: This product is toxic to fish and wildlife. Birds and other wildlife in treated areas may be killed. Keep out of lakes, streams, or ponds. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label. Do not reuse empty drum. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place.

NOTICE: Seller's guaranty shall be limited to the terms of the label and subject thereto. The buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions

E.P.A. Reg. No. 6720-232 E.P.A. Est. No. 6720-FL-1

1-20-75



•



3.73% PRAMITOL* VEGETATION KILLER

For Non-Selective Weed Control on Non-Cropland Areas such as <u>Parking</u> Lots, Fence Rows, Railroad Sidings, <u>Storage Yards</u> , Around Buildings of Siles
ACTIVE INGREDIENTS: Prometon (2,4-Bis(isopropylamino)-6-methoxy-3-triazine)*. 3.73% Petroleum distillate
*Pramitol (R), tredemark of Ciba-Geigy Corporation
CAUTION: KEEP OUT OF REACH CT FILDREN
Harmful if swallowed. Avoid contact with eyes and skin. In case of skin contact, wash with soap and water. Avoid inhalation of spray mist.
Avoid contamination of food and feed. Do not contaminate water by cleaning of equipment or dispesal of wastes. Do not contaminate water used for irrigation or domestic purposes.
Do not reuse container. Destroy when empty.
Do not use, pour, spill or store near heat or open flame.
Apply this product only as specified on this label.
Store at temperatures above 32°F.
<u>Important: Read the entire Directions for Use and the Conditions of Sale and Warranty before using this product.</u>
DIRECTIONS FOR USE

Uniformly spray one gallon of VEGETATION KILLER mixed in ten gallons of water over 1000 sq. ft. of soil surface. Thoroughly wet all foliage to insure contact action. Best results are obtained when appli-cations are made during or soon after weed emergence while growth is succulent. Where vegetation is tall and dense, use 1% gallons of

Avoid contamination of food and feed. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not contaminate water used for irrigation or domestic purposes. Do not reuse container. Destroy when empty. Do not use, pour, spill or store near heat or open flame. Apply this product only as specified on this label. Store at temperatures above 32°F.

<u>Important:</u> Read the entire Directions for Use and the Conditions of Sale and Warranty before using this product.

DIRECTIONS FOR USE

Uniformly spray one gallon of VEGETATION KILLER mixed in ten gallons of water over 1000 sq. ft. of soil surface. Thoroughly wet all foliage to insure contact action. Best results are obtained when applications are made during or soon after weed emergence while growth is succulent. Where vegetation is tall and dense, use 1½ gallons of VEGETATION KILLER in ten gallons of water per 1000 sq. ft. For control of hard-to-kill perennial weeds and grasses such as Johnsongrass, Bindweed, Puncturevine, and Wild Carrot, use 3-4½ gallons per 1000 sq. " ft. Although VEGETATION KILLER has considerable activity through foliar contact, much of its activity is through the roots and therefore, adequate rainfall to move the chemical into the root zone is necessary.

Under conditions where the rainfall is high and the growing season long, a second application may be necessary to maintain a weed-free soil surface.

Care must be taken to keep this herbicide away from desirable flower beds, lawns and shrubs. Do not apply over roots of valuable trees, or woody ornamental plants. Only apply herbicide outside of the drip line of trees.

NGTICE: Seller's guaranty shall be limited to the terms of the label and subject thereto. The buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

EPA Registration No. 6720-233

EPA Est. No. 6720-F1-1