



50% SEVIN

MULTI-PURPOSE

WETTABLE POWDER

CAUTION: KEEP OUT OF REACH OF CHILDREN

SEE OTHER PRECAUTIONS ON THE BACK PANEL.

ACTIVE INGREDIENT:

Carbaryl (1-naphthyl methylcarbamate)..... 50% by wt.

INERT INGREDIENTS..... 50% by wt.

TOTAL 100%

FOR USE ON:

- Ornamentals
- Turf and Shade Trees
- Pecans
- Vegetable Gardens
- Lawn Insects as listed

BES-TEX

INSECTICIDES CO., INC.

San Angelo, Texas — EPA Est. No. 06853
EPA Registration No. 6853-18

DO NOT FEED TO LIVESTOCK

Trademark of Union Carbide for the active ingredient 1-naphthyl methylcarbamate.



DIRECTIONS FOR USE

DO NOT use except as specified on this label. Do not use in any manner inconsistent with the statements.

PECANS To control Pecan Nutsaw, apply 5 to 10 lbs. of wettable powder per acre in 300 to 400 gallons of water.

VEGETABLE INSECT CONTROL

Apply when insects or their damage appear. Repeat at 7 to 10 day intervals or as necessary unless shorter interval is specified below.

ON CORN: Use at the rate of 9 tablespoons per 3 gallons of water per 1000 sq. ft. for control of corn earworm, corn rootworm adults, European corn borer, fall armyworm, flea beetles, Japanese beetle, sap beetles and leafhoppers. FOR larvae in which budworm damage and foliage feeders, apply to entire plant. Repeat as necessary. FOR insects attacking silks and ears, apply at 2 to 3 day intervals starting when first silks appear and continuing until silks begin to dry. Three or more applications may be required depending on severity of infestation. Timing and good coverage are absolutely essential for effective control.

ON CUCUMBER, MELONS, PUMPKIN AND SQUASH use at the rate of 5 tablespoons per 3 gallons of water to 1000 sq. ft. for cucumber beetles, flea beetles, leafhoppers and squash bugs. Avoid excessive applications. To avoid possible injury to tender foliage, do not apply when foliage is wet or when rain or excessive humidity is expected during the next 2 days.

ON POTATO, TOMATO, EGGPLANT, AND PEPPER, use 4 tablespoons per 3 gallons of water per 1000 sq. ft. on Colorado potato beetle, flea beetles and leafhoppers. Use 9 tablespoons in 3 gallons of water per 1000 sq. ft. for European corn borer, fall armyworm, lace bugs, tomato fruitworm, tomato hornworm, tarnished plant bugs and stink bugs. On potato, tomato and eggplant use 9 tablespoons in 3 gallons of water per 1000 sq. ft. for cutworm control.

ON OKRA use 9 tablespoons in 3 gallons of water per 1000 sq. ft. for corn earworm and stink bugs. Repeat at 5 to 7-day intervals or as necessary.

ON BEANS (GREEN BEANS, LIMA BEANS, SNAP BEANS, AND BLACK-EYED PEAS) use 2 tablespoons in 3 gallons of water per 1000 sq. ft. for Mexican bean beetle. Use 5 tablespoons in 3 gallons of water per 1000 sq. ft. for bean leaf beetle, cucumber beetles, flea beetles, Japanese beetle, leafhoppers, velvetbean caterpillar, and western bean cutworm. Use 6 tablespoons in 3 gallons of water per 1000 sq. ft. for armyworms, cutworms, corn earworm, stink bugs and tarnished plant bug.

ORNAMENTAL INSECT CONTROL

Use 2 pounds of SEVIN* in 150 to 200 gallons of water (6 tablespoons per 3 gallons of water) for each 5000 sq. ft. of established lawn area for the control of chinch bugs, leafhoppers, sodwebworms (lawn moths), Fall armyworm, cutworms, earwigs, ants, fleas, mosquitoes, and millipedes.

MIXING INSTRUCTIONS

Add recommended amount of SEVIN* to small amount of water and mix thoroughly before adding to applicator tank. Keep spray mixture well agitated.

WARRANTY

Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

NET WT. 2 POUNDS

Repackaged by
BES-TEX INSECTICIDES CO., INC.
San Angelo, Texas — EPA Est. No. 06853
EPA Registration No. 6853-18