

## SEVIN 5% BAIT CROP INSECTICIDE

(PELLETIZED)

**ACTIVE INGREDIENT:** 

Carbaryl (1-napthyl N-methylcarbamate) 5%
INERT INGREDIENTS: 95%
100%

Sevin is Reg. Trademark of Union Carbide Corporation. U.S. 2-903-478

## CAUTION KEEP OUT OF REACH OF CHILDREN

SMCP Sevin 5% Bait Crop Insecticide is an effective insecticide which can be used to control several common insects on a variety of crops.

Following the Rate Table below — Use as a Broadcast Application

INSECTS	CROP	RATE PER ACRE	LIMITATIONS
Armyworms, Cutworms, Crickets, Darkling beetles, Grasshoppers, Sowbugs	Alfalfa	30 lbs.	To avoid possible injury to tender foliage, do not apply when foliage is wet or when rain or excessive humidity is expected during the few 1 days.
Ants,	Apples, Cherries, Plums	40-50-Hrs.	d penarsest Interval
Crickets,	Apricots	44.50 lbs.	≯day preharvest interval∖
Cutworms, Grasshoppers, Sowbugs	Citrus (Grapefruit, Kumquats, Lemons, Limes, Nectarines, Oranges, Tangelos, Tangerines), Grapes, Peaches, Pears	10:50 lbs.	EP 27 1970  Get and for the part of the pa
Armyworms, Crickets, Cutworms,	Beans, Carrots, Corn, Okra, Tomatoes, Eggplant, Potatoes, Peppers,	Land, Digas	dde, and flow 155  nonce tor two 1 255  nonce tor t
Darkling ground beetles,	Asparagus, Strawberries	40 bseq1	le menarvest interval
Grasshoppers, Sowbugs	Beets (roots), Broccoli, Brussel Sprouts, Cabbage, Cauliflower, Lettuce, Horseradish, Parsnips, Radishes, Rutabagas, Turnips	40 1	3 day preharvest interval
Armyworms, Crickets, Cutworms, Darkling ground beetles, Grasshoppers, Sowbugs	Collards, Dandelions, Endive, Kale, Spinach, Parsley, Sugar Beets, Swiss Chard, Beets (tops)	40 lbs.	14 day preharvest interval
	Cotton, Peas	30 lbs.	
Armyworms, Crickets, Cutworms, Darkling Beetles, Grasshoppers, Sowbugs	<ul> <li>Cucumbers, Melons (Cantaloupes, Casabas, Crenshaw, Honey balls, Honeydew Melons, Muskmelons, Persian Melon, Watermelons), Peanuts, Squash</li> </ul>	•	To avoid possible injury to tender foliage, do not apply when foliage is wet or when rain or excessive humidity expected during the next 2 days. Carbaryl injures watermelons in Florida.
Crickets, Cutworms,	Tobacco (in plant beds)	1/2 to 1 lb. per 1000 sq. ft.	
Armyworms	Tobacco (in fields)	20-40 lbs. per acre	

## CAUTION

Human Cautions: Keep out of reach of children. Harmful if inhaled or swallowed. DO NOT breathe dust. Avoid contact with skin or eyes. Wear regular long sleeved clothing. Avoid storage near food or feedstuffs. Wash hands thoroughly after handling and before eating or smoking. Do not wear contaminated clothing. Wash clothing in hot water and soap before re-using.

NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

Environmental Cautions: This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label. This product is highly toxic to bees exposed to direct treatment or residues. Consult State Agricultural Extension Service for additional information.

Do not reuse container. Destroy when empty.

**NOTICE:** Seller's guaranty shall be limited to the terms of the label and subject thereto. The buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

E.P.A. Reg. No. 6720-255

E.P.A. Est. No. 6720-FI-1

**MANUFACTURED BY** 

## Southern Mill Creek PRODUCTS COMPANY, INC.

TAMPA, FLORIDA 33601