

HOME LAWN & ORNAMENTAL SPRX

containing 5.-0% Chlorpyrifos

Controls Chinch Bugs, Sod Webworms (Lawn Moths), Armyworms, Ants, Fleas, Brown Dog Ticks, Cutworms, Earnings, Crickets, Grasshoppers and Hyperodes Weevil infesting home lawns and other owners. US OFFE and other ornamental turf areas. Also controls Aphids, Bagworms, Grasshoppers, Leafhoppers, 5 Spittle Bugs, Mealybugs, Whiteflies, exposed Thrips, Mites and Cutworms infesting ornamental piants.

ACTIVE INGREDIENTS:		
Chlorpyrifos [O,O-diethyl O-(3,5,6-trich	nloro-2-	
pyridyl) phosphorothicate *		5.40%
Aromatic Petroleum Derivative Solvents		91.24%
INERT INGREDIENTS:		
TOTAL		00.00%

CAUTION: KEEP OUT OF REACH OF CHILDREN

SEE SIDE PANEL FOR OTHER INFORMATION AND CAUTIONS E.P.A. Reg. No. 6720-257 E.P.A. Est. No. 6720-FL-1

'U. S. Patent No. 3,244,586

MANUFACTURED BY outhern MILL CREEK PRODUCTS COMPANY, Inc.

DOSAGE & DILUTION FOR USE ON ORNAMENTAL PLANTS

AMOUNT O	F LAWN & ORNAMENTA WATER TO MAKE	AL SPRAY IN
1 Gal. 1 fl, oz.	3 Gal. 2 ¹ / ₂ fl. oz.	10 Gal. ¹/₂ pt. •
– 2 fl. oz.	5 fl. oz.	1 pt.
ntil spray has dried. Kee contaminated with this	ep out of lawn and garde	n fish ponds and othe
CAUTION	F	
	1 Gal. 1 fl. oz. 2 fl. oz. ntil spray has dried. Kelectontaminated with this	1 Gal. 3 Gal. 1 fl. oz. 21/2 fl. oz. 2 fl. oz. 5 fl. oz. ntil spray has dried. Keep out of lawn and garde contaminated with this product.

SPILL, OR STORE FEAR HEAT OR OPEN FLAME. Avoid contact with eyes, skin and clothing. Do not take internally, Wash thoroughly at a handling. Wash contaminated clothing before re-use. Flush contaminated eyes with pienty of water and get prompt medical attention. Avoid breathing vapors and spray mist. Avoid spraying food crops. Keep away from food. regordifs and domestic water supplies. Keep container closed. Handle concentrate in a ventilated area.

NOTE TO PHYSICIAN: Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically; atropine is an antidote.

GENERAL USE PRECAUTIONS

This product is highly toxic to bees exposed to direct treatment or residues on plants. Protective information may be obtained from your Cooperative Agricultural Extension Service.

This product is toxic to fish, birds and other wildlife. Keep cut of any body of water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning equipment or disposal of wastes. Apply this product only as specified on this label.

Promptly crush empty containers and bury them and wastes in locations away from domestic water supplies.

Do not reuse empty drum. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place away from water supplies.

NOTICE: Seller's guaranty shall be limited to the terms of the label and subject thereto. The buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

Pests of Lawns and other Ornamental and Recreational Turf Grass Areas:

To control Ants, Brown Dog Ticks, Chinch Bugs, Crickets, Cutworms, Earwigs, Fleas, Armyworms, Grasshoppers, and Sodweb Worms, (Lawn Moths). Mix SMCP LAWN & ORNAMENTAL SPRAY according to directions given in the following table and apply as a coarse low pressure spray. For best results the lawn should be moist at the time of treatment, and care should be taken to obtain complete and uniform coverage of pest infested areas. Spray when pests first appear; retreat when needed. Control for Chinch Bugs will last up to 8 weeks. To prevent damage from Chinch Bugs apply as soon as daily temperatures reach 80°F and repeat every 2 months.

DOSAGE AND DILUTION FOR USE ON TURF

Area to be Treated	Amount of SMCP Lawn & Ornamental Spray used	Dilution with water to make
500 Sq. Ft.	31 ¿ fluid ounces	15 gallons
1,000 Sq. Ft.	7 fluid ounces	30 gallons
5,000 Sq. Ft.	35 fluid ounces	150 gallons
10,000 Sq. Ft.	75 fluid ounces (2 qts. 11 oz.)	325 gallons

To control Turfgrass (Hyperodes) Weevils use twice the amount of SMCP LAWN & ORNAMENTAL SPRAY shown in the above Turf Table. Spray suspected problem areas in mid-April and again in mid-May. Consult your local Agricultural Extension Service specialist for proper time to treat in your area.

Note: To help prevent infestation of buildings by Ants, Crickets, Earwigs and Sowbugs (Pillbugs), treat a band of soil 6 to 10 feet wide around and adjacent to the building, also the building foundation to a height of 2 to 3 feet, where pests are active and may find entrance. Use the same dosage and dilution recommended for control of these pests on turf as shown in the above table. Apply as a coarse spray to thoroughly and uniformly wet the band area.

For standard 15 gallon hose-end applicators add 4 fl. oz. to quart bottle full of water. This will treat 500 sq. ft.

DO NOT MOW LAWN FOR 24 HOURS FOLLOWING TREATMENT Pests of Ornamentals:

Use SMCP LAWN & ORNAMENTAL SPRAY to control Aphids, Bagworms, Cutworms, Grasshoppers, Leafhoppers, Mealybugs, Mites, Spittlebugs, Thrips (exposed) and Whiteflies on flowers, shrubs, vines, shade and flowering trees and evergreens found to be infested with these pests. Dilute SMCP LAWN & ORNAMENTAL SPRAY with water according to directions given in the following table and apply using suitable hand or power spray equipment in a manner to provide complete and uniform coverage of infested plants. Avoid overspraying to the point of excessive run off. Treat when pests appear and repeat at 7 to 10 day intervals, if needed.

Note: Do not use on Azaleas, Camellias, Poinsettias, Rose Bushes or Variegated Ivy because of possible injury to these plants.

22