


FRONT PANEL
EPA No. 6720-.....

TRAC OFTANOL INSECTICIDE
1.5% GRANULAR
Controls Mole Crickets and Grubs



ACTIVE INGREDIENT:	
1-Methylethyl 2-[[ethoxy[(1-methylethyl)amino] oxy]	
benzoate	1.5%
INERT INGREDIENTS	98.5%
TOTAL	100.0%

U.S. PATENT No. 3,621,082 CANADIAN PATENT No. 825,198
OFTANOL is a Reg. TM of the Parent Company of Farbenfabriken Bayer
GmbH, Leverkusen.

KEEP OUT OF REACH OF CHILDREN
CAUTION
PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
Caution

Harmful if swallowed. Avoid contact with skin, eyes, or clothing. Wash contaminated clothing with soap and warm water after handling. Wash contaminated clothing with soap and hot water before reuse. Keep out of reach of children.

STATEMENTS OF PRACTICAL TREATMENT

If in eyes, flush with plenty of water. If on skin, wash with soap and hot water.
If swallowed, call a physician or Poison Control Center. Give a glass or two of water and induce vomiting by touching finger to back of throat. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and wildlife. Do not apply directly to water and wetlands. Keep out of lakes, streams and ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

MANUFACTURED BY
SOUTHERN MILL CREEK PRODUCTS CO., INC.
TAMPA, FLORIDA 33601

FRONT PANEL
TRAC OFTANOL 1.5%
EPA No. 6720-.....

STORAGE AND DISPOSAL
DO NOT CONTAMINATE WATER, FOOD OR FEED BY STORAGE AND DISPOSAL

HOUSEHOLD: Do not reuse empty container. Wrap and put in trash collection. Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of reach of children.

Handle any open container in a manner as to prevent spillage. If the container is leaking or material spilled for any reason or cause, carefully sweep material into a pile. Do not walk through spilled material. Dispose of pesticide as directed above.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

IMPORTANT: Read these entire Directions for Use and Conditions of Sale before using OFTANOL 1.5% Granular insecticide. OFTANOL 1.5% Granular Insecticide is recommended for use on lawns, and other ornamental and recreational turf grass areas.

APPLICATION EQUIPMENT

Apply OFTANOL 1.5% Granular uniformly over the turf area with suitable granular turf application equipment. Both drop-type and rotary-type spreaders may be used to apply OFTANOL 1.5% Granular. Avoid the use of spreaders which would apply the material in narrow, concentrated bands. Calibrate application equipment prior to use according to manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly.

(DIRECTIONS FOR USE CONTINUED ON BACK PANEL)

NET WEIGHT:

EPA Reg. No. 6720-600

EPA EST. No. 6720-FL-1

BACK PANEL
TRAC OFTANOL 1.5%

DIRECTIONS FOR USE (Cont'd)
APPLICATION TECHNIQUES

OFTANOL 1.5% Granular application using a pull-type drop spreader can be aided by a splash board mounted under the hopper for more uniform granule distribution. A more uniform application can be made by spreading half the required amount of OFTANOL 1.5% Granular over the area, and then immediately apply the remaining half at right angles or crosswise. The full recommended rate of OFTANOL 1.5% Granular must be used to maximum benefit.

When applying OFTANOL 1.5% Granular, use appropriate marking systems and make application when the wind is calm to prevent missed areas.

SPREADER SETTINGS

Approximate Spreader Setting's for applying of OFTANOL 1.5% Granular at the Rate of 3 pounds per 1000 sq. ft.

MODEL	SETTING
Cyclone model B-1	5
Scotts PF-1/011B	10

RECOMMENDED APPLICATIONS

CROP	PEST	DOSAGE OFTANOL 1.5% G	REMARKS
<hr/>			
	Soil Insects		
Turf Grasses	White Grub larvae (such as Japanese beetle Black turf grass ataenius, Chafers, Phyllophaga Green June beetle, Oriental beetle, Asiatic garden beetle)	3 lbs/1000 sq. ft.	Be sure to read "Application Equipment" Section of this Label Do not apply more than 6 lbs of Oftanol 1.5% Granular per 1000 sq. ft. per year or an equivalent of 1.45 ounces acti- ve ingredient per 1000 sq. ft. regardless of formulation used; Minimum interval between appli- cations must be 30 days.
	Billbug (larval stage)		For White Grub and Billbug Larval Control: Water turf area thoroughly immediately after application, using sufficient water to wet soil 1 to 1/2 inches deep.
	Hyperodes weevil (larval stage)		
	Mole Crickets		

BACK PANEL (Cont'd)
TRAC OFTANOL 1.5%
EPA No. 6720-

CROP	PEST	DOSAGE OFTANOL 1.5% G.	REMARKS
			For Hyperodes Weevil Control: Applying between full bloom periods of for- sythia and flowering dog- wood, usually April 15 to May 15, and water-in as for white grub larval control. For Mole Cricket Control: Water immediately after appli- cation using $\frac{1}{2}$ to 1 inch water.
Fire Ants	3 lbs/1000 sq. ft.		Area Fire Ant Control: Apply specified dosage uniformly over area to be treated for residual control. Water im- mediately after application using $\frac{1}{2}$ to 1 inch of water. If established fire ant mounds are present, use the mound drench treatment prior to area application. If a mound drench treatment is made only one area treatment may be made. Established Fire Ant Mounds Drench Treatments: Wet Conditions- Distribute dosage evenly over 9 sq. ft. and immediately sprinkle gently with 3 gals. of water. Dry Conditions-Treat as noted to insure penetration apply an additional 3 gals of water to the treated mound area after treatment. Make only one mound treatment. Consult your local turf, state, Agricultural Extension Service Specialists for more specific information regarding timing one mound drench treatment.
	0.5 oz per 9 sq. ft.		

BACK PANEL (Cont'd)
TRAC OFTANOL 1.5%
EPA REG. No. 6720-

CROP	PEST	DOSAGE OFTANOL 1.5% G.	REMARKS
	Surface Insects Sod webworm larvae Chinch Bug	1.5 or 3 lbs /1000 sq. ft.	Apply $\frac{1}{2}$ to 1 inch of water immediately after treatment. Reapply as necessary for pest infestations. Do not apply more than 6 lbs OFTANOL 1.5% Granular per 1000 sq. ft. per year or an equivalent amount of 1.45 ounces active ingredient per 100 sq. ft. regardless of formulation used. Minimum interval between application is 15 days for 1.5 lb rate and 30 days for the 3 lb rate. Consult your local turf, state Agricultural Experiment Station or State Extension Service Specialists for more specific information regarding timing of application.

RESTRICTIONS

Do not mow treated turf area before rainfall or irrigation has been applied as per RECOMMENDED APPLICATIONS. Do not graze treated areas or use clippings from treated areas for feed or forage. Keep children and pets off treated area until dry.

CONDITIONS OF SALE: THE DIRECTIONS ON THIS LABEL WERE DETERMINED THROUGH RESEARCH TO BE THE DIRECTIONS FOR CORRECT USE OF THIS PRODUCT. THIS PRODUCT HAS BEEN TESTED FOR A RANGE OF WEATHER CONDITIONS SIMILAR TO THOSE WEATHER CONDITIONS THAT ARE ORDINARY AND CUSTOMARY IN THE GEOGRAPHIC AREA WHERE THE PRODUCT IS USED. INSUFFICIENT CONTROL OF PESTS MAY RESULT FROM THE OCCURRENCE OF EXTRAORDINARY OR UNUSUAL WEATHER, OR FROM FAILURE TO FOLLOW LABEL DIRECTION. IN ADDITION, FAILURE TO FOLLOW LABEL DIRECTIONS MAY CAUSE INJURY TO THE CROP TO WHICH THE PRODUCT IS APPLIED, OTHER CROPS, ANIMALS, MAN OR THE ENVIRONMENT. SOUTHERN MILL CREEK PRODUCTS COMPANY, INC. AND THE SELLER OFFER, AND THE BUYER ACCEPTS AND USES THIS PRODUCT SUBJECT TO THE CONDITIONS THAT EXTRAORDINARY OR UNUSUAL WEATHER, OR FAILURE TO FOLLOW LABEL DIRECTIONS ARE BEYOND THE CONTROL OF SOUTHERN MILL CREEK PRODUCTS COMPANY INC. AND ARE, THEREFORE, THE RESPONSIBILITY OF THE BUYER