



MANEB Notice 62-22
GARDEN FUNGICIDE 11/15/62
 CONTROLS BLIGHT, ANTHRACNOSE, LEAF SPOT
 and certain other diseases on TOMATOES
 and on other Vegetables, Fruit, Flowers as listed

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS: CAUTION— Harmful if swallowed. May cause irritation of nose, throat, eyes and skin. Do not breathe dust or spray mist. Avoid contact with skin, eyes and clothes. Protective clothing (long pants, long sleeved shirt, hat, boots and gloves) must be worn during mixing. Food utensils such as a teaspoon or measuring cup should not be used for food after use with pesticide. Keep children and pets out of treated areas until dry. Do not contaminate water, food or feed. **ENVIRONMENTAL HAZARD:** This product is toxic to fish. Do not apply directly to water or wetlands. Do not apply where runoff is likely to occur or when weather conditions favor drift from treated area. Do not contaminate water by direct application or cleaning of equipment. **PHYSICAL AND CHEMICAL HAZARD:** Keep away from fire and sparks. Do not allow to become wet or overheated in storage. This may bring on a chemical change which will impair the fungicidal effectiveness and generate flammable vapors.

DIRECTIONS FOR USE:
 It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Store this product in its closed, original container in a locked area away from children and domestic animals. Do not reuse empty container. For disposal, securely wrap original container in several layers of newspaper and discard in trash.

DIRECTIONS

Controls diseases indicated on plants listed. Unless otherwise directed use 1 1/2 level tablespoons per gallon of water to make spray mix. Thoroughly spray both upper and under surfaces of foliage. Unless otherwise directed begin when disease first appears, repeat every 7 to 10 days as needed. Spray more frequently if infestation is severe. When a number in parentheses follows the plant name, do not spray that crop within the number of days of harvest shown.

VEGETABLES

TOMATOES (5): • Anthracnose, early and late blight, gray leaf spot (stemphylium), septoria leaf spot. Begin application when tomatoes on first cluster are well formed. Do not use on young plants in greenhouse. **BEANS (4):** green and dry. Downy mildew, rust. Begin when plants are small. Repeat every 5 to 7 days. **BROCCOLI:** • Alternaria leaf spot, downy mildew. If applied within 3 days of harvest remove residues by washing or trimming. **BRUSSELS SPROUTS, CAULIFLOWER, KOHLFABI**

• Alternaria leaf spot, downy mildew. **CABBAGE (7):** Alternaria leaf spot, downy mildew. Begin 7 to 10 days after planting. Repeat every 3 to 5 days. **CARROTS:** • Leaf spot diseases. Start spraying when plants are 6 weeks old or disease first appears. **CELERY (14):** • Early and late blight. Start spraying when plants are established in the garden. Remove residues at harvest by stripping, trimming and washing. **CUCUMBERS (5):** **SQUASH (5):** • Anthracnose, downy mildew, pythium fruit rot. Begin when disease first appears or runners start to form. **CANTALOUPE (5):** **WATERMELON (5):** • Anthracnose, downy mildew, cercospora leaf spot, gummy stem blight. Begin when vines start to run or disease first appears. **EGGPLANT, PEPPERS:** • Anthracnose, fruit rot. Apply as fruit begins to form. Repeat weekly through harvest. **LETTUCE (10):** **ENDIVE (10):** • Downy mildew. Remove residues at harvest from head lettuce by stripping and trimming outer leaves, and from leaf lettuce by dividing by washing. **ONIONS:** • Botrytis leaf blight, downy mildew, purple blotch. **PUMPKINS:** • Angular leaf spot, downy mildew. Use 3 tablespoons per gallon. Spray thoroughly. **SPINACH (11):** • Downy mildew, white rust. Remove residues at harvest by washing.

FLOWERS

Begin spraying when new growth starts or when leaves first unfold. Apply at weekly intervals during season.

CARNATIONS, PANSIES: anthracnose, leaf spot. **DAHLIAS, ZINNIAS:** botrytis, leaf spot. • **FLOWERING DOGWOOD:** anthracnose; apply when buds begin to open and again when bracts fall, and 4 weeks later, and again in late summer. **GLADIOLUS:** botrytis, curvularia, Stemphylium leaf spot. **PEONY:** botrytis and phytophthora blights. Apply in early spring and early fall to foliage and to soil around plants. Repeat at 7 to 10 day intervals during growing season. **ROSES:** black spot. **SNAPDRAGON:** rust.

APPLES (30)

Black rot, bitter rot, brown rot, bullseye rot, fly speck, rust, scab, sooty blotch, white rot. Begin application in the spring. Repeat in subsequent pre-bloom, bloom and post-bloom sprays in accordance with local state spray schedule. Do not spray within 30 days of harvest.

LIMITED WARRANTY: BUYER ASSUMES ALL RISKS OF STORAGE AND HANDLING OF THIS MATERIAL. NOT TO BE STRICT ACCORDANCE WITH DIRECTIONS GIVEN HEREON.

FORM 100-1000
 COPYRIGHT 1983 MILLER CHEMICAL & FERTILIZER CORPORATION

BEST AVAILABLE COPY