

# SEVIN<sup>5</sup>

## ACTIVE INGREDIENT:

Carbaryl (1-naphthyl N-methylcarbamate) ..... 5.0%

INERT INGREDIENTS: ..... 95.0%

100.0%

\*Trademark of UCC for 1-naphthyl N-methylcarbamate.

## CAUTION

- KEEP OUT OF REACH OF CHILDREN AND ANIMALS
- HARMFUL IF SWALLOWED OR INHALED.
- DO NOT TAKE INTERNALLY.
- DO NOT BREATHE DUST.
- AVOID PROLONGED BREATHING OF DUST.
- SKIN CONTACT MAY BE HARMFUL. AVOID CONTACT.
- WEAR REGULAR LONG SLEEVED WORK CLOTHING. CHANGE TO CLEAN CLOTHING DAILY.
- WASH HANDS BEFORE EATING. TAKE SHOWER OR BATH AFTER WORK.
- NOTE FOR PHYSICIANS: CARBARYL IS A MODERATE REVERSIBLE, CHOLINESTERASE INHIBITOR. ATROPINE IS ANTIDOTAL.

### ENVIRONMENTAL HAZARDS

The Federal Food and Drug Administration has set official tolerances for certain pesticide residues. In order that residues on food and forage crops will not exceed established tolerances, use only at recommended rates, and do not apply closer to harvest than specified. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Apply when air is calm, using ground equipment, aircraft, or hand duster. Make thorough coverage of plants with a uniform deposit of dust. Apply at the first sign of infestation and repeat at regular intervals when needed to maintain control.

### RE-ENTRY

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

Do not enter treated areas until dust has settled unless appropriate protective clothing is worn. Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. (Indicate specific oral warnings which inform workers of areas or fields that may not be entered without specific protective clothing, period of time field must be vacated and appropriate actions to take in case of accidental exposure.) When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information: "Caution" area treated with Sevin 5 on (date of application). Do not enter without appropriate protective clothing until dust has settled.

If accidental exposure occurs, see caution statements on left panel of label.

### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. STORAGE: Do not store in or around the home. Store in a secure, dry and temperate area. Store only in original container. Keep away from heat, sparks or open flame. In case of spill sweep up. Place in drum, label contents and dispose of according to the instructions below in PESTICIDE DISPOSAL. CAUTION: Wear a pesticide respirator and avoid forming dust clouds during clean-up.

### PESTICIDE DISPOSAL


Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

### CONTAINER DISPOSAL - PAPER BAGS

Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

E.P.A. Registration No. 6735-14

E.P.A. Est. No. 6735-TX-1



NET WT. 50 LBS.

A PRODUCT OF

**TIDE PRODUCTS, INC.**

EDINBURG, TEXAS 78539

## AN AGRICULTURAL INSECTICIDE

### RECOMMENDATIONS

#### COTTON:

For control of early season Thrips and Cotton Fleahopper:

Apply 20 to 30 lbs. of Sevin 5 Dust per acre when insects first appear.

For control of Mid and Late-season Boll Weevil, Bollworm, Cotton Leafworm, Lygus Bugs, and Cotton Leaf Perforators:

Apply 30 to 50 lbs. of Sevin 5 Dust per acre at 5-7 day intervals when insects first appear.

For control of Pink Bollworms:

Apply 45 to 50 lbs. of Sevin 5 Dust per acre at 5-7 day intervals.

Aphid populations will usually be repressed when this insecticide is used on a regular schedule of treatments. Where mite infestations are encountered, use a recommended miticide. SEVIN is believed compatible with all pesticides commonly used on cotton.

**BEANS:** For control of Mexican Bean Beetle, apply 20-30 lbs. per acre. Leafhopper, apply 30 lbs. per acre and Tarnished Plant Bug, apply 25-30 lbs. per acre at 7-10 day intervals. Must be applied the day of harvest.

**GRAPES:** For control of Grape Leaf Roller, Grape Berry Moth and Grape Leafhopper, apply 20-40 lbs. per acre at 7-10 day intervals. May be applied up to harvest.

**POTATOES:** For control of Colorado Potato Beetle, Potato Leafhopper, Six Spotted Leafhopper and Potato Flea Beetle, apply 15-20 lbs. per acre, and Lygus Bugs, apply 25-30 lbs. per acre at 7-10 day intervals. Foliage application only. May be applied up to harvest.

**STRAWBERRIES:** For control of Meadow Spittlebug and Strawberry Leaf Roller, apply 25-35 lbs. per acre at 7-10 day intervals up to 1 day before harvest.

**CUCUMBERS and SUMMER SQUASH:** For control of Striped and Spotted Cucumber Beetles, Squash bugs and Flea Beetles, apply 15-20 lbs. per acre at 7-10 day intervals. May be applied up to harvest.

**TOMATOES, EGGPLANT and PEPPERS:** For control of Flea Beetles apply 15-20 lbs. per acre at 7-10 day intervals. For control of European Corn Borer, Armyworm, Tomato Fruitworm and Tomato Hornworm apply 30-40 lbs. per acre at 7-10 day intervals. For control of Stink Bugs apply 40-50 lbs. per acre. May be applied up to harvest.

**CORN:** For control of Corn Earworm, European Corn Borer, Fall Armyworm, Sap Beetle, Japanese Beetle, apply 20-40 lbs. per acre at 7-10 day intervals up to day of harvest.

**LETTUCE:** For control of Flea Beetles, Leafhoppers, and Harlequin Bugs, apply 15-20 lbs. per acre. To control the Six-Spotted Leafhopper, apply 20-30 lbs. per acre. To control Corn Earworm, Armyworm, Cabbage Caterpillars, Spittlebugs, Lygus and Stink Bugs, apply 30-40 lbs. per acre. Make application at a 7-10 day interval. On head lettuce do not apply later than 3 days before harvest. On leaf lettuce, do not apply within 14 days of harvest.

**OKRA:** For control of Corn Earworm and Stink Bugs, apply 30-40 lbs. per acre at a 5-7 day interval up to and including the day of harvest.

**ALFALFA, CLOVERS, COWPEAS (SOUTHERN OR FIELD PEAS), PEANUTS, SOYBEANS:** For control of Blister Beetles and Mexican Bean Beetle, use 15 to 30 lbs. per acre. To control Thrips, Leafhoppers, Three-Cornered Alfalfa Hopper, Cucumber Beetles, Bean Leaf Beetle, Japanese Beetle, Alfalfa Caterpillar and Velvet Bean Caterpillar, apply 25-30 lbs. per acre. To control Armyworms, Webworms, Corn Earworm and Stink Bugs, apply 30 lbs. per acre. May be applied up to day of harvest.

**RICE, FORAGE GRASSES AND PASTURE:** For control of Thrips, Armyworms, and Stink Bugs, apply 25-35 lbs. per acre. Do not apply within 14 days of harvest on rice.

**SORGHUMS: (including MILO and GRAIN SORGHUM)** For control of Armyworms, Webworms, Corn Earworms and Stink Bugs, apply 30-40 lbs. per acre. Do not apply within 21 days of harvest on grain sorghum. May be applied day of harvest for forage.

**NOTICE:** Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty or merchantability or fitness for a particular purpose, expressed or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

GP-784-(3853)