



# SEVIN\* 10

AN AGRICULTURAL INSECTICIDE

**ACTIVE INGREDIENT**

Carbaryl (1-naphthyl N-methylcarbamate)

10.0%

**INERT INGREDIENTS:**

90.0%

100.0%

\*Trade-Mark of UCC for 1-naphthyl N-methylcarbamate.

## CAUTION

- **KEEP OUT OF REACH OF CHILDREN AND ANIMALS**
- **HARMFUL IF SWALLOWED OR INHALED.**
- **DO NOT TAKE INTERNALLY.**
- **DO NOT BREATHE DUST.**
- **AVOID PROLONGED OR REPEATED BREATHING OF DUST.**
- **SKIN CONTACT MAY BE HARMFUL. AVOID UNNECESSARY CONTACT.**
- **WEAR REGULAR LONG SLEEVED WORK CLOTHING. CHANGE TO CLEAN CLOTHING DAILY.**
- **WASH HANDS BEFORE EATING. TAKE SHOWER OR BATH AFTER WORK.**
- **NOTE FOR PHYSICIANS: CARBARYL IS A MODERATE REVERSIBLE, CHOLINESTERASE INHIBITOR. ATROPINE IS ANTIDOTAL.**

### OBSERVE THE FOLLOWING USE PRECAUTIONS

The Federal Food and Drug Administration has set certain tolerance levels for certain pesticides. In order that residue on food and forage crops will not exceed established tolerance levels, it is recommended that you do not apply insecticide to harvest crops specified.

This product is not to be used in any manner which would be injurious to man or animals. For more information, may be obtained from your Cooperative Extension Service.

### DIRECTIONS FOR USE

Apply when an insecticide is needed. Do not apply to plants with a surface wet with rain. Apply at the rate of 10-20 lbs. per acre at 7-10 day intervals when needed to control insects.

### RECOMMENDATIONS

#### COTTON

For control of European Corn Borer, Armyworm, Tomato Fruitworm and Tomato Hornworm, apply 10-15 lbs. per acre at 7-10 day intervals. May be applied up to day of harvest.

For control of Mexican Cotton Bollworm, Tobacco Budworm, Cotton Leafworm, and Cotton Leaf Perforator, apply 10-15 lbs. per acre at 7-10 day intervals. May be applied up to day of harvest.

For control of Pink Bollworm, apply 10-15 lbs. per acre at 7-10 day intervals. May be applied up to day of harvest.

Aphid populations may build up rapidly on cotton. The use of SEVIN 10 is recommended where necessary. Where necessary, SEVIN 10 is compatible with all registered insecticides for use on cotton.

**BEANS:** For control of Mexican Bean Beetle, apply 10-15 lbs. per acre. Leafhoppers, apply 18 lbs. per acre and Tarnished Plant Bug, apply 12-18 lbs. per acre at 7-10 day intervals. May be applied the day of harvest.

**GRAPES:** For control of Grape Leaf Roller, Grape Berry Moth and Grape Leafhopper, apply 12-20 lbs. per acre at 7-10 day intervals. May be applied up to harvest.

**POTATOES:** For control of Colorado Potato Beetle, Potato Leafhopper, Six Spotted Leafhopper and Potato Flea Beetle, apply 7-10 lbs. per acre, and Lygus Bugs, apply 12-18 lbs. per acre at 7-10 day intervals. Foliage application only. May be applied up to harvest.

**STRAWBERRIES:** For control of Meadow Spittlebug and Strawberry Leaf Roller, apply 12-18 lbs. per acre at 7-10 day intervals up to 1 day before harvest.

**CUCUMBERS and SUMMER SQUASH:** For control of Striped and Spotted Cucumber Beetles, Squash Bugs and Flea Beetles, apply 8-10 lbs. per acre at 7-10 day intervals. May be applied up to harvest.

**TOMATOES, EGGPLANT and PEPPERS:** For control of Flea Beetles, apply 7-10 lbs. per acre at 7-10 days interval. For control of European Corn Borer, Armyworm, Tomato Fruitworm and Tomato Hornworm, apply 10-20 lbs. per acre at 7-10 day intervals. For control of Stink Bugs, apply 25-30 lbs. per acre. May be applied up to harvest.

**CORN:** For control of Corn Earworm, European Corn Borer, Fall Armyworm, Sap Beetle, Japanese Eggrug, apply 10-20 lbs. per acre at 7-10 day intervals up to day of harvest. Except when used for storage do not apply within 7 days of harvest.

**LETTUCE:** For control of Flea Beetles, Leafhoppers, and Harlequin Bugs, apply 7-10 lbs. per acre. To control the Six-Spotted Leafhopper, apply 10-15 lbs. per acre. To control Corn Earworm, Armyworm, Cabbage Caterpillar, Spittlebugs, Lygus and Stink Bugs, apply 10-20 lbs. per acre. Make application at a 7-10 day interval. On Head lettuce do not apply later than 3 days before harvest. On Leaf Lettuce do not apply within 14 days of harvest.

**OKRA:** For control of Corn Earworm and Stink Bugs, apply 12-20 lbs. per acre at a 7-10 day interval up to and including the day of harvest.

**ALFALFA, CLOVERS, COWPEAS, SOUTHERN OR FIELD PEAS, PEANUTS, SOYBEANS:** For control of Flea Beetles and Mexican Bean Beetle, use 7 to 14 lbs. per acre. To control Thrips, Leafhoppers, Three-Cornered Alfalfa Hopper, Cucumber Beetles, Bean Leaf Beetle, Japanese Beetle, Alfalfa Caterpillar and Velvet Bean Caterpillar, apply 12-15 lbs. per acre. To control Armyworms, Webworms, Corn Earworm and Stink Bugs, apply 14-18 lbs. per acre. May be applied up to day of harvest.

**RICE, FORAGE GRASSES AND PASTURE:** For control of Thrips, Armyworm, and Stink Bug, apply 14-15 lbs. per acre. Do not apply within 14 days of harvest on rice.

**SORGHUMS:** (including **M.L.O.** and **GRAIN SORGHUM**): For control of Armyworms, Webworms, Corn Earworm and Stink Bug, apply 14-20 lbs. per acre. Do not apply within 21 days of harvest on grain sorghum. May be applied up to harvest for forage.

**NOTICE:** Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty or merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

BURN OR BURY BAG WHEN EMPTY. DO NOT REUSE.

A PRODUCT OF  
**TIDE PRODUCTS, INC.**  
EDINBURG, TEXAS