METHYL

Emulsifiable Concentrate Insecticide

PROTECT FROM TEMPERATURES BELOW 32°F

ACTIVE INGREDIENTS	
O, O-dimethyl O-p-Nitrophenyl phosphorothioate	44.21
*Related Compounds	2.28
Xylene Range Aromatic Solvent	48.24
INERT INGREDIENTS	5.27
	100.00

* Methyl Parathion

Contains
4 lbs. Methyl Parathion
per Gallon

E.P.A. Est. No. 6735-TX-10

E.P.A. Reg. No. 6735-89

DANGER-POISON-Precautions

Precautionary Statements
Hazards To Humans
(& Domestic Animals)
DANGER

Environmental Hazards
Poisonous To Fish And Wildlife

Physical or Chemical Hazards
Use Only When Wearing The Following Protective Equipment And Clothing

Work Safety Rules

POISON - POISON

Poison Signs (Symptoms)

First Aid Treatment

INFORMATION FOR PHYSICIAN

METHYL PARATHION MIXTURE, LIQUID NA2783

DIRECTIONS FOR USE

RE-ENTRY

STORAGE AND DISPOSAL

PROHIBITIONS

PESTICIDE DISPOSAL

CONTAINER DISPOSAL

METAL

CLASS A

RECOMMENDATIONS

CEREAL

BARLEY, OATS, RYE AND WHEAT

CORN

RICE

FIELD AND FORAGE CROPS

ALFALFA AND CLOVER

COTTON

GRASS Forage

HOPS

SOYBEANS

SUGAR BEETS

TOBACCO

CAUTION

VETCH

FRUIT

APPLY AT THE RATES INDICATED FOR THE FOLLOWING CROPS. MAKE NO APPLICATION CLOSER THAN 15 DAYS BEFORE HARVEST. APPLY AS FREQUENTLY AS NEEDED TO CONTROL INSECTS.

APPLES

GRAPES

PEACHES, PLUMS AND PRUNES

STRAWBERRIES

OIL SEED CROPS

SAFFLOWER

SUNFLOWER

VEGETABLES

APPLY AT THE RATES INDICATED FOR THE FOLLOWING CROPS. MAKE NO APPLICATIONS CLOSER THAN 15 DAYS BEFORE HARVEST.

POTATOES

SWEET POTATOES

APPLY AT THE RATES INDICATED FOR THE FOLLOWING CROPS. MAKE NO APPLICATIONS CLOSER THAN 7 DAYS BEFORE HARVEST.

ARTICHOKE

BROCCOLI

BRUSSELS SPROUTS

CAULIFLOWER

KOHLRABI

RUTABAGAS

CABBAGE

COLLARDS, KALE AND MUSTARD

Contains
ethyl Parathion
per Gallon

No. 6735-TX-10

g. No. 6735-89

ISON
ALL CASES OF
N
H CES AND
LOSS

ptoms)

ent

HYSICIAN
PERVISION OF THD
ON

RQ METHYL PARATHION MIXTURE, LIQUID NA2783

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

NOTE: Prepare only enough spray mixture for immediate use. Do not contaminate water supplies by cleaning of equipment or disposal of wastes.

SUGGESTED WATER RATES FOR AIRCRAFT AND GROUND APPLICATION

GAL/WATER/ACRE

SUFFICIENT WATER FOR COVERAGE

RE-ENTRY

Workers must not enter treated areas

STORAGE AND DISPOSAL

PROHIBITIONS

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL

Pesticide wastes are toxic.

CONTAINER DISPOSAL

METAL Nonrefillable container.

Triple rinse (or equivalent).

Offer for recycling or reconditioning.

GLASS

Do not reuse container.

RECOMMENDATIONS

CEREAL

BARLEY OATS RYE AND WHEAT

CORN

RICE

FIELD AND FORAGE CROPS

ALFALFA AND CLOVER

COTTON

CAUTION

GRASS forage

HOPS

SOYBEANS

SUGAR BEETS

TOBACCO

CAUTION

VETCH

FRUIT

APPLY AT THE RATES INDICATED FOR THE FOLLOWING CROPS. MAKE NO APPLICATIONS CLOSER THAN 21 DAYS BEFORE HARVEST. APPLY AS FREQUENTLY AS NEEDED TO CONTROL INSECTS.

APPLES

GRAPES

PEACHES PLUMS AND PRUNES

STRAWBERRIES

OIL SEED CROPS

SAFFLOWER

SUNFLOWER

VEGETABLES

APPLY AT THE RATES INDICATED FOR THE FOLLOWING CROPS. MAKE NO APPLICATIONS CLOSER THAN DAYS BEFORE HARVEST

POTATOES

SWEET POTATOES

APPLY AT THE RATES INDICATED FOR THE FOLLOWING CROPS. MAKE NO APPLICATIONS CLOSER THAN DAYS BEFORE HARVEST

ARTICHOKE

BROCCOLI

BRUSSELS SPROUTS

CAULIFLOWER

KOHLRABI

RUTABAGAS

APPLY AT THE RATES INDICATED FOR THE FOLLOWING CROPS. MAKE NO APPLICATIONS CLOSER THAN DAYS BEFORE HARVEST

CABBAGE

COLLARDS KALE AND MUSTARD

BEANS Dry

BEANS Dry and Green

BEETS Red

CARROTS

CELERY

CUCUMBERS

ONIONS

PEPPERS

SPINACH

TURNIPS

APPLY AT THE RATES INDICATED FOR THE FOLLOWING CROPS. MAKE NO APPLICATIONS CLOSER THAN DAYS BEFORE HARVEST

LETTUCE

TANK MIXTURES

TANK MIXTURE	INSECT	RATE/ACRE
METHYL PARATHION		
METHYL PARATHION		
METHYL PARATHION		
METHYL PARATHION		
METHYL PARATHION		
METHYL PARATHION		
METHYL PARATHION		
METHYL PARATHION		
METHYL PARATHION		
METHYL PARATHION		

NET CONTENTS GALS

A PRODUCT OF

TIDE PRODUCTS, INC.

EDINBURG, TEXAS 78539

ONLY the BEST . . . is fit to be TIDE