

# RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.

BEST DOCUMENT AVAILABLE

# ETHYL

## Emulsifiable Concentrate Insecticide

STOP — READ LABEL BEFORE USING!


NOT FOR HOME USE

### POISON



### DANGER

Keep Out  
Reach of  
Children



Contains  
4 lbs. Parathion  
per Gallon

E.P.A. Est. No. 6735-TX-10

E.P.A. Reg. No. 6735-100

#### ACTIVE INGREDIENTS:

- \*Parathion (O, O-diethyl O-p-1-trophenyl phosphorothioate) 45.07%
- \*Related Compounds from Parathion 2.16%
- Xylene Range Aromatic Solvent 48.95%
- INERT INGREDIENTS 1.72%

\*Also known as Ethyl Parathion

## DANGER-POISON-CAUTIONS



**POISONOUS IF SWALLOWED**  
This product can cause death if swallowed. Avoid contact with mouth. Do not drink from container. Do not use as a food preservative.



**POISONOUS BY SKIN CONTACT**  
This product can cause death if absorbed through the skin. Do not touch. Do not use as a food preservative.



**DO NOT BREATHE**  
This product can cause death if inhaled. Do not breathe. Do not use as a food preservative.

### Precautionary Statements Hazards To Humans (& Domestic Animals) DANGER

This product can cause death if swallowed. Avoid contact with mouth. Do not drink from container. Do not use as a food preservative. Do not touch. Do not breathe. Do not use as a food preservative.

### Environmental Hazards Poisonous To Fish And Wildlife

This product is highly toxic to fish and wildlife. Do not apply to water. Do not use in or near water. Do not use as a food preservative.

### Physical or Chemical Hazards Use Only When Wearing The Following Protective Equipment And Clothing

Wear protective clothing and equipment. Do not eat, drink, or smoke while using. Do not use as a food preservative.

### Work Safety Rules

Do not use in or near water. Do not use as a food preservative. Do not touch. Do not breathe. Do not use as a food preservative.

### POISON POISON CALL A PHYSICIAN IMMEDIATELY IN ALL CASES OF SUSPECTED POISONING. FOLLOW PROCEDURES, ANTIDOTES AND PHYSICIAN'S INSTRUCTIONS.

### Poison Signs (Symptoms)

**WARNING SYMPTOMS:** ...  
**First Aid Treatment:** ...  
**IF SWALLOWED:** ...  
**IF ON SKIN:** ...  
**IF IN EYES:** ...  
**IF INHALED:** ...

### INFORMATION FOR PHYSICIAN

This product is highly toxic. Do not use in or near water. Do not use as a food preservative. Do not touch. Do not breathe. Do not use as a food preservative.

# RQ PARATHION MIXTURE, LIQUID

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**NOTE:** Preparation of nurse tank emulsion for transfer of smaller amounts to application rig requires agitation to prevent emulsion settling.

**SUGGESTED WATER RATES FOR AIRCRAFT AND GROUND APPLICATION:** (The actual side through uniform coverage varies with plant growth at time of application. Except as specified, following rates are therefore intended to cover a broad range of conditions.)  
CROP

Vegetable and Field Crops  
FOR USE ONLY IN POWER SPRAYERS BY OPERATORS EXPERIENCED OR ADEQUATE IN THE APPLICATION OF PARATHION INSECTICIDES.

## RE-ENTRY

Workers entering treated fields within 48 hours must wear protective clothing.

## STORAGE AND DISPOSAL

- PROHIBITIONS:** Do not contaminate water, food, feed, by storage or disposal. Open dumping is prohibited.
- PESTICIDE DISPOSAL:** Pesticide spray mixture or rinse that cannot be used according to label instructions must comply with Federal, State or local procedures under the Resource Conservation and Recovery Act.
- CONTAINER DISPOSAL:** Triple rinse or equivalent and offer for recycling or reconditioning, or dispose of in a sanitary approved State and local procedures. Triple rinse or equivalent and dispose of in a sanitary landfill or by other approved State or local procedures.

## RECOMMENDATIONS

### CEREAL

**DO NOT APPLY WITHIN 22 DAYS OF HARVEST, CUTTING OR FORAGE USE ON THE FOLLOWING CROPS:** For control of European corn borers, use 1 pint per acre. Apply the first spray when silks are 1/2 to 1 inch long and follow with two additional sprays at 7 to 10 day intervals. Use sufficient water to provide coverage and be certain wheels of plant are well treated. For corn leaf aphids use 1/2 pint per acre. For control of European corn borers, armyworms up to third instar, climbing cutworms, and European spruce sawfly, use 1/2 pint per acre. To control stalk borer and spider beetles, use 1/2 pint per acre.  
**SORGHUM:** For control of European corn borers, apply at rate of 1 pint to 1 quart per acre. 2 applications, approximately 10 to 14 days apart, are preferred. For control of European corn borers, apply at rate of 1/2 pint per acre. For control of armyworms, fall armyworms, and hairy worms, use 1/2 pint per acre. For control of European spruce sawfly, use 1/2 pint per acre. For control of stalk borer, use 1/2 pint per acre. Spray a few rows a week or so before heading to test effects on plants.

**DO NOT APPLY WITHIN 22 DAYS OF HARVEST, CUTTING OR FORAGE USE ON THE FOLLOWING CROPS:** For control of European corn borers, use 1/2 pint per acre. For control of armyworms, fall armyworms, and hairy worms, use 1/2 pint per acre. For control of European spruce sawfly, use 1/2 pint per acre. For control of stalk borer, use 1/2 pint per acre.

### FIELD AND FORAGE CROPS

**DO NOT APPLY WITHIN 22 DAYS OF HARVEST, CUTTING OR FORAGE USE ON THE FOLLOWING CROPS:** For control of European corn borers, use 1/2 pint per acre. For control of armyworms, fall armyworms, and hairy worms, use 1/2 pint per acre. For control of European spruce sawfly, use 1/2 pint per acre. For control of stalk borer, use 1/2 pint per acre.

**DO NOT APPLY WITHIN 22 DAYS OF HARVEST, CUTTING OR FORAGE USE ON THE FOLLOWING CROPS:** For control of European corn borers, use 1/2 pint per acre. For control of armyworms, fall armyworms, and hairy worms, use 1/2 pint per acre. For control of European spruce sawfly, use 1/2 pint per acre. For control of stalk borer, use 1/2 pint per acre.

**DO NOT APPLY WITHIN 22 DAYS OF HARVEST, CUTTING OR FORAGE USE ON THE FOLLOWING CROPS:** For control of European corn borers, use 1/2 pint per acre. For control of armyworms, fall armyworms, and hairy worms, use 1/2 pint per acre. For control of European spruce sawfly, use 1/2 pint per acre. For control of stalk borer, use 1/2 pint per acre.

**DO NOT APPLY WITHIN 22 DAYS OF HARVEST, CUTTING OR FORAGE USE ON THE FOLLOWING CROPS:** For control of European corn borers, use 1/2 pint per acre. For control of armyworms, fall armyworms, and hairy worms, use 1/2 pint per acre. For control of European spruce sawfly, use 1/2 pint per acre. For control of stalk borer, use 1/2 pint per acre.

### FRUIT

**DO NOT APPLY WITHIN 22 DAYS OF HARVEST, CUTTING OR FORAGE USE ON THE FOLLOWING CROPS:** For control of European corn borers, use 1/2 pint per acre. For control of armyworms, fall armyworms, and hairy worms, use 1/2 pint per acre. For control of European spruce sawfly, use 1/2 pint per acre. For control of stalk borer, use 1/2 pint per acre.

**DO NOT APPLY WITHIN 22 DAYS OF HARVEST, CUTTING OR FORAGE USE ON THE FOLLOWING CROPS:** For control of European corn borers, use 1/2 pint per acre. For control of armyworms, fall armyworms, and hairy worms, use 1/2 pint per acre. For control of European spruce sawfly, use 1/2 pint per acre. For control of stalk borer, use 1/2 pint per acre.

**DO NOT APPLY WITHIN 22 DAYS OF HARVEST, CUTTING OR FORAGE USE ON THE FOLLOWING CROPS:** For control of European corn borers, use 1/2 pint per acre. For control of armyworms, fall armyworms, and hairy worms, use 1/2 pint per acre. For control of European spruce sawfly, use 1/2 pint per acre. For control of stalk borer, use 1/2 pint per acre.

# RQ PARATHION MIXTURE, LIQUID NA2783

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

NOTE: Preparation of nurse tank emulsion for transfer of pesticide to its application rate should provide sufficient agitation to prevent emulsion settling.

**SUGGESTED WATER RATES FOR AIRCRAFT AND GROUND APPLICATION:** The actual rate required will vary through uniform coverage varies with plant growth and rate of application. Except as specified, the following rates are therefore intended to cover a broad range of conditions.

CROP	GAL Water/Acre	Aircraft Ground
Vegetable and Field Crops	10-20	10-20

FOR USE ONLY IN POWER STRAYERS BY OPERATORS EXPERIENCED OR ADEQUATELY TRAINED IN THE APPLICATION OF PARATHION INSECTICIDES.

## RE-ENTRY

Workers entering treated field within 4 hours must wear protective clothing.

## STORAGE AND DISPOSAL

- PROHIBITIONS:** Do not contaminate water, food, feed, storage or disposal. Open burning is prohibited. Do not reuse empty container.
- PESTICIDE DISPOSAL:** Pesticide spray mixture or residue that cannot be used according to label instructions must be disposed of according to Federal, State or local procedures under the Hazardous Conservation and Recovery Act.
- CONTAINER DISPOSAL:** METAL: Empty container or equivalent and offer for recycling or reuse, for reuse or disposal in a sanitary landfill or by other approved State and local procedures. GLASS: Empty container or equivalent and dispose of in a sanitary landfill or by other approved State and local procedures.

## RECOMMENDATIONS

### CEREAL

**DO NOT APPLY WITHIN 30 DAYS OF HARVEST ON THE FOLLOWING CROPS:**  
**CORN:** For control of European cornborer, use 1.2 pint per acre. Apply the first spray when 75% of the corn plants show silks and the second spray when silks are 2/3 grown. For control of fall armyworm, use 1.2 pint per acre. For control of stalk borer, use 1.2 pint per acre. For control of stalk borer, use 1.2 pint per acre. For control of stalk borer, use 1.2 pint per acre.

### FIELD AND FORAGE CROPS

**DO NOT APPLY WITHIN 30 DAYS OF HARVEST ON THE FOLLOWING CROPS:**  
**SOYBEANS:** For control of soybean aphid, use 1.2 pint per acre. For control of soybean aphid, use 1.2 pint per acre. For control of soybean aphid, use 1.2 pint per acre. For control of soybean aphid, use 1.2 pint per acre.

### FRUIT

**DO NOT APPLY WITHIN 30 DAYS OF HARVEST ON THE FOLLOWING CROPS:**  
**APPLES:** For control of apple maggot, use 1.2 pint per acre. For control of apple maggot, use 1.2 pint per acre. For control of apple maggot, use 1.2 pint per acre. For control of apple maggot, use 1.2 pint per acre.

**DO NOT APPLY WITHIN 30 DAYS OF HARVEST ON THE FOLLOWING CROPS:**  
**PEACHES AND NECTARINES:** For control of green peach aphid, use 1.2 pint per acre. For control of green peach aphid, use 1.2 pint per acre. For control of green peach aphid, use 1.2 pint per acre. For control of green peach aphid, use 1.2 pint per acre.

**DO NOT APPLY WITHIN 30 DAYS OF HARVEST ON THE FOLLOWING CROPS:**  
**PEAS:** For control of pea beetle and leaf miner, use 1.2 pint per acre. For control of pea beetle and leaf miner, use 1.2 pint per acre. For control of pea beetle and leaf miner, use 1.2 pint per acre. For control of pea beetle and leaf miner, use 1.2 pint per acre.

**DO NOT APPLY WITHIN 30 DAYS OF HARVEST ON THE FOLLOWING CROPS:**  
**VEGETABLES:** For control of aphid, leaf miner, and armyworm, use 1.2 pint per acre. For control of aphid, leaf miner, and armyworm, use 1.2 pint per acre. For control of aphid, leaf miner, and armyworm, use 1.2 pint per acre. For control of aphid, leaf miner, and armyworm, use 1.2 pint per acre.

**DO NOT APPLY WITHIN 30 DAYS OF HARVEST ON THE FOLLOWING CROPS:**  
**GRAPES:** For control of grape phylloxera, use 1.2 pint per acre. For control of grape phylloxera, use 1.2 pint per acre. For control of grape phylloxera, use 1.2 pint per acre. For control of grape phylloxera, use 1.2 pint per acre.

**DO NOT APPLY WITHIN 30 DAYS OF HARVEST ON THE FOLLOWING CROPS:**  
**WATERMELONS:** For control of watermelon beetle and leaf miner, use 1.2 pint per acre. For control of watermelon beetle and leaf miner, use 1.2 pint per acre. For control of watermelon beetle and leaf miner, use 1.2 pint per acre. For control of watermelon beetle and leaf miner, use 1.2 pint per acre.

**DO NOT APPLY WITHIN 30 DAYS OF HARVEST ON THE FOLLOWING CROPS:**  
**PEACHES AND NECTARINES:** For control of green peach aphid, use 1.2 pint per acre. For control of green peach aphid, use 1.2 pint per acre. For control of green peach aphid, use 1.2 pint per acre. For control of green peach aphid, use 1.2 pint per acre.

**PEACHES AND NECTARINES (Areas other than California):** For control of green peach aphid, use 1.2 pint per acre. For control of green peach aphid, use 1.2 pint per acre. For control of green peach aphid, use 1.2 pint per acre. For control of green peach aphid, use 1.2 pint per acre.

### NUTS

**DO NOT APPLY WITHIN 30 DAYS OF HARVEST ON THE FOLLOWING CROPS UNLESS OTHERWISE SPECIFIED:**  
**ALMONDS:** For control of almond moth, use 1.2 pint per acre. For control of almond moth, use 1.2 pint per acre. For control of almond moth, use 1.2 pint per acre. For control of almond moth, use 1.2 pint per acre.

### VEGETABLES

**DO NOT APPLY WITHIN 30 DAYS OF HARVEST ON THE FOLLOWING CROPS UNLESS OTHERWISE SPECIFIED:**  
**CABBAGE:** For control of cabbage looper, use 1.2 pint per acre. For control of cabbage looper, use 1.2 pint per acre. For control of cabbage looper, use 1.2 pint per acre. For control of cabbage looper, use 1.2 pint per acre.

**DO NOT APPLY WITHIN 30 DAYS OF HARVEST ON THE FOLLOWING CROPS UNLESS OTHERWISE SPECIFIED:**  
**BEANS:** For control of bean leaf beetle and two-spotted mite, use 1.2 pint per acre. For control of bean leaf beetle and two-spotted mite, use 1.2 pint per acre. For control of bean leaf beetle and two-spotted mite, use 1.2 pint per acre. For control of bean leaf beetle and two-spotted mite, use 1.2 pint per acre.

**DO NOT APPLY WITHIN 30 DAYS OF HARVEST ON THE FOLLOWING CROPS UNLESS OTHERWISE SPECIFIED:**  
**BEETS:** For control of flea beetle and leaf miner, use 1.2 pint per acre. For control of flea beetle and leaf miner, use 1.2 pint per acre. For control of flea beetle and leaf miner, use 1.2 pint per acre. For control of flea beetle and leaf miner, use 1.2 pint per acre.

**DO NOT APPLY WITHIN 30 DAYS OF HARVEST ON THE FOLLOWING CROPS UNLESS OTHERWISE SPECIFIED:**  
**CUCUMBERS:** For control of squash vine borer, aphid, and leaf miner, use 1.2 pint per acre. For control of squash vine borer, aphid, and leaf miner, use 1.2 pint per acre. For control of squash vine borer, aphid, and leaf miner, use 1.2 pint per acre. For control of squash vine borer, aphid, and leaf miner, use 1.2 pint per acre.

**DO NOT APPLY WITHIN 30 DAYS OF HARVEST ON THE FOLLOWING CROPS UNLESS OTHERWISE SPECIFIED:**  
**GARLIC:** For control of leaf miner, use 1.2 pint per acre. For control of leaf miner, use 1.2 pint per acre. For control of leaf miner, use 1.2 pint per acre. For control of leaf miner, use 1.2 pint per acre.

**DO NOT APPLY WITHIN 30 DAYS OF HARVEST ON THE FOLLOWING CROPS UNLESS OTHERWISE SPECIFIED:**  
**ONIONS:** For control of onion thrips, use 1.2 pint per acre. For control of onion thrips, use 1.2 pint per acre. For control of onion thrips, use 1.2 pint per acre. For control of onion thrips, use 1.2 pint per acre.

**DO NOT APPLY WITHIN 30 DAYS OF HARVEST ON THE FOLLOWING CROPS UNLESS OTHERWISE SPECIFIED:**  
**PEPPERS:** For control of pepper weevil, use 1.2 pint per acre. For control of pepper weevil, use 1.2 pint per acre. For control of pepper weevil, use 1.2 pint per acre. For control of pepper weevil, use 1.2 pint per acre.

**DO NOT APPLY WITHIN 30 DAYS OF HARVEST ON THE FOLLOWING CROPS UNLESS OTHERWISE SPECIFIED:**  
**RADISHES:** For control of radish root fly, use 1.2 pint per acre. For control of radish root fly, use 1.2 pint per acre. For control of radish root fly, use 1.2 pint per acre. For control of radish root fly, use 1.2 pint per acre.

**DO NOT APPLY WITHIN 30 DAYS OF HARVEST ON THE FOLLOWING CROPS UNLESS OTHERWISE SPECIFIED:**  
**SPINACH:** For control of aphid, leaf miner, and armyworm, use 1.2 pint per acre. For control of aphid, leaf miner, and armyworm, use 1.2 pint per acre. For control of aphid, leaf miner, and armyworm, use 1.2 pint per acre. For control of aphid, leaf miner, and armyworm, use 1.2 pint per acre.

**DO NOT APPLY WITHIN 30 DAYS OF HARVEST ON THE FOLLOWING CROPS UNLESS OTHERWISE SPECIFIED:**  
**SQUASH:** For control of squash vine borer, use 1.2 pint per acre. For control of squash vine borer, use 1.2 pint per acre. For control of squash vine borer, use 1.2 pint per acre. For control of squash vine borer, use 1.2 pint per acre.

**DO NOT APPLY WITHIN 30 DAYS OF HARVEST ON THE FOLLOWING CROPS UNLESS OTHERWISE SPECIFIED:**  
**SWEET POTATOES:** For control of sweet potato weevil, use 1.2 pint per acre. For control of sweet potato weevil, use 1.2 pint per acre. For control of sweet potato weevil, use 1.2 pint per acre. For control of sweet potato weevil, use 1.2 pint per acre.

**DO NOT APPLY WITHIN 30 DAYS OF HARVEST ON THE FOLLOWING CROPS UNLESS OTHERWISE SPECIFIED:**  
**WATERMELONS:** For control of watermelon beetle and leaf miner, use 1.2 pint per acre. For control of watermelon beetle and leaf miner, use 1.2 pint per acre. For control of watermelon beetle and leaf miner, use 1.2 pint per acre. For control of watermelon beetle and leaf miner, use 1.2 pint per acre.

**DO NOT APPLY WITHIN 30 DAYS OF HARVEST ON THE FOLLOWING CROPS UNLESS OTHERWISE SPECIFIED:**  
**PEACHES AND NECTARINES:** For control of green peach aphid, use 1.2 pint per acre. For control of green peach aphid, use 1.2 pint per acre. For control of green peach aphid, use 1.2 pint per acre. For control of green peach aphid, use 1.2 pint per acre.

NET CONTENTS \_\_\_\_\_ GALS

A PRODUCT OF  
**TIDE PRODUCTS, INC.**  
EDINBURG, TEXAS 78539

ONLY the BEST . . . is fit to be TIDE