RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators or persons under their direct supervision, and only for uses covered by the Certified Applicator's certification.

PARATHION 8-E

Emulsifiable Concentrate Insecticide

ACCEPTED
OCT 2 8 1981
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No.

PROTECT FROM TEMPERATURES BELOW 32° F

STOP — READ LABEL BEFORE USING!

NOT FOR HOME USE

POISON

DANGER

Keep Out of Reach of Children

BEST DOCUMENT AVAILABLE

ACTIVE INGREDIENTS:

*Parathion (O, O-diethyl) O-p-nitrophenyl phosphorothioate	77.21%
*Related Compounds from Parathion	3.70%
Xylene Range Aromatic Solvent	9.86%
INERT INGREDIENTS	9.23%
	100.00%

**Also known as Ethyl Parathion

Contains

8 Lbs. Parathion

Per Gallon

E.P.A. Est. No. 6735-TX-10

E.P.A. Reg. No. 6735-113

DANGER-POISON-PRECAUTIONS



DO NOT SWALLOW

POISONOUS if SWALLOWED

A product can kill you if swallowed even in small amounts. Spray mist or dust may be fatal if swallowed.



POISONOUS by CONTACT

Poisonous if touched by hands or spilled or splashed on skin, in eyes or on clothing. Liquid goes through clothes.

Breathing vapors, spray mist or dust may be fatal.

Precautionary Statements
Hazards To Humans (& Domestic Animals)
DANGER

Poisonous if swallowed, inhaled or absorbed through the skin. Rapidly absorbed through the skin or eyes.

Do not get in eyes, on skin or on clothing. Do not breathe the spray mist or vapor.

Repeated inhalation or skin contact may, without symptoms, progressively increase susceptibility to poisoning.

Clothing wet with this material must be removed immediately.

Exposed persons must receive prompt medical attention or they may die.

Do not contaminate feed and foodstuffs.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

Environmental Hazards
Poisonous To Fish And Wildlife

This product is toxic to fish, birds and other wildlife. Birds and other wildlife in treated areas may be killed. Do not apply directly to water. Do not apply when weather conditions favor drift from treated areas. Do not apply where run-off is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. APPLY THIS PRODUCT ONLY AS SPECIFIED ON THIS LABEL.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Physical or Chemical Hazards
Use Only When Wearing The Following Protective Equipment And Clothing

Wear clean, heavy duty, natural rubber gloves, waterproof jacket, hat, rubber boots or rubber overshoes and safety goggles when handling, mixing or applying this product. Wear a pesticide respirator jointly approved by the Mine Safety Enforcement and Safety Administration (formerly the U.S. Bureau of Mines) and by the National Institute for Occupational Safety and Health under provisions of 30 CFR Part 11.

Work Safety Rules

Keep all unprotected persons and children away from treated area where there is danger of drift.

Do not rub eyes or mouth with hands. If you feel symptoms any way STOP work and get help right away. Call a doctor, physician, clinic, hospital immediately. Explain that the victim has been exposed to Parathion 8-E and describe his condition. After first aid is given see First Aid Treatment Section and if doctor cannot come, take victim to clinic or hospital.

IMPORTANT: Before removing gloves, wash them with soap and water. Always wash hands, face and arms with soap and water before smoking, eating or drinking.

AFTER WORK, take off all work clothes and shoes. Shower using soap and water. Wear only clean clothes when leaving work. Do not wear contaminated clothing. Wash protective clothing and protective equipment with soap and water after each use. Respirator should be cleaned and filter replaced according to instructions included with respirator.

POISON — POISON

CALL A PHYSICIAN IMMEDIATELY IN ALL CASES OF SUSPECTED POISONING

FIRST AID PROCEDURES, ANTIDOTES AND PHYSICIAN'S INSTRUCTIONS

Poison Signs (Symptoms)

WARNING SYMPTOMS include weakness, headache, tightness in the chest, altered respiration, blurred vision, nausea, abdominal cramps, diarrhea, sweating, vomiting, salivation and constricted pinpoint pupils.

First Aid Treatment

IF SWALLOWED: Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

INHALATION: Remove victim from contaminated atmosphere.

SKIN: In case of contact, immediately remove patient from the vicinity of the pesticide. Remove contaminated clothing and immediately wash skin thoroughly with soap and water.

EYES: In case of contact, immediately remove victim from the vicinity of the pesticide. Flush eyes immediately with water.

IF WARNING SYMPTOMS (See Above) OCCUR: Keep victim quiet and quiet. Start artificial respiration if victim stops breathing. Transport victim to nearest hospital immediately.

ANTIDOTE: Atropine is the specific antidote for Parathion poisoning. Give an adult 2 mg (1/30 gr) to 1.5 gram of atropine, immediately, preferably by intravenous injection, and repeat at 5 to 10 minute intervals until signs of atropinization appear: dry, flushed skin, dilated pupils, rapid pulse. Maintain full atropinization and give 1/2 mg (adult dosage) Protopam (2-PAM) intravenously in saline solution to restore normal cholinesterase levels. DO NOT GIVE MORPHINE OR TRANQUILIZERS. Methyl Parathion is a strong cholinesterase inhibitor affecting the central and peripheral nervous systems and producing cardiac and respiratory depression. At the first signs of pulmonary edema, the patient should be given supplementary oxygen and treated symptomatically. Convulsions, if the patient may survive, and fatal relapses have been reported after initial improvement. VERY CLOSE SUPERVISION OF THE PATIENT IS INDICATED FOR AT LEAST 48 HOURS.

Clear airways by postural drainage. Artificial respiration, oxygen, and other measures may be necessary.

Repeated exposure to cholinesterase inhibitors [illegible]

INFORMATION FOR PHYSICIAN

[illegible]

RQ PARATHION MIXTURE, LIQUID N

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

NOTE: Preparation of nurse tank emulsion (for transfer of smaller amounts to application rigs) should [illegible] sufficient agitation to prevent emulsion settling.

SUGGESTED WATER RATES FOR AIRCRAFT AND GROUND APPLICATION: (The actual [illegible] provide thorough uniform coverage varies with plant growth at time of application. Except as specified [illegible] following rates are therefore intended to cover a broad range of growing conditions.)

CROP

Vegetables and Field Crops

BEST DOCUMENT A

RE-ENTRY

(Workers entering treated fields within 48 hours must wear protective clothing.)

STORAGE AND DISPOSAL

1. PROHIBITIONS
Do not contaminate water, food, feed by storage or disposal. Open dumping is prohibited. Do [illegible] container.

2. PESTICIDE DISPOSAL
Pesticide spray mixture or rinsate that cannot be used according to label instructions must be [illegible] according to Federal, State or local procedures under the Resource Conservation and Recovery Act.

3. CONTAINER DISPOSAL
METAL (All Sizes)
Triple rinse (or equivalent) and offer for recycling or reconditioning, or dispose of in a sanitary [illegible] approved State and local procedures.

RECOMMENDATIONS

CEREAL

DO NOT APPLY WITHIN 12 DAYS OF HARVEST, CUTTING OR FORAGE USE ON THE FO[illegible]

CORN: For control of European corn borers, use 1/2 pint per acre. Apply the first spray when 7 [illegible] show "shot hole" feeding and follow with two additional sprays at 5 to 7 day intervals. Use sufficient [illegible] provide complete coverage and be certain whorls of plant are well treated. For corn leaf aphids, use [illegible] fall armyworms up to third instar, chinbug, cutworms, grasshoppers and Japanese beetles, use 6 ounces [illegible] control stink bugs and spider mites, use 1/2 pint per acre. To control chinch bugs, use 3/4 pint per a[illegible]

SORGHUM: To control sorghum midge, apply at rate of [illegible] pint to 1 pint per acre, 2 applications [illegible] approximately [illegible] of the heads have completely emerged from the boot or not later than start of bl[illegible] aphids and mites, use [illegible] pint per acre. For sorghum webworms, fall armyworms up to third instar, use 6 to 8 ounces per acre. To control chinch bugs, use 1/2 pint per acre. Leaf injury may occur on so[illegible] sorghums. Spray a few rows a week or so before beginning to test effects on plants.

DO NOT APPLY WITHIN 1 DAY OF HARVEST, CUTTING OR FORAGE USE ON THE FO[illegible]

SMALL GRAINS (WHEAT, OATS, BARLEY): To control armyworms up to third instar, a[illegible] winter grain rates use [illegible] pint per acre. For thrips, use 4 to 6 ounces per acre. For Sax's plant bugs [illegible] For black grass bugs, stink bugs, white spider mites, leafhoppers, chinbing cutworms, grasshopp[illegible] mites, use [illegible] pint per acre. For [illegible] bugs, false chinch bugs and banks grass mites, use 1/2 p[illegible]

FIELD AND FORAGE CROPS

DO NOT APPLY WITHIN [illegible] DAYS OF HARVEST, CUTTING OR FORAGE USE ON THE F[illegible]

SOYBEANS: For control of [illegible] pint per acre. To control velvet bean caterpillars, gre[illegible] spotted mites and stink bugs, use [illegible] pint per acre. To control corn earworms and fall armyworms [illegible] acre. To control white grubs and wireworms, broadcast [illegible] gallon per acre just prior to planting [illegible] potato and pepper [illegible]

DO NOT APPLY WITHIN [illegible] DAYS OF HARVEST, CUTTING OR FORAGE USE ON THE F[illegible]

ALFALFA, CLOVER, VETCH AND GRASS: For sweet clover aphids, three-cornered alfalfa [illegible] pilloses and spittlebugs, use [illegible] pint per acre. For aphids, alfalfa weevil larvae and adult weevils a[illegible] webworms, grasshoppers, crickets, spotted alfalfa [illegible] bugs, chinch bugs and [illegible] to [illegible] pint per acre. For alfalfa seed chalcids, conte[illegible] and Nevada regulations [illegible] This material [illegible] per acre. For [illegible] mites, blister beetles, Asiatic garden beetles, sweet clo[illegible] beetles, use [illegible] pint per acre. For lesser armyworms and cutworms, use [illegible] to [illegible] pint per acre. For [illegible] bugs, false chinch bugs and banks grass mites, use [illegible] p[illegible]

PEANUTS: [illegible] pint per acre. For [illegible] bugs, use [illegible] pint per acre. For [illegible] spray to soil surface and base of plants.

SUGAR BEETS: [illegible] aphids and [illegible] to third instar, leafhoppers, blister [illegible] grasshoppers, use [illegible] pint per acre. For [illegible] bugs, use [illegible] pint per acre. For [illegible] ground applications [illegible] stage. For [illegible] gallon per acre just prior to p[illegible]

SUGARCANE: [illegible]

DO NOT APPLY WITHIN [illegible] DAYS OF HARVEST ON THE FOLLOWING CROPS:

COTTON: [illegible] flea hoppers, garden webworms [illegible] pint per acre. For cabbage looper, use [illegible] pint per acre. For [illegible] caterpillars, use [illegible] to 1 pint per acre. For [illegible] bugs, sorghum leaf miners and Southern garden [illegible] per acre. Use [illegible] water [illegible] coverage. Make first application when insects appear at [illegible] frequent [illegible] treated with the insecticide are complete [illegible]

DO NOT APPLY WITHIN [illegible] DAYS OF HARVEST ON THE FOLLOWING CROPS:

TOBACCO: [illegible]

VEGTABLES

DO NOT APPLY WITHIN [illegible] DAYS OF HARVEST ON THE FOLLOWING CROPS [illegible] SPECIFIED:

CELERY: [illegible] pint per acre. [illegible] with no days of harvest.

ENDIVE: [illegible] per acre.

LETTUCE (LEAF AND BIBB): [illegible] cabbage loopers [illegible] spotted leaf [illegible] per acre. At the [illegible] pint rate harves[illegible]

Contains

ds. Parathion

Per Gallon

. 6735-TX-10

No. 6735-113

T BREATHE

vapors spray mist or

ds fatal

SON

ALL CASES OF

ES AND

NS

oms)

e tightness in the chest

ral cramps, diarrhea

ist pupils

nt

d induce vomiting in

ing or give anything by

n

atmosphere

t from the vicinity of

k and immediately wash

n from the vicinity of the

espiration if victim is not

l immediately

on poisoning. Obtain

r physician. Keep in

YSICIAN

ately, preferably by in

tervals until signs of

rapid pulse. Maintain

2 PAM intravenous

levels. DO NOT GIVE

Parathion is a strong

ral nervous systems

At the first sign of

mentary oxygen and

poison may occur and

ment. VERY CLOSE

FOR AT LEAST 48

oxygen administra

chest warming causes

cholinesterase inhibitor

has been administered as

RQ PARATHION MIXTURE, LIQUID NA2783

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

NOTE: Preparation of nurse tank emulsion (for transfer of smaller amounts to application rigs) should provide for efficient agitation to prevent emulsion settling

SUGGESTED WATER RATES FOR AIRCRAFT AND GROUND APPLICATION [illegible] actual rates required to provide thorough, uniform coverage vary with plant size, [illegible] the following rates are therefore intended to cover a broad range of growing conditions.

CROP	GAL Water Acre Aircraft Ground
Vegetables and Field Crops	[illegible]

BEST [illegible] AVAILABLE

RE-ENTRY

(Workers entering treated [illegible] must wait [illegible])

STORAGE AND DISPOSAL

1. **PROHIBITIONS**
Do not contaminate water, food, feed by storage or disposal. Do not reuse empty containers.

2. **PESTICIDE DISPOSAL**
Pesticide spray mixtures: [illegible] according to Federal, State or Local procedures under the Resource Conservation and Recovery Act.

3. **CONTAINER DISPOSAL**
METAL (All Sizes)
Triple rinse for equivalent and offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by other approved State and Local procedures.

RECOMMENDATIONS

CEREAL

DO NOT APPLY WITHIN 21 DAYS OF HARVEST CUTTING OR FORAGE USE ON THE FOLLOWING CROPS

CORN – For control of European corn borers, use 1 pint per acre. Apply the first spray when 5% of the corn plants show shot hole feeding and follow with two additional sprays at 5 to 7 day intervals. Use sufficient water per acre to provide complete coverage and the centers of whorls of plants are well treated. For corn leaf aphids, use 1/4 pint per acre. For fall armyworms up to third instar, chinch bugs, grasshoppers and Japanese beetles, use 6 ounces per acre. For control stink bugs and spider mites, use 1/2 pint per acre. For control chinch bugs, use 1/4 pint per acre.

SORGHUM – To control sorghum midge, apply at rate of 1/4 pint to 1 pint per acre. 2 applications 10-5 days apart when approximately 50% of the heads have completely emerged from the boot or not later than start of blooming. For corn leaf aphids and mites, use 1/4 pint per acre. For sorghum webworms, fall armyworms up to third instar and corn earworms, use 6 to 8 ounces per acre. To control chinch bugs, use 1/4 pint per acre. Leaf injury may occur on some hybrid varieties of sorghum. Spray a few rows as a test before treating to test effects on plants.

DO NOT APPLY WITHIN 7 DAYS OF HARVEST CUTTING OR FORAGE USE ON THE FOLLOWING CROPS

SMALL GRAINS (WHEAT, OATS, BARLEY) – To control armyworms up to third instar, aphids, grasshoppers and winter grain mites, use 1/4 pint per acre. For thrips, use 4 to 6 ounces per acre. For Sax's plant bugs, use 6 ounces per acre. For black grass bugs, stink bugs, wheat spider mites, leafhoppers, chinch bugs, armyworms, grasshoppers and brown wheat mites, use 1/2 pint per acre. For stink bugs, false chinch bugs and banks grass mites, use 3/4 pint per acre.

FIELD AND FORAGE CROPS

DO NOT APPLY WITHIN 21 DAYS OF HARVEST CUTTING OR FORAGE USE ON THE FOLLOWING CROPS

SOYBEANS – To control Mexican bean beetles, use 1/2 pint per acre. To control velvet bean caterpillars, green cloverworms, two-spotted mites and stink bugs, use 1 pint per acre. To control armyworms and fall armyworms, use 6 to 12 ounces per acre. Broadcast application: 1 pint per acre. Cost plant by planting and thoroughly incorporate into upper 4 to 6 inches of soil.

DO NOT APPLY WITHIN 7 DAYS OF HARVEST CUTTING OR FORAGE USE ON THE FOLLOWING CROPS

ALFALFA, CLOVER, VETCH AND GRASS – For sweet clover aphids, three cornered alfalfa hoppers, alfalfa caterpillars and spittlebugs, use 1/2 pint per acre. For aphids, alfalfa weevil larvae and adult weevils, armyworms up to third instar, third instar clover leaf weevils, chinch bugs, armyworms, webworms, grasshoppers, crickets, spotted alfalfa aphids, leafhoppers, lygus bugs, thrips and tortoise beetles, use 1/4 to 1/2 pint per acre. For alfalfa seed [illegible] Canada [illegible] grown for seed, use 1/4 to 1/2 pint per acre. California and Nevada regulations limit [illegible] per acre. For control of armyworms, cutworms, blister beetles, Asiatic garden beetles, sweet clover weevils and green June beetles, use 1/2 pint per acre. For best results, [illegible] 1/2 pint per acre. [illegible] spray legumes during [illegible] per acre.

PEANUTS – To control armyworms and redbanded stink bugs, use 1/2 pint per acre. To control [illegible] hoppers, use 1/2 pint per acre. [illegible] spray to control leafhoppers, disease [illegible] plants.

SUGAR BEETS – [illegible] aphids are [illegible] leaf miners, lygus bugs, stink bugs, webworms, [illegible] and grasshoppers, use 1/2 pint per acre. For false chinch bugs, use 3/4 pint per acre. For best results, use 1/2 pint per acre ground application over the row during seedling stage. To control white grubs and armyworms, broadcast application: 1 pint per acre. Cost plant by planting and thoroughly incorporate into [illegible] inches of soil.

SUGARCANE – [illegible] furrow at time of planting.

DO NOT APPLY WITHIN 30 DAYS OF HARVEST ON THE FOLLOWING CROPS

COTTON – To control [illegible] leafhoppers, garden webworms, [illegible] 1/2 pint per acre. For control spider mites, use 1/2 pint per acre. For cabbage loopers, use 1/2 ounces per acre. For bollworms and stink bugs, use 1/2 pint per acre. For saltmarsh caterpillars, use 1/2 pint per acre. For [illegible] cotton leaf perforator, lygus bugs, false chinch bugs, sugarbeet leaf miners and other garden leafhoppers, use 1 pint per acre. Use enough water for complete coverage. Make first application when pests appear and repeat at 7 day intervals if required. If desired, [illegible] may be tank mixed with [illegible]

DO NOT APPLY WITHIN [illegible] DAYS OF HARVEST ON THE FOLLOWING CROPS

TOBACCO – [illegible]

VEGETABLES

DO NOT APPLY WITHIN 21 DAYS OF HARVEST ON THE FOLLOWING CROPS UNLESS OTHERWISE SPECIFIED

CELERY – [illegible] 1/2 pint per acre. [illegible] whiteflies and leafhoppers, use 1/2 pint per acre. [illegible] days of harvest.

ENDIVE – [illegible] 1/2 pint per acre.

LETTUCE (LEAF AND BIBB) – [illegible] cabbage loopers and potato leafhoppers, use 1/2 pint per acre. [illegible] spotted leafhoppers, use [illegible] per acre. For harlequin bugs and [illegible] worms, use 1/2 pint per acre. [illegible] days of application.

OKRA – To control leaf miners and spider mites, use 1/2 to 3/4 pint per acre. For aphids and stink bugs, use 3/4 pint per acre.

SWISS CHARD – To control aphids and serpentine leaf miners, use 3/4 pint per acre.

DO NOT APPLY WITHIN 15 DAYS OF HARVEST ON THE FOLLOWING CROPS UNLESS OTHERWISE SPECIFIED

BEANS – For control of bean leaf beetles and two-spotted mites, use 3/4 pint per acre. Use 4 to 6 ounces per acre to control thrips and lima pod borers. To control stink bugs, plant bugs, Mexican bean beetles, leaf rollers, leaf miners, leaf hoppers, aphids, red spider mites and armyworms up to third instar, use 3/4 pint per acre.

BEETS – To control flea beetles and leaf miners, use 4-8 ounces per acre. For aphids, blister beetles and webworms, use 1/2 pint per acre. If greens are used for feed, do not use within 21 days of harvest.

BLACKEYED PEAS – To control aphids, leaf miners, bean leaf rollers and stink bugs, use 3/4 pint per acre.

CARROTS – To control leaf miners, use 4 to 6 ounces per acre. To control leafhoppers, use 6 ounces per acre. Use 3/4 pint per acre to control aphids, vegetable weevils, stink bugs and petrobia mites. To control rust fly maggots (first brood), mix [illegible] per acre (2 gallons per acre) and distribute to furrow at planting time. To control rust fly maggots (second brood), use [illegible] per acre as a foliage spray. Do not feed tops.

CUCUMBERS – [illegible] squash vine borers, aphids, cucumber beetles, leaf miners, pickleworms, mites and thrips, use 3/4 to [illegible] squash bugs, stink bugs, flea beetles and leafhoppers, use 3/4 pint per acre. Do not apply prior to vining.

EGGPLANT – To control thrips, leaf miners, blister beetles and flea beetles, use 4-6 ounces per acre. To control Colorado potato beetles, use [illegible] per acre. To control spider mites and lace bugs, use 6 to 8 ounces per acre. To control aphids, [illegible] and stink bugs, use 1/2 pint per acre.

ONIONS – To control thrips, use 1/2 pint per acre. To control onion maggot flies, use 6 ounces per acre. To control aphids, [illegible] and potato leafhoppers, use 3/4 pint per acre. To control brown wheat mites, use 10 ounces per acre.

PEPPERS – To control thrips, use [illegible] 4 ounces per acre. To control aphids, leaf miners and western potato flea beetles, use 1/2 pint per acre.

RADISHES – To control aphids, harlequin bugs and harlequin bugs, use 1/2 to 3/4 pint per acre. To control cabbage loopers and [illegible], use 3/4 pint per acre.

SPINACH – To control aphids, leaf miners, armyworms up to third instar, cabbage loopers, vegetable weevils, harlequin bugs, stink bugs, corn maggots, webworms and leafhoppers, use 3/4 pint per acre.

SQUASH – To control aphids, potato flea beetles, aphids, stink bugs, melonworms, pickleworms, climbing cutworms, serpentine leaf miners and squash vine borers, use 3/4 to 1 pint per acre. To control squash bugs, flea beetles and leafhoppers, use 3/4 pint per acre.

SWEET POTATOES – To control aphids, spider mites, leafhoppers and stink bugs, use 3/4 pint per acre. To control serpentine leaf miners and morning glory leaf miners, use 1/2 to 3/4 pint per acre.

DO NOT APPLY WITHIN 15 DAYS OF HARVEST ON THE FOLLOWING CROPS

SWEET CORN – To control corn borers feeding in the bud, fall armyworms, aphids and silkflies, use 3/4 pint per acre. To control aphids, leafhoppers and spider mites, use 3/4 pint per acre. To control chinch bugs, use 3/4 pint per acre.

DO NOT APPLY WITHIN 10 DAYS OF HARVEST ON THE FOLLOWING CROPS UNLESS OTHERWISE SPECIFIED

PEAS – To control aphids, pea weevils, spider mites, stink bugs, thrips, armyworms up to third instar, climbing cutworms, leaf miners, alfalfa loopers and celery loopers, use 3/4 pint per acre. If vines are to be used for forage, do not harvest for 7 days after treatment.

PUMPKINS – To control aphids, cucumber beetles, climbing cutworms, squash bugs and squash vine borers, use 3/4 to 1 pint per acre.

COLLARDS, KALE AND MUSTARD GREENS – To control aphids, leaf miners, armyworms up to third instar, cabbage loopers, vegetable weevils, harlequin bugs, seed corn maggots, cross striped cutworms and leafhoppers, use 3/4 pint per acre. On fall [illegible] 15 days of harvest.

TOMATOES – To control blister beetles, use 3/4 pint per acre. To control hornworms, leafhoppers and psyllids, use 6 ounces per acre. For aphids, leaf miners, whiteflies, armyworms up to third instar, grasshoppers, mites, tomato russet mites, leaffoot bugs, stink bugs, stink bugs, tarnished plant bugs, use 3/4 pint per acre.

TURNIPS – To control aphids, cabbage webworms, climbing cutworms, vegetable weevils, false chinch bugs and harlequin bugs, use 1/2 pint per acre. To control cabbage loopers, use 1/2 pint per acre. If greens are used for food, do not apply within [illegible] days of harvest.

DO NOT APPLY WITHIN 7 DAYS OF HARVEST ON THE FOLLOWING CROPS UNLESS OTHERWISE SPECIFIED

CABBAGE AND COLE CROPS (BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER) – To control aphids, thrips, flea beetles, [illegible] spotted cabbageworms and armyworms up to third instar, use 3/4 to 1 pint per acre. To control [illegible] vegetable weevils, climbing cutworms and flea beetles, use 3/4 pint per acre. Rates above 3/4 pint [illegible] applied to cabbage [illegible] than 10 days until harvest.

KOHLRABI – To control aphids, use 1/2 pint per acre.

LETTUCE (HEAD) – To control aphids, cabbage loopers, imported cabbageworms, banded cucumber beetles, lygus bugs, webworms and [illegible] use 3/4 to 1 pint per acre. To control six spotted leafhoppers, use 6 ounces per acre. For [illegible] worms and leaf miners, use 1/2 pint per acre. To control garden symphylans, broadcast [illegible] and thoroughly incorporate into upper 6 to 9 inches of soil.

MELONS – To control aphids, leaf miners and false chinch bugs, use 3/4 pint per acre. For aphids, leafhoppers, cucumber beetles, [illegible] use 3/4 to 1 pint per acre. To control thrips, squash bugs and stink bugs, use 1/2 pint per acre.

DO NOT APPLY WITHIN [illegible] DAYS OF HARVEST ON THE FOLLOWING CROP

POTATOES – To control aphids, flea beetles, Colorado potato beetles, leaf miners, mites, plant bugs, potato psyllid, [illegible] leafhoppers, use 3/4 to 1 pint per acre. For armyworms up to third instar, cabbage loopers, [illegible] per acre. For leafhoppers, stink bugs and flea beetles, use 3/4 pint per acre.

NOTICE: Seller warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions not reasonably foreseeable to seller, or [illegible] use.

NET CONTENTS _____ GALS

A PRODUCT OF

TIDE PRODUCTS, INC.

EDINBURG, TEXAS 78539

ONLY the BEST . . . is fit to be TIDE

BEST [illegible] AVAILABLE

GP 881 11767