

BUDMOR 42

EMULSIFIABLE LIQUID AGRICULTURAL INSECTICIDE

ACTIVE INGREDIENTS:

O-O-dimethyl O-p-nitrophenyl thiophosphate 41.88%
 O-Ethyl O-p-nitrophenyl phenylphosphonothioate 20.93%
 Xylene 29.19%

INERT INGREDIENTS:

Methyl Parathion 8.00%

CONTAINS 2 LBS. EPN PLUS 4 LBS. METHYL PARATHION PER GALLON
 CONTAINS DUPONT EPN 100.00%

DANGER-POISON

Keep Out of Reach of Children



WARNING SYMPTOMS: Symptoms include weakness, headache, tightness in chest, blurred vision, non-reactive pin-point pupils, salivation, sweating, nausea, vomiting, diarrhea, and abdominal cramps.

TREATMENT: Atropine is the specific therapeutic antagonist of choice against para-sympathetic nervous stimulation. If there are signs of para-sympathetic stimulation, atropine sulfate should be injected at 10-minute intervals, in doses 1 to 2 milligrams, until complete atropinization has occurred. 2-PAM is also antidotal and may be administered in conjunction with atropine. Morphine is contraindicated. Clear chest by postural drainage. Oxygen administration may be necessary. Observe patient continuously for 48 hours. Repeated exposure to cholinesterase inhibitors may without warning cause prolonged susceptibility to very small doses of any cholinesterase inhibitor. Allow no further exposure until time for cholinesterase regeneration has been allowed as determined by blood test.

Wear gloves and goggles when handling.

In case of contact, wash immediately with soap and water.

Wash hands, arms and face thoroughly with soap and water before eating or smoking.

Wash all contaminated clothing with soap and hot water before reuse.

Repeated inhalation or skin contact may, without symptoms, progressively increase susceptibility to Budmor poisoning.

If you suspect exposure, see a doctor.

Highly toxic to fish and wildlife. Birds and other wildlife in treated areas may be killed. Do not contaminate any body of water or apply to areas or crops other than those for which directions are given on the label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your cooperative Agricultural Extension Service.

DO NOT APPLY WHERE RUNOFF WILL OCCUR. DO NOT CONTAMINATE WATER BY CLEANING OF EQUIPMENT. OR DISPOSAL OF WASTES APPLY THIS PRODUCT ONLY AS SPECIFIED ON THIS LABEL.

Never allow spray mist to drift onto property of others.

The Federal Food and Drug Administration has set of-ten tolerances for certain pesticide residues. In order that residues on food and forage crops will not exceed established tolerances, use only at recommended rates, and do not apply closer to harvest than specified.

Do not contaminate feed or foodstuffs.

Do not splash and do not inhale.

Whenever handling or applying Budmor, wear natural rubber gloves, clean waterproof or freshly laundered protective clothing (coveralls, cap, etc.) gloves and goggles.

Clothing must not get wet through.

Destroy and replace gloves frequently.

Wear a pesticide respirator jointly approved by the Mining Enforcement and Safety Administration (formerly the U.S. Bureau of Mines) and by the National Institute for Occupational Safety and Health under the provisions of 30 CFR Part 11.

Airplane pilots should wear full face canister-type mask.

Do not smoke while spraying or handling Budmor.

Before applying Budmor, remove unprotected persons and domestic animals from operational and adjoining areas where spray may drift.

Work to windward.

For handling indoors, provide mechanical exhaust ventilation.

Burn or destroy containers, never re-use. Stay out of smoke and fumes.

Bury spillage; clean up area with strong lye solution.

Bathe promptly with soap and water after using or handling.

Wash clothing with soap and water before reuse.

Prohibit re-entry until drift and vapor dissipate.

To avoid injury to bees, application of Budmor should not be made while trees or plants are in bloom.

CONTAINER DISPOSAL

1. Empty container into spray tank. Then drain in verti- cle position for 30 seconds.
2. Add a measured amount of rinse water so container is 1/4 to 1/5 full.
3. Rinse container thoroughly, pour into tank, and drain for 30 seconds. Repeat 3 times. Add enough fluid (wa- ter) to bring tank up to level.
4. Crush pesticide container. Sell as scrap for recycling or bury. Do not reuse.

If symptoms or signs of poisoning include blurred vision, abdominal cramps, and tightness in the chest, don't wait for doctor, but give two atropine tablets (each 1/100 gram) at once. Remove patient immediately from the area where Budmor is present. IF SWALLOWED—Induce vomiting by giving warm salt water. Have victim lie down, keep warm and quiet. Never give anything by mouth to an unconscious patient. SKIN CONTACT—Remove contaminated clothing and wash skin clean with plenty of soap and water to remove all traces of Budmor. EYES—Wash immediately with clear water for at least 10 minutes.

A PRODUCT OF

TIDE PRODUCTS, INC.

EDINBURG, TEXAS 78539

ONLY the BEST . . . is fit to be TIDE

Applicatio
 approved
 equipment
 nearly ill
 Keep agri
 tions. D
 hot mix
 do not u
 time with
 tire has
 thorough
 water be

SUGGEST
GROUND
 provide
 growth at
 rain uses
 cover a b

Crop
 Vegetable

COTTON
 worm, C
 Spider M
 water fo
 of Boll
 rints pe
 coverage
 day inte
 hand pic
 cation,
 (100 ap)
 ing.

E.P.A. R
 E.P.A. E

BUDMOR 42

EMULSIFIABLE LIQUID AGRICULTURAL INSECTICIDE

ACTIVE INGREDIENTS:

- *O-O-dimethyl O-p-nitrophenyl thiophosphate
- O-Ethyl O-p-nitrophenyl phenylphosphonothioate
- Xylene

41.88%

20.93%

29.19%

INERT INGREDIENTS:

- *Methyl Parathion

8.00%

100.00%

CONTAINS 2 LBS. EPN PLUS 4 LBS. METHYL PARATHION PER GALLON
CONTAINS DUPONT EPN

DANGER-POISON

Keep Out of Reach of Children



DIRECTIONS FOR USE

Application should be made only by trained personnel using approved protective equipment. Do not apply with hand equipment. Pour specified amount of this product into nearly filled spray tank. Add balance of water to fill tank. Keep agitator running during filling and spraying operations. Do not allow mixture to stand. If mixture does not mix readily, but tends to separate as an oily layer, do not use as injury to plants may result. Do not combine with wettable powders unless previous use of mixture has proven physical compatible and safe. Always thoroughly emulsify this product with at least half of total water before adding wettable powder.

SUGGESTED WATER RATES FOR AIRCRAFT AND GROUND APPLICATION. (The actual rate required to provide thorough, uniform coverage varies with plant growth at time of application. Except as specified for certain uses, the following rates are therefore intended to cover a broad range of conditions.)

Crop	Gal. Water/Acre	
	Aircraft	Ground
Vegetable and Field Crops	1-20	5-100

RECOMMENDATIONS

COTTON: — Cotton Aphid, Thrip, Cotton Leafworm, Cotton Fleahopper and certain species of Spider Mites apply 3/4 pint per acre in sufficient water for thorough coverage. For the control of Boll Weevil and Cotton Bollworm apply 2 pints per acre in sufficient water for thorough coverage. Application should be made at 4-5 day intervals until control is obtained. Do not hand pick or harvest within 7 days of application. Workers entering field within 24 hours after application should wear protective clothing.

SOYBEANS: (Southeast States only) For cotton bollworm control, apply 2 pints per acre in sufficient water to obtain thorough plant coverage. Begin application when counts show one bollworm per 3 feet of soybean row; repeat as necessary. Do not graze or feed forage from treated fields to livestock within 21 days after last application. Do not make more than two applications per growing season.

FOR USE ONLY IN POWER SPRAYERS BY OPERATORS EXPERIENCED OR ADEQUATELY TRAINED IN THE APPLICATION OF PARATHION INSECTICIDES.

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

A PRODUCT OF
TIDE PRODUCTS, INC.
EDINBURG, TEXAS 78539

ONLY the BEST . . . is fit to be TIDE

E.P.A. Registration No. 6735-154
E.P.A. E.S.T. No. 6735-TX-1

GAL. NET

If symptoms or signs of poisoning include blurred vision, abdominal cramps, and tightness in the chest, don't wait for doctor, but give two atropine tablets (each 1/100 grain) at once. Remove patient immediately from the area where Budmor is present. **IF SWALLOWED**—Induce vomiting by giving warm salt water. Have victim lie down, keep warm and quiet. Never give anything by mouth to an unconscious patient. **SKIN CONTACT** — Remove contaminated clothing and wash skin clean with plenty of soap and water to remove all traces of Budmor. **EYES** — Wash immediately with clear water for at least 10 minutes.

CAUTION
DO NOT USE FOR...
3/75