



TIDE COPPER-SULFUR 3.4-70 DUST

AN AGRICULTURAL FUNGICIDE

ACTIVE INGREDIENTS:	
Copper expressed as Metallic (from 53% Basic Copper Sulfate)	3.4%
Sulfur	70.0%
INERT INGREDIENTS	26.6%
Sulfur Fineness 93% passing No. 325 USS Sieve	100.0%

CAUTION

Keep Out of Reach of Children

MAY CAUSE IRRITATION OF NOSE, THROAT AND SKIN.
DO NOT BREATHE DUST.

OBSERVE THE FOLLOWING USE PRECAUTIONS

Sulfur will cause severe fruit and leaf injury to sulfur-sensitive crops. Do not apply to apricots, Anjou and Comice pears, cranberries, cucurbits (cucumber, cantaloupes, melons, squash), walnuts, or other sensitive plants.

Sulfur may burn foliage and fruit during periods of high temperature and under certain other climatic conditions. Do not apply sulfur to sensitive crops when shade temperatures exceed or are likely to exceed 90° F.

Consult State Agricultural Experiment Stations or State Agricultural Extension Service for additional information as the timing, number and rate of applications needed will vary with local conditions.

Apply this product only as specified on the label.

DIRECTIONS FOR USE

Apply the specified amounts of this product uniformly to the foliage when air is calm, using ground equipment or aircraft. Apply at the first sign of infection and repeat at 5- to 10-day intervals, depending on severity of disease, but observe use limits given for specific crops.

When using ground equipment or aircraft apply at the rate of 20 to 40 lb. per acre, depending upon degree of infection and stage of plant development.

RECOMMENDATIONS

BEANS: Powdery Mildew and Rust — Apply as preventive dusting when plants are about 5 inches high. Repeat as necessary. Start with 20 lb. per acre and increase rate to up to 40 pounds to compensate for plant development.

PEANUTS: Peanut Leaf Spot — Apply dust about 60 days after planting, or as soon as spots appear on the leaf.

First Dust — Apply 15 to 20 lb. per acre

Later Dusts — Apply 20 to 30 lb. per acre.

A minimum of 3 to 4 applications should be made at 2-week intervals. Consult your County Agent or Experiment Station for complete details on the peanut-dusting program best suited to your local conditions.

ROSES: Black Spot, Powdery Mildew, Rust — Rose bushes should be dusted before the appearance of diseases on foliage. Apply at rate of 15 to 25 lb. per acre.

Repeat applications at weekly intervals until July (or until hot, dry weather begins); thereafter dust within 24 hours after each rain, but not more than once a week, until growth of plant is checked by cool weather.

BURN EMPTY BAG — KEEP OUT OF SMOKE

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonable fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty or merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

USDA Reg. No. 6735-163

GPS-1-70-1M-(3834) B-48

50 LBS. NET

A PRODUCT OF
TIDE PRODUCTS, INC.
EDINBURG, TEXAS 78539